

G_2 -instantons over twisted connected sums

Henrique Sá Earp

Thomas Walpuski

2015-11-03

Abstract

We introduce a method to construct G_2 -instantons over compact G_2 -manifolds arising as the twisted connected sum of a matching pair of building blocks [Kov03; KL11; CHNP15]. Our construction is based on gluing G_2 -instantons obtained from holomorphic vector bundles over the building blocks via the first named author's work [Sá15]. We require natural compatibility and transversality conditions which can be interpreted in terms of certain Lagrangian subspaces of a moduli space of stable bundles on a $K3$ surface.

Keywords: G_2 -manifolds, G_2 -instantons, twisted connected sum, gauge theory, holomorphic bundles, gluing construction

MSC2010: 53C07, 53C25, 53C38

Changes to the published version This article was first published in *Geom. Topol.* 19 (2015), no. 3, 1263–1285. The present version differs only in aesthetic aspects and Remark 1.10 has been updated to reflect recent progress.

1 Introduction

A G_2 -manifold (Y, g) is a Riemannian 7-manifold whose holonomy group $\text{Hol}(g)$ is contained in the exceptional Lie group G_2 or, equivalently, a 7-manifold Y together with torsion-free G_2 -structure, that is, a non-degenerate 3-form ϕ satisfying a certain non-linear partial differential equation, see, e.g., [Joy96, Part I]. An important method to produce examples of compact G_2 -manifolds with $\text{Hol}(g) = G_2$ is the **twisted connected sum construction**, suggested by Donaldson, pioneered by Kovalev [Kov03] and later extended and improved by Kovalev and Lee [KL11] and Corti, Haskins, Nordström, and Pacini [CHNP15]. Here is a brief summary of this construction: A **building block** consists of a projective 3-fold Z and a smooth anti-canonical $K3$ surface $\Sigma \subset Z$ with trivial normal bundle, see Definition 2.8. Given a choice of hyperkähler structure $(\omega_I, \omega_J, \omega_K)$ on Σ such that $\omega_J + i\omega_K$ is of type $(2, 0)$ and $[\omega_I]$ is the restriction of a Kähler class on Z , one can make $V := Z \setminus \Sigma$

into an asymptotically cylindrical (ACyl) Calabi–Yau 3–fold, that is, a non-compact Calabi–Yau 3–fold with a tubular end modelled on $\mathbf{R}_+ \times S^1 \times \Sigma$, see Haskins–Hein–Nordström [HHN15]. Then $Y := S^1 \times V$ is an ACyl G_2 –manifold with a tubular end modelled on $\mathbf{R}_+ \times T^2 \times \Sigma$.

Definition 1.1. Given a pair of building blocks (Z_\pm, Σ_\pm) , a collection

$$= \{(\omega_{I,\pm}, \omega_{J,\pm}, \omega_{K,\pm}), \mathfrak{r}\}$$

consisting of a choice of hyperkähler structures on Σ_\pm such that $\omega_{J,\pm} + i\omega_{K,\pm}$ is of type $(2, 0)$ and $[\omega_{I,\pm}]$ is the restriction of a Kähler class on Z_\pm as well as a hyperkähler rotation $\mathfrak{r} : \Sigma_+ \rightarrow \Sigma_-$ is called **matching data** and (Z_\pm, Σ_\pm) are said to **match** via \cdot . Here a **hyperkähler rotation** is a diffeomorphism $\mathfrak{r} : \Sigma_+ \rightarrow \Sigma_-$ such that

$$(1.2) \quad \mathfrak{r}^* \omega_{I,-} = \omega_{J,+}, \quad \mathfrak{r}^* \omega_{J,-} = \omega_{I,+} \quad \text{and} \quad \mathfrak{r}^* \omega_{K,-} = -\omega_{K,+}.$$

Given a matching pair of building blocks, one can glue Y_\pm by interchanging the S^1 –factors at infinity and identifying Σ_\pm via \mathfrak{r} . This yields a simply-connected compact 7–manifold Y together with a family of torsion-free G_2 –structures $(\phi_T)_{T \geq T_0}$, see Kovalev [Kov03, Section 4]. From the Riemannian viewpoint (Y, ϕ_T) contains a “long neck” modelled on $[-T, T] \times T^2 \times \Sigma_+$; one can think of the twisted connected sum as reversing the degeneration of the family of G_2 –manifolds that occurs as the neck becomes infinitely long.

If (Z, Σ) is a building block and $\mathcal{E} \rightarrow Z$ is a holomorphic vector bundle such that $\mathcal{E}|_\Sigma$ is stable, then $\mathcal{E}|_\Sigma$ carries a unique ASD instanton compatible with the holomorphic structure [Don85]. The first named author showed that in this situation $\mathcal{E}|_V$ can be given a Hermitian–Yang–Mills (HYM) connection asymptotic to the ASD instanton on $\mathcal{E}|_\Sigma$ [Sá15]. The pullback of a HYM connection over V to $S^1 \times V$ is a G_2 –instanton, i.e., a connection A on a G –bundle over a G_2 –manifold such that $F_A \wedge \psi = 0$ with $\psi := *\phi$. It was pointed out by Simon Donaldson and Richard Thomas in their seminal article on gauge theory in higher dimensions [DT98] that, formally, G_2 –instantons are rather similar to flat connections over 3–manifolds; in particular, they are critical points of a Chern–Simons type functional and there is hope that counting them could lead to an enumerative invariant for G_2 –manifolds not unlike the Casson invariant for 3–manifolds, see [DS11, Section 6] and [Wal13a, Chapter 6]. The main result of this article is the following theorem, which gives conditions for a pair of such G_2 –instantons over $Y_\pm = S^1 \times V_\pm$ to be glued to give a G_2 –instanton over (Y, ϕ_T) .

Theorem 1.3. *Let (Z_\pm, Σ_\pm) be a pair of building blocks that match via \cdot . Denote by Y the compact 7–manifold and by $(\phi_T)_{T \geq T_0}$ the family of torsion-free G_2 –structures obtained from the twisted connected sum construction. Let $\mathcal{E}_\pm \rightarrow Z_\pm$ be a pair of holomorphic vector bundles such that the following hold:*

- $\mathcal{E}_\pm|_{\Sigma_\pm}$ is stable. Denote the corresponding ASD instanton by $A_{\infty,\pm}$.
- There is a bundle isomorphism $\bar{\tau}: \mathcal{E}_+|_{\Sigma_+} \rightarrow \mathcal{E}_-|_{\Sigma_-}$ covering the hyperkähler rotation τ such that $\bar{\tau}^* A_{\infty,-} = A_{\infty,+}$.
- There are no infinitesimal deformations of \mathcal{E}_\pm fixing the restriction to Σ_\pm :

$$(1.4) \quad H^1(Z_\pm, \mathcal{E}nd_0(\mathcal{E}_\pm)(-\Sigma_\pm)) = 0.$$

- Denote by $\text{res}_\pm: H^1(Z_\pm, \mathcal{E}nd_0(\mathcal{E}_\pm)) \rightarrow H^1(\Sigma_\pm, \mathcal{E}nd_0(\mathcal{E}_\pm|_{\Sigma_\pm}))$ the restriction map and by $\lambda_\pm: H^1(Z_\pm, \mathcal{E}nd_0(\mathcal{E}_\pm)) \rightarrow H^1_{A_{\infty,\pm}}$ the composition of res_\pm with the isomorphism from Remark 1.6. The images of λ_+ and $\bar{\tau}^* \circ \lambda_-$ intersect trivially in $H^1_{A_{\infty,+}}$:

$$(1.5) \quad \text{im}(\lambda_+) \cap \text{im}(\bar{\tau}^* \circ \lambda_-) = \{0\}.$$

Then there exists a non-trivial $\text{PU}(n)$ -bundle E over Y , a constant $T_1 \geq T_0$ and for each $T \geq T_1$ an irreducible and unobstructed¹ G_2 -instanton A_T on E over (Y, ϕ_T) .

Remark 1.6. If A is an ASD instanton on a $\text{PU}(n)$ -bundle E over a Kähler surface Σ corresponding to a holomorphic vector bundle \mathcal{E} , then

$$H_A^1 := \ker(d_A^* \oplus d_A^+: \Omega^1(\Sigma, \mathfrak{g}_E) \rightarrow (\Omega^0 \oplus \Omega^+)(\Sigma, \mathfrak{g}_E)) \cong H^1(\Sigma, \mathcal{E}nd_0(\mathcal{E})),$$

see Donaldson and Kronheimer [DK90, Section 6.4]. Here \mathfrak{g}_E denotes the adjoint bundle associated with E .

Remark 1.7. If

$$(1.8) \quad H^1(\Sigma_+, \mathcal{E}nd_0(\mathcal{E}_+|_{\Sigma_+})) = \{0\},$$

then (1.5) is vacuous. If, moreover, the topological bundles underlying \mathcal{E}_\pm are isomorphic, then the existence of $\bar{\tau}$ is guaranteed by a theorem of Mukai [HL97, Theorem 6.1.6].

Since $H^2(Z_\pm, \mathcal{E}nd_0(\mathcal{E}_\pm)) \cong H^1(Z_\pm, \mathcal{E}nd_0(\mathcal{E}_\pm)(-\Sigma_\pm))$ vanish by (1.4), there is a short exact sequence

$$0 \rightarrow H^1(Z_\pm, \mathcal{E}nd_0(\mathcal{E}_\pm)) \xrightarrow{\text{res}_\pm} H^1(\Sigma_\pm, \mathcal{E}nd_0(\mathcal{E}_\pm|_{\Sigma_\pm})) \rightarrow H^2(Z_\pm, \mathcal{E}nd_0(\mathcal{E}_\pm)(-\Sigma_\pm)) \rightarrow 0.$$

This sequence is self-dual under Serre duality. It was pointed out by Tyurin [Tyu08, p. 176 ff.] that this implies that

$$\text{im} \lambda_\pm \subset H^1_{A_{\infty,\pm}}$$

¹See Definition 3.12.

is a complex Lagrangian subspace with respect to the complex symplectic structure induced by $\Omega_{\pm} := \omega_{J,\pm} + i\omega_{K,\pm}$ or, equivalently, Mukai's complex symplectic structure on $H^1(Z_{\pm}, \mathcal{E}nd_0(\mathcal{E}_{\pm}))$. Under the assumptions of Theorem 1.3 the moduli space $\mathcal{M}(\Sigma_+)$ of holomorphic vector bundles over Σ_+ is smooth near $[\mathcal{E}_+|_{\Sigma_+}]$ and so are the moduli spaces $\mathcal{M}(Z_{\pm})$ of holomorphic vector bundles over Z_{\pm} near $[\mathcal{E}_{\pm}]$. Locally, $\mathcal{M}(Z_{\pm})$ embeds as a complex Lagrangian submanifold into $\mathcal{M}(\Sigma_{\pm})$. Since $r^*\omega_{K,-} = -\omega_{K,+}$, both $\mathcal{M}(Z_+)$ and $\mathcal{M}(Z_-)$ can be viewed as Lagrangian submanifolds of $\mathcal{M}(\Sigma_+)$ with respect to the symplectic form induced by $\omega_{K,+}$. Equation (1.5) asks for these Lagrangian submanifolds to intersect transversely at the point $[\mathcal{E}_+|_{\Sigma_+}]$. If one thinks of G_2 -manifolds arising via the twisted connected sum construction as analogues of 3-manifolds with a fixed Heegaard splitting, then this is much like the geometric picture behind Atiyah–Floer conjecture in dimension three [Ati88].

Remark 1.9. The hypothesis (1.5) appears natural in view of the above discussion. Assuming (1.8) instead would slightly simplify the proof, see Remark 3.38; however, it would also substantially restrict the applicability of Theorem 1.3 and, hence, the chance of finding new examples of G_2 -instantons because (1.8) is a very strong assumption.

Remark 1.10. Using Theorem 1.3 in a situation with (1.8), the first example of an irreducible and unobstructed G_2 -instanton over a twisted connected sum has been constructed by the second named author in [Wal15]. Recent joint work by Grégoire Menet, Johannes Nordström and the first named author [MNS15] constructs a further example of an irreducible and unobstructed G_2 -instanton using Theorem 1.3 in a situation where (1.8) fails.

Outline We recall the salient features of the twisted connected sum construction in Section 2. The expert reader may wish to skim through it to familiarise with our notation. The objective of Section 3 is to prove Theorem 3.24, which describes hypotheses under which a pair of G_2 -instantons over a matching pair of ACyl G_2 -manifolds can be glued. Finally, in Section 4 we explain how these hypotheses can be verified for G_2 -instantons obtained via the first named author's construction. Theorem 1.3 is then proved by combining Theorem 3.24 and Theorem 4.2 with Proposition 4.3.

Acknowledgements. We are grateful to Simon Donaldson for suggesting the problem solved in this article. We thank Marcos Jardim. Moreover, we thank the anonymous referee for many helpful comments and suggestions. TW was supported by ERC Grant 247331 and Unicamp-Faepex grant 770/13.

2 The twisted connected sum construction

In this section we review the twisted connected sum construction using the language introduced by Corti–Haskins–Nordström–Pacini [CHNP15].

2.1 Gluing ACyl G_2 -manifolds

We begin with gluing matching pairs of ACyl G_2 -manifolds.

Definition 2.1. Let (Z, ω, Ω) be a compact Calabi–Yau 3-fold. Here ω denotes the Kähler form and Ω denotes the holomorphic volume form. A G_2 -manifold (Y, ϕ) is called **asymptotically cylindrical (ACyl)** with asymptotic cross-section (Z, ω, Ω) if there exist a constant $\delta < 0$, a compact subset $K \subset Y$, a diffeomorphism $\pi: Y \setminus K \rightarrow \mathbf{R}_+ \times Z$ and a 2-form ρ on $\mathbf{R}_+ \times Z$ such that

$$\pi_*\phi = dt \wedge \omega + \operatorname{Re} \Omega + d\rho$$

and

$$\nabla^k \rho = O(e^{\delta t})$$

for all $k \in \mathbf{N}_0$. Here t denotes the coordinate on \mathbf{R}_+ .

Remark 2.2. Unfortunately, Z is the customary notation both for building blocks and asymptotic cross-sections of ACyl G_2 -manifolds. To avoid confusion we point out that, unlike asymptotic cross-sections, building blocks always come in pair with a divisor, e.g., (Z, Σ) .

Definition 2.3. A pair of ACyl G_2 -manifolds (Y_{\pm}, ϕ_{\pm}) with asymptotic cross-sections $(Z_{\pm}, \omega_{\pm}, \Omega_{\pm})$ is said to **match** if there exists a diffeomorphism $f: Z_+ \rightarrow Z_-$ such that

$$f^*\omega_- = -\omega_+ \quad \text{and} \quad f^*\operatorname{Re} \Omega_- = \operatorname{Re} \Omega_+.$$

Let (Y_{\pm}, ϕ_{\pm}) be a matching pair of ACyl G_2 -manifolds. Fix $T \geq 1$. Define $F: [T, T+1] \times Z_+ \rightarrow [T, T+1] \times Z_-$ by

$$F(t, z) := (2T+1-t, f(z)).$$

Denote by Y_T the compact 7-manifold obtained by gluing together

$$Y_{T,\pm} := K_{\pm} \cup \pi_{\pm}^{-1}([0, T+1] \times Z_{\pm})$$

via F . Fix a non-decreasing smooth function $\chi: \mathbf{R} \rightarrow [0, 1]$ with $\chi(t) = 0$ for $t \leq 0$ and $\chi(t) = 1$ for $t \geq 1$. Define a 3-form $\tilde{\phi}_T$ on Y_T by

$$\tilde{\phi}_T := \phi_{\pm} - d[\pi_{\pm}^*(\chi(t-T+1)\rho_{\pm})]$$

on $Y_{T,\pm}$. If $T \gg 1$, then $\tilde{\phi}_T$ defines a closed G_2 -structure on Y_T . Clearly, all the Y_T for different values of T are diffeomorphic; hence, we often drop the T from the notation. The G_2 -structure $\tilde{\phi}_T$ is not torsion-free yet, but can be made so by a small perturbation:

Theorem 2.4 (Kovalev [Kov03, Theorem 5.34]). *In the above situation there exist a constant $T_0 \geq 1$ and for each $T \geq T_0$ there exists a 2-form η_T on Y_T such that $\phi_T := \tilde{\phi}_T + d\eta_T$ defines a torsion-free G_2 -structure; moreover, for some $\delta < 0$*

$$(2.5) \quad \|d\eta_T\|_{C^{0,\alpha}} = O(e^{\delta T}).$$

2.2 ACyl Calabi–Yau 3-folds from building blocks

The twisted connected sum is based on gluing ACyl G_2 -manifolds arising as the product of ACyl Calabi–Yau 3-folds with S^1 .

Definition 2.6. Let $(\Sigma, \omega_I, \omega_J, \omega_K)$ be a hyperkähler surface. A Calabi–Yau 3-fold (V, ω, Ω) is called **asymptotically cylindrical (ACyl)** with asymptotic cross-section $(\Sigma, \omega_I, \omega_J, \omega_K)$ if there exist a constant $\delta < 0$, a compact subset $K \subset V$, a diffeomorphism $\pi: V \setminus K \rightarrow \mathbf{R}_+ \times S^1 \times \Sigma$, a 1-form ρ and a 2-form σ on $\mathbf{R}_+ \times S^1 \times \Sigma$ such that

$$\begin{aligned} \pi_*\omega &= dt \wedge d\alpha + \omega_I + d\rho, \\ \pi_*\Omega &= (d\alpha - idt) \wedge (\omega_J + i\omega_K) + d\sigma \end{aligned}$$

and

$$\nabla^k \rho = O(e^{\delta t}) \quad \text{as well as} \quad \nabla^k \sigma = O(e^{\delta t})$$

for all $k \in \mathbf{N}_0$. Here t and α denote the respective coordinates on \mathbf{R}_+ and S^1 .

Given an ACyl Calabi–Yau 3-fold (V, ω, Ω) , taking the product with S^1 , with coordinate β , yields an ACyl G_2 -manifold

$$(Y := S^1 \times V, \phi := d\beta \wedge \omega + \text{Re } \Omega)$$

with asymptotic cross-section

$$(T^2 \times \Sigma, d\alpha \wedge d\beta + \omega_K, (d\alpha - id\beta) \wedge (\omega_J + i\omega_I)).$$

Let V_\pm be a pair of ACyl Calabi–Yau 3-folds with asymptotic cross-section Σ_\pm and suppose that $r: \Sigma_+ \rightarrow \Sigma_-$ is a hyperkähler rotation, see (1.2). Then $Y_\pm := V_\pm \times S^1$ match via the diffeomorphism $f: T^2 \times \Sigma_+ \rightarrow T^2 \times \Sigma_-$ defined by

$$f(\alpha, \beta, x) := (\beta, \alpha, r(x)).$$

Remark 2.7. If f did not interchange the S^1 -factors, then Y would have infinite fundamental group and, hence, could not carry a metric with holonomy equal to G_2 [Joy00, Proposition 10.2.2].

ACyl Calabi–Yau 3-folds can be obtained from the following building blocks:

Definition 2.8 (Corti, Haskins, Nordström, and Pacini [CHNP13, Definition 5.1]). A **building block** is a smooth projective 3-fold Z together with a projective morphism $f: Z \rightarrow \mathbf{P}^1$ such that the following hold:

- The anticanonical class $-K_Z \in H^2(Z)$ is primitive.
- $\Sigma := f^{-1}(\infty)$ is a smooth $K3$ surface and $\Sigma \sim -K_Z$.
- If N denotes the image of $H^2(Z)$ in $H^2(\Sigma)$, then the embedding $N \hookrightarrow H^2(\Sigma)$ is primitive.
- $H^3(Z)$ is torsion-free.

Remark 2.9. The existence of the fibration $f: Z \rightarrow \mathbf{P}^1$ is equivalent to Σ having trivial normal bundle. This is crucial because it means that $Z \setminus \Sigma$ has a cylindrical end. The last two conditions in the definition of a building block are not essential; they have been made to facilitate the computation of certain topological invariants in [CHNP13].

In his original work Kovalev [Kov03] used building blocks arising from Fano 3-folds by blowing-up the base-locus of a generic anti-canonical pencil. This method was extended to the much larger class of semi Fano 3-folds (a class of weak Fano 3-folds) by Corti, Haskins, Nordström, and Pacini [CHNP15]. Kovalev and Lee [KL11] construct building blocks starting from $K3$ surfaces with non-symplectic involutions, by taking the product with \mathbf{P}^1 , dividing by \mathbf{Z}_2 and blowing up the resulting singularities.

Theorem 2.10 (Haskins, Hein, and Nordström [HHN15, Theorem D]). *Let (Z, Σ) be a building block and let $(\omega_I, \omega_J, \omega_K)$ be a hyperkähler structure on Σ such that $\omega_J + i\omega_K$ is of type $(2, 0)$. If $[\omega_I] \in H^{1,1}(\Sigma)$ is the restriction of a Kähler class on Z , then there is an asymptotically cylindrical Calabi–Yau structure (ω, Ω) on $V := Z \setminus \Sigma$ with asymptotic cross-section $(\Sigma, \omega_I, \omega_J, \omega_K)$.*

Remark 2.11. This result was first claimed by Kovalev [Kov03, Theorem 2.4]; see the discussion in [HHN15, Section 4.1].

Combining the results of Kovalev and Haskins–Hein–Nordström, each matching pair of building blocks (see Definition 1.1) yields a one-parameter family of G_2 -manifolds. This is called the **twisted connected sum construction**.




Figure 1: The twisted connected sum of a matching pair of building blocks.

3 Gluing G_2 -instantons over ACyl G_2 -manifolds

In this section we discuss when a pair of G_2 -instantons over a matching pair of ACyl G_2 -manifolds Y_{\pm} can be glued to give a G_2 -instanton over (Y, ϕ_T) .

3.1 Linear analysis on ACyl manifolds

We recall some results about linear analysis on ACyl Riemannian manifolds. The references for the material in this subsection are Maz'ya and Plamenevskii [MP78] and Lockhart and McOwen [LM85].

3.1.1 Translation-invariant operators on cylindrical manifolds

Let $E \rightarrow X$ be a Riemannian vector bundle over a compact Riemannian manifold. By slight abuse of notation we also denote by E its pullback to $\mathbf{R} \times X$. Denote by t the coordinate function on \mathbf{R} . For $k \in \mathbf{N}_0$, $\alpha \in (0, 1)$ and $\delta \in \mathbf{R}$ we define

$$\|\cdot\|_{C_{\delta}^{k,\alpha}} := \|e^{-\delta t} \cdot\|_{C^{k,\alpha}}$$

and denote by $C_{\delta}^{k,\alpha}(\mathbf{R} \times X, E)$ the closure of $C_0^{\infty}(\mathbf{R} \times X, E)$ with respect to this norm.

We set $C_{\delta}^{\infty} := \bigcap_k C_{\delta}^{k,\alpha}$.

Let $D: C^{\infty}(X, E) \rightarrow C^{\infty}(X, E)$ be a linear self-adjoint elliptic operator of first order. The operator

$$L_{\infty} := \partial_t - D$$

extends to a bounded linear operator $L_{\infty, \delta}: C_{\delta}^{k+1,\alpha}(\mathbf{R} \times X, E) \rightarrow C_{\delta}^{k,\alpha}(\mathbf{R} \times X, E)$.

Theorem 3.1 (Maz'ya and Plamenevskii [MP78, Theorem 5.1]). *The linear operator $L_{\infty, \delta}$ is invertible if and only if $\delta \notin \text{spec}(D)$.*

Elements $a \in \ker L_{\infty}$ can be expanded as

$$(3.2) \quad a = \sum_{\delta \in \text{spec } D} e^{\delta t} a_{\delta}$$

where a_δ are δ -eigensections of D , see [Dono2, Section 3.1]. One consequence of this is the following result:

Proposition 3.3. *Denote by λ_+ and λ_- the first positive and negative eigenvalue of D , respectively. If $a \in \ker L_\infty$ and*

$$a = O(e^{\delta t}) \text{ as } t \rightarrow \infty$$

with $\delta < \lambda_+$, then there exists $a_0 \in \ker D$ such that

$$\nabla^k(a - a_0) = O(e^{\lambda_- t}) \text{ as } t \rightarrow \infty$$

for all $k \in \mathbf{N}_0$. If $a \in L^\infty(\mathbf{R} \times X, E)$, then $a = a_0$.

3.1.2 Asymptotically translation-invariant operators on ACyl manifolds

Let M be a Riemannian manifold together with a compact set $K \subset M$ and a diffeomorphism $\pi: M \setminus K \rightarrow \mathbf{R}_+ \times X$ such that the push-forward of the metric on M is asymptotic to the metric on $\mathbf{R}_+ \times X$, this means here and in what follows that their difference and all of its derivatives are $O(e^{\delta t})$ as $t \rightarrow \infty$ with $\delta < 0$. Let F be a Riemannian vector bundle and let $\tilde{\pi}: F|_{M \setminus K} \rightarrow E$ be a bundle isomorphism covering π such that the push-forward of the metric on F is asymptotic to the metric on E . Denote by $t: M \rightarrow [1, \infty)$ a smooth positive function which agrees with $t \circ \pi$ on $\pi^{-1}([1, \infty) \times X)$. We define

$$\|\cdot\|_{C_\delta^{k,\alpha}} := \|e^{-\delta t} \cdot\|_{C^{k,\alpha}}$$

and denote by $C_\delta^{k,\alpha}(M, F)$ the closure of $C_0^\infty(M, F)$ with respect to this norm.

Let $L: C_0^\infty(M, E) \rightarrow C_0^\infty(M, E)$ be an elliptic operator asymptotic to $L_\infty = \partial_t - D$, that is, the coefficients of the push-forward of L to $\mathbf{R}_+ \times X$ are asymptotic to the coefficients of L_∞ . The operator L extends to a bounded linear operator $L_\delta: C_\delta^{k+1,\alpha}(M, E) \rightarrow C_\delta^{k,\alpha}(M, E)$.

Proposition 3.4 ([HHN15, Proposition 2.4]). *If $\delta \notin \text{spec}(D)$, then L_δ is Fredholm.*

Elements in the kernel of L still have an asymptotic expansion analogous to (3.2). We need the following result which extracts the constant term of this expansion.

Proposition 3.5. *There is a constant $\delta_0 > 0$ such that, for all $\delta \in [0, \delta_0]$, $\ker L_\delta = \ker L_0$ and there is a linear map $\iota: \ker L_0 \rightarrow \ker D$ such that*

$$\nabla^k(\tilde{\pi}_* a - \iota(a)) = O(e^{-\delta_0 t}) \text{ as } t \rightarrow \infty$$

for all $k \in \mathbf{N}_0$; in particular,

$$\ker \iota = \ker L_{-\delta_0}.$$

Proof. Let λ_{\pm} be the first positive/negative eigenvalue of D . Pick $0 < \delta_0 < \min(\lambda_+, -\lambda_-)$ such that the decay conditions made above hold with $-2\delta_0$ instead of δ . Given $a \in \ker L_{\delta_0}$, set $\tilde{a} := \chi(t)\tilde{\pi}_*a_{\pm}$ with χ as in Definition 2.1. Then $L_{\infty}\tilde{a} \in C_{-\delta_0}^{\infty}$. By Theorem 3.1 there exists a unique $b \in C_{-\delta_0}^{\infty}$ such that $L_{\infty}(\tilde{a} - b) = 0$. By Proposition 3.3 $(\tilde{a} - b)_0 \in \ker D$ and $\tilde{a} - b - (\tilde{a} - b)_0 = O(e^{\lambda_- t})$ as t tends to infinity. From this it follows that $a \in \ker L_0$; hence, the first part of the proposition. With $\iota(a) := (\tilde{a} - b)_0$ the second part also follows. \square

3.2 Hermitian–Yang–Mills connections over Calabi–Yau 3–folds

Suppose (Z, ω, Ω) is Calabi–Yau 3–fold and $(Y := \mathbf{R} \times Z, \phi := dt \wedge \omega + \operatorname{Re} \Omega)$ is the corresponding cylindrical G_2 –manifold. In this section we relate translation-invariant G_2 –instantons over Y with Hermitian–Yang–Mills connections over Z . Let G denote a compact semi-simple Lie group.

Definition 3.6. Let (Z, ω) be a Kähler manifold and let E be a G –bundle over Z . A connection A on E is **Hermitian–Yang–Mills (HYM) connection** if

$$(3.7) \quad F_A^{0,2} = 0 \quad \text{and} \quad \Lambda F_A = 0.$$

Here Λ is the dual of the Lefschetz operator $L := \omega \wedge \cdot$.

Remark 3.8. We are mostly interested in the special case of $U(n)$ –bundles; however, for $G = U(n)$, (3.7) is too restrictive as it forces $c_1(E) = 0$. There are two customary ways to circumnavigate this issue: One is to change (3.7) and instead of the second part require that ΛF_A be equal to a constant in $\mathfrak{u}(1)$, the centre of $\mathfrak{u}(n)$, which is determined by the degree of $\det E$; the other one is to work with the induced $PU(n)$ –bundle. These view points are essentially equivalent and we adopt the latter.

Remark 3.9. By the first part of (3.7) a HYM connection induces a holomorphic structure on E . If Z is compact, then there is a one-to-one correspondence between gauge equivalence classes of HYM connections on E and isomorphism classes of polystable holomorphic $G^{\mathbb{C}}$ –bundles \mathcal{E} whose underlying topological bundle is E , see Donaldson [Don85] and Uhlenbeck–Yau [UY86].

On a Calabi–Yau 3–fold (3.7) is equivalent to

$$F_A \wedge \operatorname{Im} \Omega = 0 \quad \text{and} \quad F_A \wedge \omega \wedge \omega = 0;$$

hence, using $\psi = *\phi = *(dt \wedge \omega + \operatorname{Re} \Omega) = \frac{1}{2}\omega \wedge \omega - dt \wedge \operatorname{Im} \Omega$ one easily derives:

Proposition 3.10 ([Sá 15, Proposition 8]). *Denote by $\pi_Z: Y \rightarrow Z$ the canonical projection. A is a HYM connection if and only if π_Z^*A is a G_2 –instanton.*

In general, if A is a G_2 -instanton on a G -bundle E over a G_2 -manifold (Y, ϕ) , then the moduli space \mathcal{M} of G_2 -instantons near $[A]$, i.e., the space of gauge equivalence classes of G_2 -instantons near $[A]$ is the space of small solutions $(\xi, a) \in (\Omega^0 \oplus \Omega^1)(Y, \mathfrak{g}_E)$ of the system of equations

$$d_A^* a = 0 \quad \text{and} \quad d_{A+a} \xi + *(F_{A+a} \wedge \psi) = 0$$

modulo the action of $\Gamma_A \subset \mathcal{G}$, the stabiliser of A in the gauge group of E —assuming Y is compact or appropriate assumptions are made regarding the growth of ξ and a . The linearisation $L_A: (\Omega^0 \oplus \Omega^1)(Y, \mathfrak{g}_E) \rightarrow (\Omega^0 \oplus \Omega^1)(Y, \mathfrak{g}_E)$ of this equation is

$$(3.11) \quad L_A := \begin{pmatrix} & d_A^* \\ d_A & *(\psi \wedge d_A) \end{pmatrix}.$$

It controls the infinitesimal deformation theory of A .

Definition 3.12. A is called **irreducible and unobstructed** if L_A is surjective.

If A is irreducible and unobstructed, then \mathcal{M} is smooth at $[A]$. If Y is compact, then L_A has index zero; hence, is surjective if and only if it is invertible; therefore, irreducible and unobstructed G_2 -instantons form isolated points in \mathcal{M} . If Y is non-compact, the precise meaning of \mathcal{M} and L_A depends on the growth assumptions made on ξ and a and \mathcal{M} may very well be positive-dimensional.

Proposition 3.13. *If A is HYM connection on a bundle E over a G_2 -manifold $Y := \mathbf{R} \times Z$ as in Proposition 3.10, then the operator $L_{\pi_Z^* A}$ defined in (3.11) can be written as*

$$L_{\pi_Z^* A} = \tilde{I} \partial_t + D_A$$

where

$$\tilde{I} := \begin{pmatrix} & -1 \\ 1 & I \end{pmatrix}$$

and $D_A: (\Omega^0 \oplus \Omega^0 \oplus \Omega^1)(Z, \mathfrak{g}_E) \rightarrow (\Omega^0 \oplus \Omega^0 \oplus \Omega^1)(Z, \mathfrak{g}_E)$ is defined by

$$(3.14) \quad D_A := \begin{pmatrix} & d_A^* & \\ d_A & -Id_A & -*(\text{Im } \Omega \wedge d_A) \end{pmatrix}.$$

(Note that $TY = \underline{\mathbf{R}} \oplus \pi_Z^* TZ$.)

Proof. Plugging $\psi = \frac{1}{2} \omega \wedge \omega - dt \wedge \text{Im } \Omega$ into the definition of $L_{\pi_Z^* A}$ and using the fact that the complex structure acts via

$$(3.15) \quad I = \frac{1}{2} * (\omega \wedge \omega \wedge \cdot)$$

on $\Omega^1(Z, \mathfrak{g}_E)$ the assertion follows by a direct computation. \square

Definition 3.16. Let A be a HYM connection on a G -bundle E over a Kähler manifold (Z, ω) . Set

$$\mathcal{H}_A^i := \ker \left(\bar{\partial}_A \oplus \bar{\partial}_A^* : \Omega^{0,i} \left(Z, \mathfrak{g}_E^{\mathbb{C}} \right) \rightarrow \left(\Omega^{0,i+1} \oplus \Omega^{0,i-1} \right) \left(Z, \mathfrak{g}_E^{\mathbb{C}} \right) \right).$$

\mathcal{H}_A^0 is called the space of **infinitesimal automorphisms** of A and \mathcal{H}_A^1 is called the space of **infinitesimal deformations** of A .

Remark 3.17. If Z is compact and A is a connection on a $\mathrm{PU}(n)$ -bundle E corresponding to a holomorphic vector bundle \mathcal{E} , then $\mathcal{H}_A^i \cong H^i(Z, \mathcal{E} \mathrm{nd}_0(\mathcal{E}))$.

Proposition 3.18. *If (Z, ω, Ω) is a compact Calabi–Yau 3-fold and A is a HYM connection on a G -bundle $E \rightarrow Z$, then*

$$\ker D_A \cong \mathcal{H}_A^0 \oplus \mathcal{H}_A^1$$

where D_A is as in (3.14).

Proof. If $s \in \mathcal{H}_A^0$ and $\alpha \in \mathcal{H}_A^1$, then $D_A(\mathrm{Re} s, \mathrm{Im} s, \alpha + \bar{\alpha}) = 0$. Conversely, if $(\xi, \eta, a) \in \ker D_A$, then applying d_A^* (resp. $d_A^* \circ I$) to

$$d_A \xi - \mathrm{Id}_A \eta - *(\mathrm{Im} \Omega \wedge d_A a) = 0,$$

using (3.15), taking the L^2 inner product with ξ (resp. η) and integrating by parts yields $d_A \xi = 0$ (resp. $d_A \eta = 0$). Thus $\xi + i\eta \in \mathcal{H}_A^0$ and

$$d_A^* a = 0, \quad \Lambda d_A a = 0 \quad \text{and} \quad \mathrm{Im} \Omega \wedge d_A a = 0$$

which implies $a := a^{0,1} \in \mathcal{H}_A^1$ because $d_A^* = \partial_A^* + \bar{\partial}_A^*$ and $\Lambda d_A = -i\partial_A^* + i\bar{\partial}_A^*$. \square

3.3 G_2 -instantons over ACyl G_2 -manifolds

Definition 3.19. Let (Y, ϕ) be an ACyl G_2 -manifold with asymptotic cross-section (Z, ω, Ω) . Let A_∞ be a HYM connection on a G -bundle $E_\infty \rightarrow Z$. A G_2 -instanton A on a G -bundle $E \rightarrow Y$ is called **asymptotic** to A_∞ if there exist a constant $\delta < 0$ and a bundle isomorphism $\bar{\pi} : E|_{Y \setminus K} \rightarrow E_\infty$ covering $\pi : Y \setminus K \rightarrow \mathbf{R}_+ \times Z$ such that

$$(3.20) \quad \nabla^k (\bar{\pi}_* A - A_\infty) = O(e^{\delta t})$$

for all $k \in \mathbf{N}_0$. Here by a slight abuse of notation we also denote by E_∞ and A_∞ their respective pullbacks to $\mathbf{R}_+ \times Z$.

Definition 3.21. Let (Y, ϕ) be an ACyl G_2 -manifold and let A be a G_2 -instanton on a G -bundle over (Y, ϕ) asymptotic to A_∞ . For $\delta \in \mathbf{R}$ we set

$$\mathcal{T}_{A,\delta} := \ker L_{A,\delta} = \left\{ \underline{a} \in \ker L_A : \nabla^k \bar{\pi}_* \underline{a} = O(e^{\delta t}) \text{ for all } k \in \mathbf{N}_0 \right\}.$$

where $\underline{a} = (\xi, a) \in (\Omega^0 \oplus \Omega^1)(Y, \mathfrak{g}_E)$. Set $\mathcal{T}_A := \mathcal{T}_{A,0}$.

Proposition 3.22. *Let (Y, ϕ) be an ACyl G_2 -manifold and let A be a G_2 -instanton asymptotic to A_∞ . Then there is a constant $\delta_0 > 0$ such that for all $\delta \in [0, \delta_0]$, $\mathcal{T}_{A, \delta} = \mathcal{T}_A$ and there is a linear map $\iota: \mathcal{T}_A \rightarrow \mathcal{H}_{A_\infty}^0 \oplus \mathcal{H}_{A_\infty}^1$ such that*

$$\nabla^k (\tilde{\pi}_* \underline{a} - \iota(\underline{a})) = O(e^{-\delta_0 t})$$

for all $k \in \mathbb{N}_0$; in particular,

$$\ker \iota = \mathcal{T}_{A, -\delta_0}.$$

Proof. By Proposition 3.13, L_A is asymptotic to $\tilde{I}(\partial_t - \tilde{I}D_A)$. Since $\tilde{I}D_A$ is self-adjoint and $\ker \tilde{I}D_A = \ker D_A$, we can apply Proposition 3.5 to obtain a linear map $\iota: \mathcal{T}_A \rightarrow \ker D_{A_\infty}$ and use the isomorphism $\ker D_{A_\infty} \cong \mathcal{H}_{A_\infty}^0 \oplus \mathcal{H}_{A_\infty}^1$ from Proposition 3.18. \square

Proposition 3.23. *Let (Y, ϕ) be an ACyl G_2 -manifold and let A be a G_2 -instanton asymptotic to A_∞ . Then*

$$\dim \operatorname{im} \iota = \frac{1}{2} \dim \left(\mathcal{H}_{A_\infty}^0 \oplus \mathcal{H}_{A_\infty}^1 \right)$$

and, if $\mathcal{H}_{A_\infty}^0 = 0$, then $\operatorname{im} \iota \subset \mathcal{H}_{A_\infty}^1$ is Lagrangian with respect to the symplectic structure on $\mathcal{H}_{A_\infty}^1$ induced by ω .

Proof. By Lockhart and McOwen [LM85, Theorem 7.4] for $0 < \delta \ll 1$

$$\dim \operatorname{im} \iota = \operatorname{index} L_{A, \delta} = \frac{1}{2} \dim \ker D_{A_\infty}.$$

Suppose $\mathcal{H}_{A_\infty}^0 = 0$. If $(\xi, a) \in \mathcal{T}_A$, then $d_A^* d_A \xi = 0$ and, by Proposition 3.22, ξ decays exponentially. Integration by parts shows that $d_A \xi = 0$; hence, $\xi = 0$. Therefore, $\mathcal{T}_A \subset \Omega^1(Y, \mathfrak{g}_E)$

We show that $\operatorname{im} \iota$ is isotropic: For $a, b \in \mathcal{T}_A$

$$\frac{1}{2} \int_Z \langle \iota(a) \wedge \iota(b) \rangle \wedge \omega \wedge \omega = \int_Y d(\langle a \wedge b \rangle \wedge \psi) = 0$$

because $d_A a \wedge \psi = d_A b \wedge \psi = 0$. \square

3.4 Gluing G_2 -instantons over ACyl G_2 -manifolds

In the situation of Proposition 3.23, if $\ker \iota = 0$ and $\mathcal{H}_{A_\infty}^0 = 0$, then one can show that the moduli space $\mathcal{M}(Y)$ of G_2 -instantons near $[A]$ which are asymptotic to some HYM connection is smooth. Although the moduli space $\mathcal{M}(Z)$ of HYM connections near $[A_\infty]$ is not necessarily smooth, formally, it still makes sense to talk about its symplectic structure and view $\mathcal{M}(Y)$ as a Lagrangian submanifold. The following theorem shows, in particular, that transverse intersections of a pair of such Lagrangians give rise to G_2 -instantons.

Theorem 3.24. *Let (Y_{\pm}, ϕ_{\pm}) be a pair of ACyl G_2 -manifolds that match via $f: Z_+ \rightarrow Z_-$. Denote by $(Y_T, \phi_T)_{T \geq T_0}$ the resulting family of compact G_2 -manifolds arising from the construction in Section 2.1. Let A_{\pm} be a pair of G_2 -instantons on E_{\pm} over (Y_{\pm}, ϕ_{\pm}) asymptotic to $A_{\infty, \pm}$. Suppose that the following hold:*

- *There is a bundle isomorphism $\tilde{f}: E_{\infty, +} \rightarrow E_{\infty, -}$ covering f such that $\tilde{f}^* A_{\infty, -} = A_{\infty, +}$,*
- *The maps $\iota_{\pm}: \mathcal{T}_{A_{\pm}} \rightarrow \ker D_{A_{\infty, \pm}}$ constructed in Proposition 3.22 are injective and their images intersect trivially*

$$(3.25) \quad \text{im}(\iota_+) \cap \text{im}(\tilde{f}^* \circ \iota_-) = \{0\} \subset \mathcal{H}_{A_{\infty, +}}^0 \oplus \mathcal{H}_{A_{\infty, +}}^1.$$

Then there exists $T_1 \geq T_0$ and for each $T \geq T_1$ there exists an irreducible and unobstructed G_2 -instanton A_T on a G -bundle E_T over (Y_T, ϕ_T) .

Proof. The proof proceeds in three steps. We first produce an approximate G_2 -instanton \tilde{A}_T by an explicit cut-and-paste procedure. This reduces the problem to solving the non-linear partial differential equation

$$(3.26) \quad d_{\tilde{A}_T}^* a = 0 \quad \text{and} \quad d_{\tilde{A}_T + a} \xi + *_T(F_{\tilde{A}_T + a} \wedge \psi_T) = 0.$$

for $a \in \Omega^1(Y_T, \mathfrak{g}_{E_T})$ and $\xi \in \Omega^0(Y_T, \mathfrak{g}_{E_T})$ where $\psi_T := *_T \phi_T$. Under the hypotheses of Theorem 3.24 we will show that we can solve the linearisation of (3.26) in a uniform fashion. The existence of a solution of (3.26) then follows from a simple application of Banach's fixed-point theorem.

Step 1. *There exists a $\delta < 0$ and for each $T \geq T_0$ there exists a connection \tilde{A}_T on a G -bundle E_T over Y_T such that*

$$(3.27) \quad \|F_{\tilde{A}_T} \wedge \psi_T\|_{C^{0, \alpha}} = O(e^{\delta T}).$$

The bundle E_T is constructed by gluing $E_{\pm}|_{Y_{T, \pm}}$ via \tilde{f} and the connection \tilde{A}_T is defined by

$$\tilde{A}_T := A_{\pm} - \tilde{\pi}_{\pm}^*[\chi(t - T + 1)a_{\pm}]$$

over $Y_{T, \pm}$ where

$$a_{\pm} := \tilde{\pi}_{\pm, *} A_{\pm} - A_{\infty, \pm},$$

$\tilde{\pi}_{\pm}$ is as in Definition 3.19 and χ is as in Section 2.1. Then (3.27) is a straight-forward consequence of (2.5) and (3.20).

Step 2. *Define a linear operator $L_T: C^{1, \alpha} \rightarrow C^{0, \alpha}$ by (3.11) with $A = \tilde{A}_T$ and $\phi = \phi_T$. Then there exist constants $\tilde{T}_1, c > 0$ such that for all $T \geq \tilde{T}_1$ the operator L_T is invertible and*

$$(3.28) \quad \|L_T^{-1} \underline{a}\|_{C^{1, \alpha}} \leq ce^{\frac{|\delta|}{4}T} \|\underline{a}\|_{C^{0, \alpha}}.$$

Step 2.1. *There exists a constant $c > 0$ such that for all $T \geq T_0$*

$$(3.29) \quad \|\underline{a}\|_{C^{1,\alpha}} \leq c \left(\|L_T \underline{a}\|_{C^{0,\alpha}} + \|\underline{a}\|_{L^\infty} \right).$$

This is an immediate consequence of standard interior Schauder estimates because of (2.5) and (3.20).

Step 2.2. *There exist constants $\tilde{T}_1 \geq T_0$ and $c > 0$ such that for $T \in [\tilde{T}_1, \infty)$*

$$(3.30) \quad \|\underline{a}\|_{L^\infty} \leq c e^{\frac{|\delta|}{4}T} \|L_T \underline{a}\|_{C^{0,\alpha}}.$$

Suppose not; then there exist a sequence (T_i) tending to infinity and a sequence (\underline{a}_i) such that

$$(3.31) \quad \|\underline{a}_i\|_{L^\infty} = 1 \quad \text{and} \quad \lim_{i \rightarrow \infty} e^{\frac{|\delta|}{4}T_i} \|L_{T_i} \underline{a}_i\|_{C^{0,\alpha}} = 0.$$

Then by (3.29)

$$(3.32) \quad \|\underline{a}_i\|_{C^{1,\alpha}} \leq 2c.$$

Hence, by Arzelà–Ascoli we can assume (passing to a subsequence) that the sequence $\underline{a}_i|_{Y_{T_i,\pm}}$ converges in $C_{\text{loc}}^{1,\alpha/2}$ to some section $\underline{a}_{\infty,\pm}$ of $(\Lambda^0 \oplus \Lambda^1) \otimes \mathfrak{g}_{E_\pm}$ over Y_\pm , which is bounded and satisfies

$$L_{A_\pm} \underline{a}_{\infty,\pm} = 0$$

because of (2.5) and (3.20). Using standard elliptic estimates it follows that $\underline{a}_{\infty,\pm} \in \mathcal{T}_{A_\pm}$.

Proposition 3.33. *In the above situation*

$$\lim_{i \rightarrow \infty} \left\| (\underline{a}_i|_{Y_{T_i,\pm}}) - (\underline{a}_{\infty,\pm}|_{Y_{T_i,\pm}}) \right\|_{L^\infty(Y_{T_i,\pm})} = 0.$$

The proof of this proposition will be given at the end of this section. Accepting it as a fact for now, it follows immediately that

$$\iota_+(\underline{a}_{\infty,+}) = \tilde{f}^* \circ \iota_-(\underline{a}_{\infty,-})$$

because $Y_{T_i,+} \cap Y_{T_i,-} = [T_i, T_i + 1] \times Z_+$. Now, by (3.25) we must have $\iota_\pm(\underline{a}_{\infty,\pm}) = 0$; hence, $\underline{a}_{\infty,\pm} = 0$, since ι_\pm are injective.

However, by (3.31) there exist $x_i \in Y_{T_i}$ such that $|\underline{a}_{T_i}|(x_i) = 1$. By passing to a further subsequence and possibly changing the rôles of + and – we can assume that each $x_i \in Y_{T_i,+}$; hence, by Proposition 3.33, $\underline{a}_{\infty,+} \neq 0$, contradicting what was derived above. This proves (3.30).

Step 2.3. We complete the proof of Step 2.

Combining (3.29) and (3.30) yields

$$\|\underline{a}\|_{C^{1,\alpha}} \leq ce^{\frac{|\delta|}{4}T} \|L_T \underline{a}\|_{C^{0,\alpha}}.$$

Therefore, L_T is injective; hence, also surjective since L_T is formally self-adjoint.

Step 3. There exists a constant $T_1 \geq \tilde{T}_1$ and for each $T \geq T_1$ a smooth solution $\underline{a} = \underline{a}_T$ of (3.26) such that $\lim_{T \rightarrow \infty} \|\underline{a}_T\|_{C^{1,\alpha}} = 0$.

We can write (3.26) as

$$(3.34) \quad L_T \underline{a} + Q_T(\underline{a}) + \varepsilon_T = 0$$

where $Q_T(\underline{a}) := \frac{1}{2} *_T ([a \wedge a] \wedge \psi_T) + [a, \xi]$ and $\varepsilon_T := *_T (F_{\tilde{A}_T} \wedge \psi_T)$. We make the ansatz $\underline{a} = L_T^{-1} \underline{b}$. Then (3.34) becomes

$$(3.35) \quad \underline{b} + \tilde{Q}_T(\underline{b}) + \varepsilon_T = 0$$

where $\tilde{Q}_T = Q_T \circ L_T^{-1}$. By (3.28)

$$\|\tilde{Q}_T(\underline{b}_1) - \tilde{Q}_T(\underline{b}_2)\|_{C^{0,\alpha}} \leq ce^{\frac{|\delta|}{2}T} (\|\underline{b}_1\|_{C^{0,\alpha}} + \|\underline{b}_2\|_{C^{0,\alpha}}) \|\underline{b}_1 - \underline{b}_2\|_{C^{0,\alpha}}$$

for some constant $c > 0$ independent of $T \geq \tilde{T}_1$. By Step 1, $\|\varepsilon_T\|_{C^{0,\alpha}} = O(e^{\delta T})$. Now, Lemma 3.36 yields the desired solution of (3.35) and thus of (3.26) provided $T \geq T_1$ for a suitably large $T_1 \geq \tilde{T}_1$. By elliptic regularity \underline{a} is smooth. \square

Lemma 3.36 (Donaldson and Kronheimer [DK90, Lemma 7.2.23]). *Let X be a Banach space and let $T: X \rightarrow X$ be a smooth map with $T(0) = 0$. Suppose there is a constant $c > 0$ such that*

$$\|Tx - Ty\| \leq c(\|x\| + \|y\|) \|x - y\|.$$

If $y \in X$ satisfies $\|y\| \leq \frac{1}{10c}$, then there exists a unique $x \in X$ with $\|x\| \leq \frac{1}{5c}$ solving

$$x + Tx = y.$$

Moreover, this $x \in X$ satisfies $\|x\| \leq 2\|y\|$.

To complete the proof of Theorem 3.24 it now remains to prove Proposition 3.33 for which we require the following result.

Proposition 3.37. *In the situation of Theorem 3.24, there is a $\gamma_0 > 0$ such that for each $\gamma \in (0, \gamma_0)$ the linear operator $L_{A_\pm}: C_Y^{1,\alpha} \rightarrow C_Y^{0,\alpha}$ has a bounded right inverse.*

Proof. By Proposition 3.4, $L_{A_{\pm}} : C_Y^{1,\alpha} \rightarrow C_Y^{0,\alpha}$ is Fredholm whenever $\gamma > 0$ is sufficiently small. The cokernel of $L_{A_{\pm}}$ can be identified to be $\mathcal{T}_{A_{\pm}, -\gamma}$, which is trivial by hypothesis. \square

Proof of Proposition 3.33. We restrict to the + case; the – case is identical. It follows from the construction of $\underline{a}_{\infty,+}$ that for each fixed compact subset $K \subset Y_+$

$$\lim_{i \rightarrow \infty} \left\| (\underline{a}_i|_K) - (\underline{a}_{\infty,+}|_K) \right\|_{L^\infty(K)} = 0.$$

To strengthen this to an estimate on all of $Y_{T_i,+}$ the factor $e^{\frac{|\delta|}{4}T}$ in (3.31) will be important, even though it is clearly not optimal.

With χ as in Definition 2.1 define a cut-off function $\chi_T : Y_+ \rightarrow [0, 1]$ by $\chi_T(x) := 1 - \chi(t_+(x) - \frac{3}{2}T)$. For each sufficiently small $\gamma > 0$ we have

$$\|L_{A_+}(\chi_{T_i}\underline{a}_i)\|_{C_Y^{0,\alpha}(Y_+)} = O(e^{-\frac{3}{2}\gamma T_i})$$

using the estimates (2.5), (3.20), (3.31) and (3.32). Using Proposition 3.37 we construct $\underline{b}_i \in C_Y^{1,\alpha}$ such that $\underline{a}_{\infty,+}^i := \chi_{T_i}\underline{a}_i + \underline{b}_i \in \mathcal{T}_{A_+,\gamma}$ and $\|\underline{b}_i\|_{C_{0,\gamma}^{1,\alpha}} = O(e^{-\frac{3}{2}\gamma T_i})$. Hence,

$$\left\| (\underline{a}_i|_{Y_{T_i,+}}) - (\underline{a}_{\infty,+}^i|_{Y_{T_i,+}}) \right\|_{L^\infty(Y_{T_i,+})} = O(e^{-\frac{1}{2}\gamma T_i}).$$

Moreover, $\lim_{i \rightarrow \infty} \left\| (\underline{a}_{\infty,+}^i|_K) - (\underline{a}_{\infty,+}|_K) \right\|_{L^\infty(K)} = 0$ and since both $\|\cdot\|_{L^\infty(K)}$ and $\|\cdot\|_{L^\infty(Y_+)}$ are norms on the finite dimensional vector space $\mathcal{T}_{A_+,\gamma} = \mathcal{T}_{A_+}$ it also follows that

$$\lim_{i \rightarrow \infty} \|\underline{a}_{\infty,+}^i - \underline{a}_{\infty,+}\|_{L^\infty(Y_+)} = 0.$$

Therefore,

$$\lim_{i \rightarrow \infty} \left\| (\underline{a}_i|_{Y_{T_i,+}}) - (\underline{a}_{\infty,+}|_{Y_{T_i,+}}) \right\|_{L^\infty(Y_{T_i,+})} = 0. \quad \square$$

Remark 3.38. The proof of Theorem 3.24 slightly simplifies assuming $\mathcal{H}_{A_{\infty,+}}^0 \oplus \mathcal{H}_{A_{\infty,+}}^1 = \{0\}$ instead of (3.25): We can directly conclude that $\iota_{\pm}(\underline{a}_{\infty,\pm}) = 0$ and, hence, $\underline{a}_{\infty,\pm} = 0$; thus making Proposition 3.33 unnecessary. In particular, (3.30) holds without the additional factor of $e^{\frac{|\delta|}{4}T}$.

4 From holomorphic vector bundles over building blocks to G_2 -instantons over ACyl G_2 -manifolds

We now discuss how to deduce Theorem 1.3 from Theorem 3.24.

Definition 4.1. Let (V, ω, Ω) be an ACyl Calabi–Yau 3–fold with asymptotic cross-section $(\Sigma, \omega_I, \omega_J, \omega_K)$. Let A_∞ be an ASD instanton on a G –bundle E_∞ over Σ . A HYM connection A on a G –bundle E over V is called **asymptotic** to A_∞ if there exist a constant $\delta < 0$ and a bundle isomorphism $\tilde{\pi}: E|_{V \setminus K} \rightarrow E_\infty$ covering $\pi: V \setminus K \rightarrow \mathbf{R}_+ \times S^1 \times \Sigma$ such that

$$\nabla^k(\tilde{\pi}_*A - A_\infty) = O(e^{\delta t})$$

for all $k \in \mathbf{N}_0$. Here by a slight abuse of notation we also denote by E_∞ and A_∞ their respective pullbacks to $\mathbf{R}_+ \times S^1 \times \Sigma$.

The following theorem can be used to produce examples of HYM connections A on $\mathrm{PU}(n)$ –bundles over ACyl Calabi–Yau 3–folds asymptotic to ASD instantons A_∞ ; hence, by taking the product with S^1 , examples of G_2 –instantons π_V^*A asymptotic to $\pi_\Sigma^*A_\infty$ over the ACyl G_2 –manifold $S^1 \times V$. Here $\pi_V: S^1 \times V \rightarrow V$ and $\pi_\Sigma: T^2 \times \Sigma \rightarrow \Sigma$ denote the canonical projections.

Theorem 4.2 (Sá Earp [Sá 15, Theorem 59]). *Let Z and Σ be as in Theorem 2.10 and let $(V := Z \setminus \Sigma, \omega, \Omega)$ be the resulting ACyl Calabi–Yau 3–fold. Let \mathcal{E} be a holomorphic vector bundle over Z and let A_∞ be an ASD instanton on $\mathcal{E}|_\Sigma$ compatible with the holomorphic structure. Then there exists a HYM connection A on $\mathcal{E}|_V$ which is compatible with the holomorphic structure on $\mathcal{E}|_V$ and asymptotic to A_∞ .*

By slight abuse of notation we also denote by A_∞ the ASD instanton on the $\mathrm{PU}(n)$ –bundle associated with $\mathcal{E}|_\Sigma$ and by A the HYM connection on the $\mathrm{PU}(n)$ –bundle associated with $\mathcal{E}|_V$. Theorem 3.24 and Theorem 4.2 together with the following result immediately imply Theorem 1.3.

Proposition 4.3. *In the situation of Theorem 4.2, suppose $H^0(\Sigma, \mathcal{E}\mathrm{nd}_0(\mathcal{E}|_\Sigma)) = 0$. Then*

$$(4.4) \quad \mathcal{H}_{\pi_\Sigma^*A_\infty}^1 = H_{A_\infty}^1,$$

see Definition 3.16 and Remark 1.6, and for some small $\delta > 0$ there exist injective linear maps

$$\begin{aligned} \kappa_-: \mathcal{T}_{\pi_V^*A, -\delta} &\rightarrow H^1(Z, \mathcal{E}\mathrm{nd}_0(\mathcal{E})(-\Sigma)) \\ \text{and } \kappa: \mathcal{T}_{\pi_V^*A} &\rightarrow H^1(Z, \mathcal{E}\mathrm{nd}_0(\mathcal{E})) \end{aligned}$$

such that the following diagram commutes:

$$(4.5) \quad \begin{array}{ccccc} \mathcal{T}_{\pi_V^*A, -\delta} & \longrightarrow & \mathcal{T}_{\pi_V^*A} & \xrightarrow{l} & \mathcal{H}_{\pi_\Sigma^*A_\infty}^1 \\ \downarrow \kappa_- & & \downarrow \kappa & & \downarrow \cong \\ H^1(Z, \mathcal{E}\mathrm{nd}_0(\mathcal{E})(-\Sigma)) & \longrightarrow & H^1(Z, \mathcal{E}\mathrm{nd}_0(\mathcal{E})) & \longrightarrow & H^1(\Sigma, \mathcal{E}\mathrm{nd}_0(\mathcal{E}|_\Sigma)). \end{array}$$

Equation (4.4) is a direct consequence of $\mathcal{H}_{A_\infty}^0 = 0$. The proof of the remaining assertions requires some preparation.

4.1 Comparing infinitesimal deformations of π_V^*A and A

Proposition 4.6. *If A is a HYM connection asymptotic to A_∞ , then there exists a $\delta_0 > 0$ such that for all $\delta \leq \delta_0$*

$$(4.7) \quad \mathcal{T}_{\pi_V^*A, \delta} = \left\{ \underline{a} \in \ker D_A : \nabla^k \bar{\pi}_* \underline{a} = O(e^{\delta t}) \text{ for all } k \in \mathbf{N}_0 \right\}$$

with D_A as in (3.14).

Proof. We can write $L_A = \tilde{I}\partial_\beta + D_A$ where β denotes the coordinate on S^1 . For $\delta \leq 0$, (4.7) follows by an application of Lemma A.1 in [Wal13b]. The right-hand side is contained in the left-hand side of (4.7) which, by Proposition 3.22, is independent of $\delta \in [0, \delta_0]$. \square

Proposition 4.8. *In the situation of Proposition 4.3, there exists a constant $\delta_0 > 0$ such that, for all $\delta \leq \delta_0$, $\mathcal{H}_{A, \delta}^0 = 0$ and*

$$\mathcal{T}_{\pi_V^*A, \delta} \cong \mathcal{H}_{A, \delta}^1$$

where

$$\mathcal{H}_{A, \delta}^i := \left\{ \alpha \in \mathcal{H}_A^i : \nabla^k \bar{\pi}_* \alpha = O(e^{\delta t}) \text{ for all } k \in \mathbf{N}_0 \right\}.$$

Proof. If $\delta \leq \delta_0$ (cf. Proposition 3.22) and $(\xi, \eta, a) \in \mathcal{T}_{A, \delta}$, then $\iota(\xi, \eta, a) \in \{0\} \oplus \mathcal{H}_{A_\infty}^1$. Hence ξ and η decay exponentially and one can use Proposition 4.6 and argue as in the proof of Proposition 3.18; it also follows that $\mathcal{H}_{A, \delta}^0 = 0$. \square

4.2 Acyclic resolutions via forms of exponential growth/decay

In view of the above what is missing to prove Proposition 4.3 is a way to relate $\mathcal{H}_{A, \delta}^1$ with the cohomology of (twists of) $\mathcal{E} \text{nd}_0(\mathcal{E})$. This is what the following result provides.

Proposition 4.9. *Let (Z, Σ) be a building block and let $V := Z \setminus \Sigma$ be the ACyl Calabi-Yau 3-fold constructed via Theorem 2.10. Suppose that \mathcal{E} is a holomorphic vector bundle over Z and suppose that A is a HYM connection on \mathcal{E} compatible with the holomorphic structure and asymptotic to an ASD instanton on $\mathcal{E}|_\Sigma$.*

For $\delta \in \mathbf{R}$ define a complex of sheaves $(\mathcal{A}_\delta^\bullet, \bar{d})$ on Z by

$$\mathcal{A}_\delta^i(U) = \left\{ \alpha \in \Omega^{0, i}(V \cap U, \mathcal{E}) : \nabla^k \bar{\pi}_* \alpha = O(e^{\delta t}) \text{ for all } k \in \mathbf{N}_0 \right\}.$$

If $\delta \in \mathbf{R} \setminus \mathbf{Z}$, then the complex of sheaves $(\mathcal{A}_\delta^\bullet, \bar{\partial})$ is an acyclic resolution of $\mathcal{E}(\lfloor \delta \rfloor \Sigma)$. In particular, setting $\kappa_\delta^i(\alpha) := [\alpha]$ one obtains maps

$$\kappa_\delta^i: \mathcal{H}_{A, \delta}^i \rightarrow H^i(\Gamma(\mathcal{A}_\delta^\bullet), \bar{\partial}) \cong H^i(Z, \mathcal{E}(\lfloor \delta \rfloor \Sigma)).$$

Remark 4.10. In Proposition 4.9, $\lfloor \delta \rfloor$ denotes the largest integer not greater than δ ; in particular, $\lfloor \delta \rfloor \Sigma$ is a divisor on Z .

Remark 4.11. We state Proposition 4.9 in dimension three; however, it works *mutatis mutandis* in all dimensions.

Proof of Proposition 4.9. The proof consists of three steps.

Step 1. The sheaves $\mathcal{A}_\delta^\bullet$ are C^∞ -modules; hence, acyclic, see [Dem12, Chapter IV Corollary 4.19].

Step 2. $\mathcal{E}(\lfloor \delta \rfloor \Sigma) = \ker(\bar{\partial}: \mathcal{A}_\delta^0 \rightarrow \mathcal{A}_\delta^1)$.

Let $x \in Z$ and let $U \subset Z$ denote a small open neighbourhood of x . An element $s \in \ker(\bar{\partial}: \Gamma(U, \mathcal{A}_\delta^0) \rightarrow \Gamma(U, \mathcal{A}_\delta^1))$ corresponds to a holomorphic section of $\mathcal{E}|_{V \cap U}$ such that $|z|^{-\delta} s$ stays bounded. Here z is a holomorphic function on U vanishing to first order along $\Sigma \cap U$, whose existence follows from Definition 2.8. Then $z^{-\lfloor \delta \rfloor} s$ is weakly holomorphic in U . By elliptic regularity $z^{-\lfloor \delta \rfloor} s$ extends across $U \cap \Sigma$ and thus s defines an element of $\Gamma(U, \mathcal{E}(\lfloor \delta \rfloor \Sigma))$. Conversely, it is clear that $\Gamma(U, \mathcal{E}(\lfloor \delta \rfloor \Sigma)) \subset \ker(\bar{\partial}: \Gamma(U, \mathcal{A}_\delta^0) \rightarrow \Gamma(U, \mathcal{A}_\delta^1))$.

Step 3. The complex of sheaves $(\mathcal{A}_\delta^\bullet, \bar{\partial})$ is exact.

Away from Σ the exactness follows from the usual $\bar{\partial}$ -Poincaré Lemma. If $x \in \Sigma$, then since Z is fibred over \mathbf{P}^1 , by Definition 2.8, there exist a small open neighbourhood U of x in Z , a polydisc $D \subset \Sigma$ centred at x and a biholomorphic map $\pi: V \cap U \rightarrow \mathbf{R}_+ \times S^1 \times D$ such that the push-forward of the Kähler metric on $V \cap U$ via π is asymptotic to the metric induced by that on D . The necessary version of the $\bar{\partial}$ -Poincaré Lemma can now be proved along the lines of [GH94, p. 25] provided the linear operator

$$\bar{\partial}: C_\delta^\infty \Omega^0(\mathbf{R} \times S^1) \rightarrow C_\delta^\infty \Omega^{0,1}(\mathbf{R} \times S^1)$$

is invertible. This, however, is a simple consequence of Theorem 3.1 since $\bar{\partial} = \partial_t + i\partial_\alpha$ and the spectrum of $i\partial_\alpha$ on $S^1 = \mathbf{R}/\mathbf{Z}$ is \mathbf{Z} . \square

4.3 Proof of Proposition 4.3

In view of Proposition 4.8 we only need to establish (4.5) with $\mathcal{H}_{A,\delta}^1$ instead of $\mathcal{T}_{\pi_V^*A,\delta}$. By Proposition 4.9 applied to $\mathcal{E}nd_0(\mathcal{E})$, we have linear maps

$$\kappa_\delta^1: \mathcal{H}_{A,\delta}^1 \rightarrow H^1(Z, \mathcal{E}nd_0(\mathcal{E})([\delta]\Sigma)) \quad \text{for } \delta \in \mathbf{R} \setminus \mathbf{Z};$$

hence, linear maps

$$\begin{aligned} \kappa_-: \mathcal{H}_{A,-\delta}^1 &\rightarrow H^1(Z, \mathcal{E}nd_0(\mathcal{E})(-\Sigma)) \\ \text{and } \kappa: \mathcal{H}_A^1 &= \mathcal{H}_{A,\delta}^1 \rightarrow H^1(Z, \mathcal{E}nd_0(\mathcal{E})) \end{aligned}$$

for some small $\delta > 0$ making the following diagram commute:

$$\begin{array}{ccccc} \mathcal{H}_{A,-\delta}^1 & \longrightarrow & \mathcal{H}_A^1 & \xrightarrow{l} & H_{A_\infty}^1 \\ \downarrow \kappa_- & & \downarrow \kappa & & \downarrow \cong \\ H^1(Z, \mathcal{E}nd_0(\mathcal{E})(-\Sigma)) & \longrightarrow & H^1(Z, \mathcal{E}nd_0(\mathcal{E})) & \longrightarrow & H^1(\Sigma, \mathcal{E}nd_0(\mathcal{E}|_\Sigma)). \end{array}$$

The map κ_- is injective, because if $\kappa_- a = 0$, then $a = \bar{\partial}s$ for some $s \in \Gamma(Z, \mathcal{A}_{-\delta}^0)$ and thus

$$\int_V \|a\|^2 = \int_V \langle a, \bar{\partial}s \rangle = \int_V \langle \bar{\partial}^* a, s \rangle = 0.$$

Since $H^0(\Sigma, \mathcal{E}nd_0(\mathcal{E}|_\Sigma)) = 0$, the first map on the bottom is injective and because the rows are exact a simple diagram chase proves shows that κ is injective. \square

References

- [Ati88] M. Atiyah. “New invariants of 3– and 4–dimensional manifolds”. In: *The mathematical heritage of Hermann Weyl (Durham, NC, 1987)*. Vol. 48. Proc. Sympos. Pure Math. Providence, RI, 1988, pp. 285–299. MR: 974342 (cit. on p. 4).
- [CHNP13] A. Corti, M. Haskins, J. Nordström, and T. Pacini. “Asymptotically cylindrical Calabi–Yau 3–folds from weak Fano 3–folds”. In: *Geom. Topol.* 17.4 (2013), pp. 1955–2059. DOI: 10.2140/gt.2013.17.1955. MR: 3109862. Zbl: 1273.14081 (cit. on p. 7).
- [CHNP15] A. Corti, M. Haskins, J. Nordström, and T. Pacini. “ G_2 –manifolds and associative submanifolds via semi-Fano 3–folds”. In: *Duke Math. J.* 164.10 (2015), pp. 1971–2092. DOI: 10.1215/00127094-3120743. MR: 3369307. Zbl: 06486366 (cit. on pp. 1, 5, 7).

- [Dem12] J.-P. Demailly. *Complex Analytic and Differential Geometry*. 2012. URL: <https://www-fourier.ujf-grenoble.fr/~demailly/manuscripts/agbook.pdf> (cit. on p. 20).
- [DK90] S. K. Donaldson and P. B. Kronheimer. *The geometry of four-manifolds*. Oxford Mathematical Monographs. Oxford Science Publications. New York, 1990, pp. x+440. MR: MR1079726. Zbl: 0904.57001 (cit. on pp. 3, 16).
- [Don02] S. K. Donaldson. *Floer homology groups in Yang–Mills theory*. Vol. 147. Cambridge Tracts in Mathematics. With the assistance of M. Furuta and D. Kotschick. Cambridge, 2002, pp. viii+236. DOI: 10.1017/CB09780511543098. MR: 1883043 (cit. on p. 9).
- [Don85] S. K. Donaldson. “Anti self-dual Yang–Mills connections over complex algebraic surfaces and stable vector bundles”. In: *Proc. London Math. Soc.* (3) 50.1 (1985), pp. 1–26. DOI: 10.1112/plms/s3-50.1.1. MR: 765366. Zbl: 0529.53018 (cit. on pp. 2, 10).
- [DS11] S. K. Donaldson and E. P. Segal. “Gauge theory in higher dimensions, II”. In: *Surveys in differential geometry. Volume XVI. Geometry of special holonomy and related topics*. Vol. 16. Surv. Differ. Geom. 2011, pp. 1–41. arXiv: 0902.3239. MR: 2893675. Zbl: 1256.53038 (cit. on p. 2).
- [DT98] S. K. Donaldson and R. P. Thomas. “Gauge theory in higher dimensions”. In: *The geometric universe (Oxford, 1996)*. Oxford, 1998, pp. 31–47. MR: MR1634503. Zbl: 0926.58003. URL: <http://www.ma.ic.ac.uk/~rpwt/skd.pdf> (cit. on p. 2).
- [GH94] P. Griffiths and J. Harris. *Principles of algebraic geometry*. Wiley Classics Library. Reprint of the 1978 original. New York, 1994, pp. xiv+813. MR: 1288523 (cit. on p. 20).
- [HHN15] M. Haskins, H.-J. Hein, and J. Nordström. “Asymptotically cylindrical Calabi–Yau manifolds”. In: *J. Differential Geom.* 101.2 (2015), pp. 213–265. MR: 3399097. URL: <http://projecteuclid.org/euclid.jdg/1442364651> (cit. on pp. 2, 7, 9).
- [HL97] D. Huybrechts and M. Lehn. *The geometry of moduli spaces of sheaves*. Aspects of Mathematics, E31. Braunschweig, 1997, pp. xiv+269. MR: 1450870 (cit. on p. 3).
- [Joy00] D. D. Joyce. *Compact manifolds with special holonomy*. Oxford Mathematical Monographs. Oxford, 2000, pp. xii+436. MR: 1787733. Zbl: 1027.53052 (cit. on p. 7).

- [Joy96] D. D. Joyce. “Compact Riemannian 7-manifolds with holonomy G_2 . I”. In: *J. Differential Geom.* 43.2 (1996), pp. 291–328. DOI: 10.4310/jdg/1214458109. MR: MR1424428. Zbl: 0861.53022 (cit. on p. 1).
- [KL11] A. Kovalev and N.-H. Lee. “ $K3$ surfaces with non-symplectic involution and compact irreducible G_2 -manifolds”. In: *Math. Proc. Cambridge Philos. Soc.* 151.2 (2011), pp. 193–218. DOI: 10.1017/S030500411100003X. MR: 2823130. Zbl: 1228.53064 (cit. on pp. 1, 7).
- [Kovo03] A. Kovalev. “Twisted connected sums and special Riemannian holonomy”. In: *J. Reine Angew. Math.* 565 (2003), pp. 125–160. DOI: 10.1515/crll.2003.097. MR: MR2024648. Zbl: 1043.53041 (cit. on pp. 1, 2, 6, 7).
- [LM85] R. B. Lockhart and R. C. McOwen. “Elliptic differential operators on noncompact manifolds”. In: *Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4)* 12.3 (1985), pp. 409–447. MR: 837256. URL: http://www.numdam.org/item?id=ASNSP_1985_4_12_3_409_0 (cit. on pp. 8, 13).
- [MNS15] G. Menet, J. Nordström, and H. N. Sá Earp. *Construction of G_2 -instantons via twisted connected sums*. 2015. arXiv: 1510.03836 (cit. on p. 4).
- [MP78] V. G. Maz’ya and B. A. Plamenevskiĭ. “Estimates in L_p and in Hölder classes, and the Miranda–Agmon maximum principle for the solutions of elliptic boundary value problems in domains with singular points on the boundary”. In: *Math. Nachr.* 81 (1978), pp. 25–82. MR: 0492821 (cit. on p. 8).
- [Sá 15] H. N. Sá Earp. “ G_2 -instantons over asymptotically cylindrical manifolds”. In: *Geom. Topol.* 19.1 (2015), pp. 61–111. DOI: 10.2140/gt.2015.19.61. MR: 3318748. Zbl: 1312.53044 (cit. on pp. 1, 2, 10, 18).
- [Tyuo8] A. Tyurin. *Vector bundles*. Collected works. Volume I, Edited by Fedor Bogomolov, Alexey Gorodentsev, Victor Pidstrigach, Miles Reid and Nikolay Tyurin. 2008, p. 330. MR: 2742585 (cit. on p. 3).
- [UY86] K. K. Uhlenbeck and S.-T. Yau. “On the existence of Hermitian–Yang–Mills connections in stable vector bundles”. In: *Comm. Pure Appl. Math.* 39.S, suppl. (1986). *Frontiers of the mathematical sciences: 1985* (New York, 1985), S257–S293. DOI: 10.1002/cpa.3160390714. MR: 861491 (cit. on p. 10).
- [Wal13a] T. Walpuski. “Gauge theory on G_2 -manifolds”. PhD thesis. Imperial College London, 2013. URL: <https://spiral.imperial.ac.uk/bitstream/10044/1/14365/1/Walpuski-T-2013-PhD-Thesis.pdf> (cit. on p. 2).

- [Wal13b] T. Walpuski. “ G_2 -instantons on generalised Kummer constructions”. In: *Geom. Topol.* 17.4 (2013), pp. 2345–2388. DOI: 10.2140/gt.2013.17.2345. MR: 3110581. Zbl: 1278.53051 (cit. on p. 19).
- [Wal15] T. Walpuski. “ G_2 -instantons over twisted connected sums: an example”. In: *Mathematical Research Letters* 23.2 (2015), pp. 529–544. DOI: 10.4310/MRL.2016.v23.n2.a11. arXiv: 1505.01080. Zbl: 06609380 (cit. on p. 4).