

# $G_2$ -instantons over twisted connected sums: an example

Thomas Walpuski

2016-07-31

## Abstract

Using earlier work of Sá Earp and the author [SW15] we construct an irreducible unobstructed  $G_2$ -instanton on an  $SO(3)$ -bundle over a twisted connected sum  $G_2$ -manifold recently discovered by Crowley and Nordström [CN14].

**Changes to the published version** A former version of this article has been published in *Mathematical Research Letters*, Volume 23, Issue 2, pp. 529–544 (2016). The present version is identical to the published article, except for aesthetic changes and the addition of a paragraph in Section 2.3 justifying the application of [Moř67, Theorem 7.5] in more detail.

## 1 Introduction

In order to put this note into context and help the reader appreciate its significance, we (very) briefly recall some ideas from the study of gauge theory on  $G_2$ -manifolds.

**Definition 1.1.** A connection  $A \in \mathcal{A}(E)$  on a  $G$ -bundle  $E$  over a  $G_2$ -manifold  $Y$  is called a  $G_2$ -instanton if its curvature satisfies

$$(1.2) \quad F_A \wedge \psi = 0$$

with  $\psi := *\phi$  and  $\phi \in \Omega^3(Y)$  denoting the  $G_2$ -structure on  $Y$ .

In their visionary article [DT98] Donaldson and Thomas speculated that “counting”  $G_2$ -instantons might lead to an interesting enumerative invariant. Although almost two decades have passed, it is still not understood what the precise definition of this invariant ought to be; however, see Donaldson and Segal [DS11], the author [Wal17; Wal13b], and Haydys and the author [HW15] for some recent progress. What is clear, nonetheless, is that irreducible unobstructed  $G_2$ -instantons should contribute with  $\pm 1$  (depending on orientations).

**Definition 1.3.** A  $G_2$ -instanton  $A \in \mathcal{A}(E)$  is called **irreducible (unobstructed)** if the elliptic complex

$$\Omega^0(Y, \mathfrak{g}_E) \xrightarrow{d_A} \Omega^1(Y, \mathfrak{g}_E) \xrightarrow{\psi \wedge d_A} \Omega^6(Y, \mathfrak{g}_E) \xrightarrow{d_A} \Omega^7(Y, \mathfrak{g}_E)$$

has vanishing cohomology in degree zero (one).

In [SW15] Sá Earp and the author developed a method for constructing irreducible unobstructed  $G_2$ -instantons over twisted connected sums. So far, however, we were unable to find a single instance of the input required for this construction. This brief note is meant to ameliorate this disgraceful situation by showing that our method can be used to produce at least one example.

Let us briefly recall the twisted connected sum construction, a rich source of  $G_2$ -manifolds, which was suggested by Donaldson, pioneered by Kovalev [Kov03] and later extended and improved by Kovalev and Lee [KL11], and Corti, Haskins, Nordström and Pacini [CHNP15].

**Definition 1.4.** A **building block** is a non-singular algebraic 3-fold  $Z$  together with a projective morphism  $f: Z \rightarrow \mathbf{P}^1$  such that:

- the anticanonical class  $-K_Z \in H^2(Z)$  is primitive,
- $\Sigma := f^{-1}(\infty)$  is a smooth K3 surface and  $\Sigma \sim -K_Z$ .

A **framing** of a building block  $(Z, \Sigma)$  consists of a hyperkähler structure  $\omega = (\omega_I, \omega_J, \omega_K)$  on  $\Sigma$  such that  $\omega_J + i\omega_K$  is of type  $(2, 0)$  as well as a Kähler class on  $Z$  whose restriction to  $\Sigma$  is  $[\omega_I]$ .<sup>1</sup>

For the purpose of this article we are mostly interested in the following class of building blocks introduced by Kovalev [Kov03].

**Definition 1.5.** A building block is said to be of **Fano type** if it is obtained by blowing-up a Fano 3-fold  $W$  along the base locus of general pencil  $|\Sigma_0, \Sigma_\infty| \subset |-K_W|$ . (See Section 2.1 for more details on this construction.)

Given a framed building block  $(Z, \Sigma, \omega)$ , using the work of Haskins, Hein and Nordström [HHN15], we can make  $V := Z \setminus \Sigma$  into an asymptotically cylindrical (ACyl) Calabi–Yau 3-fold with asymptotic cross-section  $S^1 \times \Sigma$ ; hence,  $Y := S^1 \times V$  is an ACyl  $G_2$ -manifold with asymptotic cross-section  $T^2 \times \Sigma$ .

**Definition 1.6.** A **matching** of pair of framed building blocks  $(Z_\pm, \Sigma_\pm, \omega_\pm)$  is a hyperkähler rotation  $r: \Sigma_+ \rightarrow \Sigma_-$ , i.e., a diffeomorphism such that

$$r^* \omega_{I,-} = \omega_{J,+}, \quad r^* \omega_{J,-} = \omega_{I,+} \quad \text{and} \quad r^* \omega_{K,-} = -\omega_{K,+}.$$

Given a matched pair of framed building blocks  $(Z_\pm, \Sigma_\pm, \omega_\pm; r)$ , the twisted connected sum construction produces a simply-connected compact 7-manifold  $Y$  together with a family of torsion-free  $G_2$ -structures  $\{\phi_T : T \gg 1\}$  by gluing truncations of  $Y_\pm$  along their boundaries via interchanging the circle factors and  $r$ .

Sá Earp [Sá15] proved that given a holomorphic vector bundle  $\mathcal{E}$  on a building block with  $\mathcal{E}|_\Sigma$   $\mu$ -stable,<sup>2</sup> the smooth vector bundle underlying  $\mathcal{E}|_V$  can be equipped with a Hermitian Yang–Mills connection which is asymptotic at infinity to the anti-self-dual connection  $A_\infty$  inducing the

<sup>1</sup>The existence of such a class is not guaranteed a priori.

<sup>2</sup>Recall that a holomorphic bundle  $\mathcal{E}$  on a compact Kähler  $n$ -fold  $(X, \omega)$  is  $\mu$ -**(semi)stable** if for each torsion-free coherent subsheaf  $\mathcal{F} \subset \mathcal{E}$  with  $0 < \text{rk } \mathcal{F} < \text{rk } \mathcal{E}$  we have  $(\mu(\mathcal{F}) \leq \mu(\mathcal{E})) \mu(\mathcal{F}) < \mu(\mathcal{E})$ . Here  $\mu(\mathcal{E}) := \langle c_1(\mathcal{E}) \cup [\omega]^{n-1}, [X] \rangle / \text{rk } \mathcal{E}$  is the **slope** of  $\mathcal{E}$  (and similarly for  $\mathcal{F}$ ).

holomorphic structure on  $\mathcal{E}|_\Sigma$  [Don85]. Building on this, Sá Earp and the author [SW15] developed a method for constructing  $G_2$ -instantons over twisted connected sums provided a pair  $\mathcal{E}_\pm$  of such bundles and a lift  $\bar{r}: \mathcal{E}_+|_{\Sigma_+} \rightarrow \mathcal{E}_-|_{\Sigma_-}$  of the hyperkähler rotation  $r$ , which pulls back  $A_{\infty,-}$  to  $A_{\infty,+}$  (and assuming certain transversality conditions). The following is a *very special* case of the main result of [SW15].

**Theorem 1.7.** *Let  $(Z_\pm, \Sigma_\pm, \omega_\pm; r)$  be a matched pair of framed building blocks. Denote by  $Y$  the compact 7-manifold and by  $\{\phi_T : T \gg 1\}$  the family of torsion-free  $G_2$ -structures obtained from the twisted connected sum construction. Let  $\mathcal{E}_\pm \rightarrow Z_\pm$  be a pair of rank  $r$  holomorphic vector bundles such that the following hold:*

- $c_1(\mathcal{E}_+|_{\Sigma_+}) = r^*c_1(\mathcal{E}_-|_{\Sigma_-})$  and  $c_2(\mathcal{E}_+|_{\Sigma_+}) = r^*c_2(\mathcal{E}_-|_{\Sigma_-})$ .
- $\mathcal{E}_\pm|_{\Sigma_\pm}$  is  $\mu$ -stable with respect to  $\omega_{I,\pm}$  and spherical, i.e.,

$$H^*(\Sigma_\pm, \mathcal{E}nd_0(\mathcal{E}_\pm|_{\Sigma_\pm})) = 0.$$

- $\mathcal{E}_\pm$  is infinitesimally rigid:

$$(1.8) \quad H^1(Z_\pm, \mathcal{E}nd_0(\mathcal{E}_\pm)) = 0.$$

Then there exists a  $U(r)$ -bundle  $E$  over  $Y$  with

$$(1.9) \quad c_1(E) = Y(c_1(\mathcal{E}_+), c_1(\mathcal{E}_-)) \quad \text{and} \quad c_2(E) = Y(c_2(\mathcal{E}_+), c_2(\mathcal{E}_-))$$

and a family of connections  $\{A_T : T \gg 1\}$  on the associated  $PU(r)$ -bundle with  $A_T$  being an irreducible unobstructed  $G_2$ -instanton over  $(Y, \phi_T)$ .

*Remark 1.10.* The map

$$Y: \{([\alpha_+], [\alpha_-]) \in H^{\text{ev}}(Z_+) \times H^{\text{ev}}(Z_-) : [\alpha_+]|_{\Sigma_+} = r^*([\alpha_-]|_{\Sigma_-})\} \rightarrow H^{\text{ev}}(Y)$$

is the natural patching map denoted by  $Y$  in [CHNP15, Definition 4.15].

Let  $\text{res}_\pm: H^2(Z_\pm) \rightarrow H^2(\Sigma_\pm)$  denote the restriction maps associated with the inclusions  $\Sigma_\pm \subset Z_\pm$  and set

$$N_\pm := \text{im res}_\pm.$$

If  $\mathcal{E}_\pm|_{\Sigma_\pm}$  is spherical, then  $c_1(\mathcal{E}_\pm|_{\Sigma_\pm})$  must be non-zero; hence, Theorem 1.7 cannot be applied in situations where  $N_+ \cap r^*N_- = 0$ . In particular, this rules out all the examples in [Kov03; KL11] as well as the mass-produced examples in [CHNP15]. This means that the list of currently known  $G_2$ -manifolds to which Theorem 1.7 could potentially be applied is relatively short. Moreover, it has proved rather difficult to find suitable  $\mathcal{E}_\pm$ .

Crowley and Nordström [CN14] systematically studied twisted connected sums of building blocks arising from Fano 3-folds with Picard number two; in particular, those that arise from matchings with  $N_+ \cap r^*N_- \neq 0$ . This note shows that for one such twisted connected sum the hypotheses of Theorem 1.7 can be satisfied.

**Theorem 1.11.** *There exists a twisted connected sum  $Y$  of a pair of Fano type building blocks  $(Z_{\pm}, \Sigma_{\pm})$ , arising from #13 and #14 in Mori and Mukai’s classification of Fano 3–folds with Picard number two [MM81, Table 2], admitting a pair of rank 2 holomorphic vector bundles  $\mathcal{E}_{\pm}$  as required by Theorem 1.7. In particular, each of the resulting twisted connected sums  $(Y, \phi_T)$  with  $T \gg 1$  carries an irreducible unobstructed  $G_2$ –instanton on an  $SO(3)$ –bundle.*

*Remark 1.12.* In earlier work [Wal13a] the author constructed examples of irreducible unobstructed  $G_2$ –instantons over  $G_2$ –manifolds arising from Joyce’s generalised Kummer construction [Joy96a; Joy96b]. To the author’s best knowledge, Theorem 1.11 provides the first example of an irreducible unobstructed  $G_2$ –instanton over a twisted connected sum.

The method of proof relies mostly on certain arithmetic properties enjoyed by the twisted connected sum listed as [CN14, Table 4, Line 16] by Crowley and Nordström. A more abstract existence theorem is stated as Theorem 3.14. It is an interesting question to ask whether there are any further twisted connected sums to which this result can be applied.

Finally, it should be pointed out that there is a very recent preprint by Menet, Nordström and Sá Earp [MNS17] in which they use the more general main result of [SW15] to construct one  $G_2$ –instanton.

**Acknowledgements** The author is grateful to Johannes Nordström for pointing out the  $G_2$ –manifold constructed in [CN14] on which the above example lives; moreover, he thanks the anonymous referees for thoughtful comments and suggestions. The author gratefully acknowledges support from the Simons Center for Geometry and Physics, Stony Brook University at which part of the research for this paper was carried out.

## 2 The twisted connected sum

In this section we provide further details on Fano type building blocks, explain how to construct matching pairs of framed building blocks and describe the twisted connected sum mentioned in Theorem 1.11.

### 2.1 Building blocks of Fano type

If  $W$  is a Fano 3–fold, then according to Shokurov [Šok79] a general divisor  $\Sigma \in |-K_W|$  is a smooth  $K_3$  surface. Given a general pencil  $|\Sigma_0, \Sigma_{\infty}| \subset |-K_W|$ , blowing-up its base locus yields a smooth 3–fold  $Z$  together with a base-point free anti-canonical pencil spanned by the proper transforms of  $\Sigma_0$  and  $\Sigma_{\infty}$ . The resulting projective morphism  $f: Z \rightarrow \mathbf{P}^1$  makes  $(Z, \Sigma_{\infty})$  into a building block with

$$(2.1) \quad N := \text{im}(\text{res}: H^2(Z) \rightarrow H^2(\Sigma)) \cong \text{Pic}(W),$$

see [Kov03, Proposition 6.42] and [CHNP15, Proposition 3.15].

Mořezon [Moř67, Theorem 7.5] showed that if  $-K_W$  is very ample for a very general<sup>3</sup>  $\Sigma \in |-K_W|$  we have  $\text{Pic}(\Sigma) = \text{Pic}(W)$ . Moreover, according to Kovalev [Kovo09, Proposition 2.14] (see also, Voisin [Voio07, Corollary 2.10]) we can assume that  $f: Z \rightarrow \mathbf{P}^1$  is a **rational double point (RDP)  $K3$  fibration**, by which we mean that it has at only finitely many singular fibres and the singular fibres have only RDP singularities. (In fact, Kovalev asserts that generically the singular fibres have only ordinary double points.)

## 2.2 Matching building blocks

Fix a lattice  $L$  which is isomorphic to  $(H^2(\Sigma), \cup)$  for  $\Sigma$  a  $K3$  surface. Using the Torelli theorem and Yau's solution to the Calabi conjecture, Corti, Haskins, Nordström, and Pacini [CHNP15, section 6] showed that a set of framings of a pair of building blocks  $Z_{\pm}$  together with a matching is equivalent (up to the action of  $O(L)$ ) to lattice isomorphisms  $h_{\pm}: L \rightarrow H^2(\Sigma_{\pm})$  and an orthonormal triple  $(k_+, k_-, k_0)$  of positive classes in  $L_{\mathbf{R}} := L \otimes_{\mathbf{Z}} \mathbf{R}$  with  $h_{\pm}(k_{\pm})$  the restriction of a Kähler class on  $Z_{\pm}$  and  $\langle k_{\mp}, \pm k_0 \rangle$  the period point of  $(\Sigma_{\pm}, h_{\pm})$ . (The corresponding framings have  $[\omega_{I, \pm}] = h_{\pm}(k_{\pm})$  and the matching is such that  $r^* = h_+ \circ h_-^{-1}$ .) The following definition is useful to further simplify the matching problem.

**Definition 2.2.** Let  $\mathcal{Z}$  be a family of building blocks with constant  $N$  and a fixed primitive isometric embedding  $N \subset L$ . Let  $\text{Amp}$  be an open subcone of the positive cone in  $N_{\mathbf{R}}$ .  $\mathcal{Z}$  is called  $(N, \text{Amp})$ -**generic** if there exists a subset  $U_{\mathcal{Z}} \subset D_N := \{\Pi \in \mathbf{P}(N_{\mathbf{C}}^{\perp}) : \Pi \bar{\Pi} > 0\}$  with complement a countable union of complex analytic submanifolds of positive codimension and with the property that for any  $\Pi \in U_{\mathcal{Z}}$  and  $k \in \text{Amp}$  there exists a  $(Z, \Sigma) \in \mathcal{Z}$  and a marking  $h: L \rightarrow H^2(\Sigma)$  such that  $\Pi$  is the period point of  $(\Sigma, h)$  and  $h(k)$  is the restriction to  $\Sigma$  of a Kähler class on  $Z$ .

This definition slightly deviates from [CHNP15, Definition 6.17]. There it is required that the complement of  $U_{\mathcal{Z}}$  is a *locally finite* union of complex analytic submanifolds of positive codimension. The above slightly weaker condition still suffices for the proof of the next proposition to carry over verbatim.

**Proposition 2.3** ([CHNP15, Proposition 6.18]). *Let  $N_{\pm} \subset L$  be a pair of primitive sublattices of signature  $(1, r_{\pm} - 1)$  and let  $\mathcal{Z}_{\pm}$  be a pair of  $(N_{\pm}, \text{Amp}_{\pm})$ -generic families of building blocks. Suppose that  $W := N_+ + N_-$  is an orthogonal pushout.<sup>4</sup> Set  $T_{\pm} := N_{\pm}^{\perp}$  and  $W_{\pm} := N_{\pm} \cap T_{\mp}$ . If*

$$\text{Amp}_{\pm} \cap W_{\pm} \neq \emptyset,$$

*then there exist  $(Z_{\pm}, \Sigma_{\pm}) \in \mathcal{Z}_{\pm}$ , markings  $h_{\pm}: L \rightarrow H^2(\Sigma_{\pm})$  compatible with the given embeddings  $N_{\pm} \subset L$  and an orthonormal triple  $(k_+, k_-, k_0)$  of positive classes in  $L_{\mathbf{R}}$  with:*

- $k_{\pm} \in \text{Amp}_{\pm} \cap W_{\pm, \mathbf{R}}$  and  $k_0 \in W^{\perp}$ ,

<sup>3</sup>Here **very general** means that the set of  $\Sigma \in |-K_W|$  not satisfying the asserted condition is a countable union of complex analytic submanifolds of positive codimension in  $\text{PH}^0(-K_W)$ .

<sup>4</sup>This means that  $W_{\mathbf{R}} = W_{+, \mathbf{R}} \oplus W_{-, \mathbf{R}} \oplus (N_{+, \mathbf{R}} \cap N_{-, \mathbf{R}})$ .

- $h_{\pm}(k_{\pm})$  the restriction of a Kähler class on  $Z_{\pm}$ , and
- $\langle k_{\mp}, \pm k_0 \rangle$  the period point of  $(\Sigma_{\pm}, h_{\pm})$ .

If  $\mathcal{Z}$  is a family of building blocks arising from a full deformation type of Fano 3–folds, then we can always find an open subcone  $\text{Amp}$  of the positive cone such that  $\mathcal{Z}$  is  $(N, \text{Amp})$ –generic [CHNP13, Proposition 6.9]. (Also Definition 2.2 allows to slightly shrink  $\mathcal{Z}$  from a full deformation type by imposing very general conditions in the sense of Footnote 3.) This reduces finding a matching of a pair such families of building blocks to the arithmetic problem of embedding  $N_{\pm}$  into  $L$  compatible with Proposition 2.3.

### 2.3 An example due to Crowley and Nordström

We will now describe the twisted connected sum found by Crowley and Nordström [CN14, Table 4, Line 16] which we referred to in Theorem 1.11.

Consider the following pair of Fano 3–folds:

- Denote by  $Q \subset \mathbf{P}^4$  a smooth quadric. Let  $W_+ \rightarrow Q$  denote the blow-up of  $Q$  in a degree 6 genus 2 curve [MM81, Table 2, #13].
- Denote by  $V_5$  a section of the Plücker-embedded Grassmannian  $\text{Gr}(2, 5) \subset \mathbf{P}^9$  by a subspace of codimension 3. Let  $W_- \rightarrow V_5$  denote the blow-up of  $V_5$  in an elliptic curve that is the intersection of two hyperplane sections [MM81, Table 2, #14].

The anticanonical divisors  $-K_{W_{\pm}}$  both are very ample. To see this note that by according to [Isk78, Section 1] if  $W$  is an index  $r$  Fano 3–fold and  $-K_W$  is not very ample, then either  $|\frac{1}{r}K_W|$  has a base point or  $W$  is hyperelliptic. According to [IP99, Remarks preceding Table 12.3] neither is the case for the Fano 3–folds under consideration; see also [IP99, Theorem 2.1.16 and Theorem 2.4.5].

$W_{\pm}$  both have Picard number 2 with  $\text{Pic}(W_{\pm})$  generated by  $H_{\pm}$ , the pullback of a generator of  $\text{Pic}(Q)$  and  $\text{Pic}(V_5)$  respectively, and the exceptional divisor  $E_{\pm}$ . With respect to the bases  $(H_{\pm}, E_{\pm})$  the intersection forms on  $N_{\pm} = \text{Pic}(W_{\pm})$ , see (2.1), can be written as

$$\begin{pmatrix} 6 & 6 \\ 6 & 2 \end{pmatrix} \quad \text{and} \quad \begin{pmatrix} 10 & 5 \\ 5 & 0 \end{pmatrix}$$

respectively.

$N_{\pm}$  can be thought of as the overlattices  $\mathbf{Z}^2 + \frac{1}{5}(3, -1)\mathbf{Z}$  and  $\mathbf{Z}^2 + \frac{1}{6}(1, 1)\mathbf{Z}$  of  $\mathbf{Z}^2$ , generated by

$$(2.4) \quad \begin{aligned} A_+ &= 3H_+ - E_+ & \text{and} & & B_+ &= 4H_+ - 3E_+, \\ \text{and } A_- &= 3H_- - 2E_- & \text{and} & & B_- &= 3H_- - 4E_-, \end{aligned}$$

with intersection forms

$$\begin{pmatrix} 20 & 0 \\ 0 & -30 \end{pmatrix} \quad \text{and} \quad \begin{pmatrix} 30 & 0 \\ 0 & -30 \end{pmatrix}$$

respectively. The overlattice  $W := \mathbf{Z}^3 + \frac{1}{5}(3, 0, -1)\mathbf{Z} + \frac{1}{6}(0, 1, 1)\mathbf{Z}$  of  $\mathbf{Z}^3$  with intersection form

$$\begin{pmatrix} 20 & 0 & 0 \\ 0 & 30 & 0 \\ 0 & 0 & -30 \end{pmatrix}$$

is an orthogonal pushout of  $N_{\pm}$  along  $R = N_+ \cap N_- = (-30)$ .

By Nikulin [Nik79, Theorem 1.12.4 and Corollary 1.12.3] the lattice  $W$  (and thus also  $N_{\pm}$ ) can be embedded primitively into  $L$ . Since we can choose  $\text{Amp}_{\pm}$  such that  $\text{Amp}_{\pm} \cap W_{\pm}$  is spanned by  $A_{\pm}$ , Proposition 2.3 yields matching data with  $k_{\pm}$  a multiple of  $A_{\pm}$  for a pair of building blocks  $(Z_{\pm}, \Sigma_{\pm})$  of Fano type arising from  $W_{\pm}$ . Moreover, the resulting matching  $r$  is such that  $B_+ = r^*B_-$  (which generates  $N_+ \cap r^*N_-$ ). By the discussion at the end of Section 2.1 we may assume that for all but countably many  $b \in \mathbf{P}^1$  the fibre  $\Sigma_{\pm, b} := f_{\pm}^{-1}(b)$  satisfies  $\text{Pic}(\Sigma_{\pm, b}) = N_{\pm}$ ; in particular, we may assume that this holds for  $\Sigma_{\pm} = \Sigma_{\pm, \infty}$ . Moreover, we may assume that  $f_{\pm}: Z_{\pm} \rightarrow \mathbf{P}^1$  is an RDP  $K3$  fibration.

### 3 Bundles on the building blocks

We will now construct holomorphic vector bundles  $\mathcal{E}_{\pm}$  over the building blocks  $Z_{\pm}$  such that the hypotheses of Theorem 1.7 are satisfied.

The following theorem provides a spherical  $\mu$ -semistable vector bundle  $\mathcal{E}_{\pm, b}$  with

$$(3.1) \quad \text{rk } \mathcal{E}_{\pm, b} = 2, \quad c_1(\mathcal{E}_{\pm, b}) = B_{\pm} \quad \text{and} \quad c_2(\mathcal{E}_{\pm, b}) = -6$$

with  $B_{\pm}$  as in (2.4) on each non-singular fibre  $\Sigma_{\pm, b} := f_{\pm}^{-1}(b)$ .

**Theorem 3.2** (Kuleshov [Kul90, Theorem 2.1]). *Let  $(\Sigma, A)$  be a polarised smooth  $K3$  surface. If  $(r, c_1, c_2) \in \mathbf{N} \times H^{1,1}(\Sigma, \mathbf{Z}) \times \mathbf{Z}$  are such that*

$$(3.3) \quad 2rc_2 - (r-1)c_1^2 - 2(r^2-1) = 0,^5$$

*then there exists a spherical  $\mu$ -semistable vector bundle  $\mathcal{E}$  on  $\Sigma$  with*

$$\text{rk } \mathcal{E} = r, \quad c_1(\mathcal{E}) = c_1 \quad \text{and} \quad c_2(\mathcal{E}) = c_2.$$

*Remark 3.4.* By Hirzebruch–Riemann–Roch, (3.3) is equivalent to  $\chi(\mathcal{E}\text{nd}_0(\mathcal{E})) = 0$ , a necessary condition for  $\mathcal{E}$  to be spherical.

Set

$$U_{\pm} := \{b \in \mathbf{P}^1 : \Sigma_{\pm, b} \text{ is non-singular and } \text{Pic}(\Sigma_{\pm, b}) \cong N_{\pm}\}.$$

---

<sup>5</sup>Here and in the following, for  $x \in H^2(\Sigma)$ , we write  $x^2 \in \mathbf{Z}$  to denote  $x \cup x \in H^4(\Sigma) \cong \mathbf{Z}$ .

Since  $A_{\pm}^{\perp} \subset N_{\pm}$  is generated by  $B_{\pm}$  and  $B_{\pm}^2 = -30 < -6$ , for  $b \in U_{\pm}$  the following guarantees that  $\mathcal{E}_{\pm, b}$  is indeed  $\mu$ -stable (and thus stable<sup>6</sup>).

**Proposition 3.5.** *In the situation of Theorem 3.2, if the divisibilities of  $r$  and  $c_1$  are coprime and for all non-zero  $x \in H^{1,1}(\Sigma, \mathbf{Z})$  perpendicular to  $c_1(A)$  we have*

$$(3.6) \quad x^2 < -\frac{r^2(r^2 - 1)}{2},$$

then  $\mathcal{E}$  is  $\mu$ -stable.

*Proof.* Suppose  $\mathcal{F}$  were a destabilising sheaf, i.e., a torsion-free subsheaf  $\mathcal{F} \subset \mathcal{E}$  with  $0 < \text{rk } \mathcal{F} < \text{rk } \mathcal{E}$  and  $\mu(\mathcal{F}) = \mu(\mathcal{E})$ . Since  $c_1(\mathcal{E})c_1(A) = \text{rk } \mathcal{E} \cdot \mu(\mathcal{E})$  (and similarly for  $\mathcal{F}$ ),  $x := \text{rk } \mathcal{E} \cdot c_1(\mathcal{F}) - \text{rk } \mathcal{F} \cdot c_1(\mathcal{E}) \in c_1(A)^{\perp}$ . The discriminant of  $\mathcal{E}$  is

$$\Delta(\mathcal{E}) := 2 \text{rk } \mathcal{E} \cdot c_2(\mathcal{E}) - (\text{rk } \mathcal{E} - 1)c_1(\mathcal{E})^2 = 2(r^2 - 1)$$

by (3.3). According to [HL10, Theorem 4.C.3] we must have either

$$-\frac{(\text{rk } \mathcal{E})^2}{4} \Delta(\mathcal{E}) \leq x^2,$$

which violates (3.6), or  $x = 0$ .

The latter, however, implies

$$\text{rk } \mathcal{E} \cdot c_1(\mathcal{F}) = \text{rk } \mathcal{F} \cdot c_1(\mathcal{E}),$$

which is impossible because the divisibilities of  $\text{rk } \mathcal{E}$  and  $c_1(\mathcal{E})$  are coprime.  $\square$

As a consequence of this and the following, for  $b \in U_{\pm}$  the moduli space of semistable bundles on  $\Sigma_{\pm, b}$  satisfying (3.1) is a reduced point.

**Theorem 3.7** (Mukai [HL10, Theorem 6.1.6]). *Let  $(\Sigma, A)$  be a polarised smooth K3 surface. Suppose that  $\mathcal{E}$  is a stable sheaf satisfying (3.3) with  $r = \text{rk } \mathcal{E}$ ,  $c_1 = c_1(\mathcal{E})$  and  $c_2 = c_2(\mathcal{E})$ . Then  $\mathcal{E}$  is locally free and any other semistable sheaf satisfying the same condition must be isomorphic to  $\mathcal{E}$ .*

If we were able construct holomorphic vector bundles  $\mathcal{E}_{\pm}$  on  $Z_{\pm}$  whose restrictions to the fibres  $\Sigma_{\pm, b}$  with  $b \in U_{\pm}$  agree with  $\mathcal{E}_{\pm, b}$  and which satisfy (1.8), then we could apply Theorem 1.7 and the proof of Theorem 1.11 would be complete. To see this, note that  $\infty \in U_{\pm}$  and thus  $\mathcal{E}_{\pm}|_{\Sigma_{\pm, \infty}}$  have the same rank, their characteristic classes are identified by  $\iota^*$  (since  $\iota^*B_- = B_+$  by construction) and both are  $\mu$ -stable. The construction of  $\mathcal{E}_{\pm}$  is achieved using the following tool. (Note that  $\frac{1}{2}B_{\pm}^2 + 6 = -9$ ; hence, (3.9) holds in our situation in view of (3.1).)

---

<sup>6</sup>Recall that a torsion-free coherent sheaf  $\mathcal{E}$  on a projective variety  $(X, \mathcal{O}(1))$  is called **(semi)stable** if for each torsion-free coherent subsheaf  $\mathcal{F} \subset \mathcal{E}$  with  $0 < \text{rk } \mathcal{F} < \text{rk } \mathcal{E}$  we have  $(p_{\mathcal{F}} \leq p_{\mathcal{E}}) p_{\mathcal{F}} < p_{\mathcal{E}}$ . Here  $p_{\mathcal{E}}$  denotes the **reduced Hilbert polynomial** of  $\mathcal{E}$ , the unique polynomial satisfying  $p_{\mathcal{E}}(m) = \chi(\mathcal{E} \otimes \mathcal{O}(m)) / \text{rk } \mathcal{E}$  for all  $m \in \mathbf{Z}$ , and we compare polynomials using the lexicographical order of their coefficients.

The notions of  $\mu$ -stability and stability are closely related in case  $(X, \mathcal{O}(1))$  is smooth (and thus Kähler): because  $p_{\mathcal{E}}(m) = \deg \mathcal{O}(1) / n! \cdot m^n + (\mu(\mathcal{E}) + \frac{1}{2} \deg(K_X)) / (n-1)! \cdot m^{n-1} + \dots$ ,  $\mu$ -stable implies stable (and semistable implies  $\mu$ -semistable).

**Proposition 3.8.** *Let  $f: Z \rightarrow B$  be RDP K3 fibration from a projective 3-fold  $Z$  to a smooth curve  $B$  and set  $S := \{b \in B : \Sigma_b := f^{-1}(b) \text{ is singular}\}$ . Let  $(r, c_1, c_2) \in \mathbf{N} \times \text{im}(\text{res}: H^2(Z) \rightarrow H^2(\Sigma_b)) \times \mathbf{Z}$  for some  $b \notin S$  be such that (3.3) holds and*

$$(3.9) \quad \gcd\left(r, \frac{1}{2}c_1^2 - c_2\right) = 1.$$

*Suppose that there is a set  $U \subset B \setminus S$  whose complement is countable and for each  $b \in U$  the moduli space  $M_b$  of semistable bundles  $\mathcal{E}_b$  on  $\Sigma_b$  with*

$$(3.10) \quad \text{rk } \mathcal{E}_b = r, \quad c_1(\mathcal{E}_b) = c_1 \quad \text{and} \quad c_2(\mathcal{E}_b) = c_2$$

*consists of a single reduced point:  $M_b = \{[\mathcal{E}_b]\}$ . Then there exists a holomorphic vector bundle  $\mathcal{E}$  over  $Z$  such that, for all  $b \in U$ ,  $\mathcal{E}|_{\Sigma_b} \cong \mathcal{E}_b$ .  $\mathcal{E}$  is spherical, i.e.,  $H^*(\mathcal{E}\text{nd}_0(\mathcal{E})) = 0$  and unique up to twisting by a line bundle pulled-back from  $B$ .*

*Remark 3.11.* Note that by Hirzebruch–Riemann–Roch  $\chi(\mathcal{E}_b) = \frac{1}{2}c_1^2 - c_2 + 2 \text{rk } \mathcal{E}_b$ , so (3.9) is asking that  $\text{rk } \mathcal{E}_b$  and  $\chi(\mathcal{E}_b)$  be coprime.

This result is essentially contained in Thomas’ work on sheaves on K3 fibrations [Tho00, Theorem 4.5]. Its proof heavily relies on the following generalisation of Theorem 3.7.

**Theorem 3.12** (Thomas [Tho00, Proof of Theorem 4.5]). *Let  $(\Sigma, A)$  be a polarised K3 surface with at worst RDP singularities. If  $\mathcal{E}$  is a stable coherent sheaf on  $\Sigma$  with  $\chi(\mathcal{E}\text{nd}_0(\mathcal{E})) = 0$ , then  $\mathcal{E}$  is locally free.*

We also use the following simple observation.

**Proposition 3.13.** *If  $\mathcal{E}$  is a semistable sheaf with  $\text{rk } \mathcal{E}$  and  $\chi(\mathcal{E})$  coprime, then  $\mathcal{E}$  is stable.*

*Proof.* If  $\mathcal{E}$  is destabilised by  $\mathcal{F} \subset \mathcal{E}$  with  $0 < \text{rk } \mathcal{F} < \text{rk } \mathcal{E}$ , then  $p_{\mathcal{F}} = p_{\mathcal{E}}$ . In particular, evaluating at  $m = 0$  we have

$$\text{rk } \mathcal{E} \cdot \chi(\mathcal{F}) = \text{rk } \mathcal{F} \cdot \chi(\mathcal{E}).$$

This contradicts  $\text{rk } \mathcal{E}$  and  $\chi(\mathcal{E})$  being coprime. □

*Proof of Proposition 3.8.* Consider the moduli functor  $\underline{\mathbf{M}}: \text{Sch}_B^{\text{op}} \rightarrow \text{Set}$  which assigns to a  $B$ -scheme  $U$  the set

$$\underline{\mathbf{M}}(U) := \{\mathcal{E} \text{ a coherent sheaf over } Z \times_B U \text{ satisfying } (\clubsuit)\} / \sim .$$

Here  $(\clubsuit)$  means that  $\mathcal{E}$  is flat over  $U$ , for each  $b \in U$ ,  $\mathcal{E} \otimes_{\mathcal{O}_U} k(b)$  is semistable and its Hilbert polynomial  $P$  agrees with that of a sheaf on a smooth fibre with characteristic classes given by (3.10). We write  $\mathcal{E} \sim \mathcal{F}$  if and only if there exists a line bundle  $\mathcal{L}$  over  $U$  such that  $\mathcal{E}$  and  $\mathcal{F} \otimes \mathcal{L}$  are  $S$ -equivalent; cf. Maruyama [Mar78, p. 561] and Huybrechts and Lehn [HL10, Section 4.1].

$\underline{\mathbf{M}}$  is universally corepresented by a proper and separated  $B$ -scheme  $\mathbf{M}$ , i.e., the moduli problem has a proper and separated coarse moduli space, see Simpson [Sim94, Section 1]. The fibre of  $\mathbf{M}$  over  $b \in B$  is the coarse moduli space of semistable sheaves on  $\Sigma_b$  with Hilbert polynomial  $P$ .

Denote by  $M$  the component of  $\mathbf{M}$  whose fibres over  $B \setminus S$  are the coarse moduli space  $M_b$  of semistable sheaves  $\mathcal{E}$  on  $\Sigma_b$  satisfying (3.10). By assumption, for each  $b \in U$ ,  $M_b$  consists of a single reduced point  $[\mathcal{E}_b]$ . By (3.9) and Proposition 3.13,  $\mathcal{E}_b$  is stable and, hence, spherical because  $\chi(\mathcal{E}nd_0(\mathcal{E}_b)) = 0$ . By Theorem 3.7 it is locally free. Using deformation theory, see, e.g., Hartshorne [Har10, Section 7], one can show that  $M \rightarrow B$  is surjective onto a open neighbourhood of each  $b \in U$  and thus to all of  $B$ , since it is proper. Using (3.9) and Proposition 3.13 as well as Theorem 3.7 again we see that for each  $b \notin S$  the fibre  $M_b$  is a reduced point. Since  $M$  is separated, it follows that  $M = B$ .

By [HL10, Corollary 4.6.7], (3.9) guarantees the existence a universal sheaf  $\mathcal{E}$  on  $Z \times_B M = Z$ .<sup>7</sup> By flatness, for each  $b \in B$ ,  $\chi(\mathcal{E}|_{\Sigma_b}) = \frac{1}{2}c_1^2 - c_2 + 2r$  and  $\chi(\mathcal{E}nd_0(\mathcal{E}|_{\Sigma_b})) = 0$ . From (3.9), Proposition 3.13 and Theorem 3.12 (resp. Theorem 3.7) it follows that  $\mathcal{E}|_{\Sigma_b}$  is locally free and spherical for arbitrary  $b \in B$ . Therefore,  $\mathcal{E}$  is also locally free by [Sim94, Lemma 1.27] and spherical by Grothendieck's spectral sequence.

The asserted uniqueness property follows from the fact that  $\mathcal{E}$  is a universal sheaf and the definition of the moduli functor.  $\square$

This completes the construction of the bundles  $\mathcal{E}_{\pm}$  and thus the proof of Theorem 1.11. Clearly, the above argument also proves the following more abstract result.

**Theorem 3.14.** *Let  $(Z_{\pm}, \Sigma_{\pm}, \omega_{\pm}; r)$  be a matched pair of framed building blocks. Suppose that  $f_{\pm}: Z_{\pm} \rightarrow \mathbf{P}^1$  are RDP K3 fibrations and that for all but countably many  $b \in \mathbf{P}^1$  we have*

$$\text{Pic}(f_{\pm}^{-1}(b)) \cong N_{\pm} := \text{im}(\text{res}_{\pm}: H^2(Z_{\pm}) \rightarrow H^2(\Sigma_{\pm})).$$

Suppose there exists a  $(r, c_1, c_2) \in \mathbf{N} \times (N_+ \cap r^*N_-) \times \mathbf{Z}$  such that

$$2rc_2 - (r-1)c_1^2 - 2(r^2-1) = 0$$

and

$$\gcd\left(r, \frac{1}{2}c_1^2 - c_2\right) = 1.$$

If  $[\omega_{I, \pm}] \in H^2(\Sigma_{\pm}, \mathbf{Q})$  and for all non-zero  $x \in [\omega_{I, \pm}]^{\perp} \subset N_{\pm}$  we have

$$x^2 < -\frac{r^2(r^2-1)}{2},$$

then there exists rank  $r$  holomorphic vector bundles  $\mathcal{E}_{\pm}$  on  $Z_{\pm}$  with

$$c_1(\mathcal{E}_+|_{\Sigma_+}) = r^*c_1(\mathcal{E}_-|_{\Sigma_-}) = c_1 \quad \text{and} \quad c_2(\mathcal{E}_+|_{\Sigma_+}) = r^*c_2(\mathcal{E}_-|_{\Sigma_-}) = c_2$$

satisfying the hypotheses of Theorem 1.7.

<sup>7</sup>Strictly speaking, the quoted result only provides the universal sheaf over  $f^{-1}(B \setminus S)$ ; however, the argument of [Thoo0, second paragraph in the proof of Theorem 4.5] shows why the argument works uniformly on  $B$ . Alternatively, the existence of the universal sheaf can be deduced from [Sim94, Theorem 1.21] and Tsen's theorem  $H_{\text{et}}^2(B, \mathcal{O}^*) = 0$ .

## References

- [CHNP13] A. Corti, M. Haskins, J. Nordström, and T. Pacini. *Asymptotically cylindrical Calabi–Yau 3–folds from weak Fano 3–folds*. *Geometry and Topology* 17.4 (2013), pp. 1955–2059. DOI: 10.2140/gt.2013.17.1955. MR: 3109862. Zbl: 1273.14081 (cit. on p. 6).
- [CHNP15] A. Corti, M. Haskins, J. Nordström, and T. Pacini.  *$G_2$ –manifolds and associative submanifolds via semi-Fano 3–folds*. *Duke Mathematical Journal* 164.10 (2015), pp. 1971–2092. DOI: 10.1215/00127094-3120743. MR: 3369307. Zbl: 06486366 (cit. on pp. 2–5).
- [CN14] D. Crowley and J. Nordström. *Exotic  $G_2$ –manifolds*. 2014. arXiv: 1411.0656 (cit. on pp. 1, 3, 4, 6).
- [Don85] S. K. Donaldson. *Anti self-dual Yang–Mills connections over complex algebraic surfaces and stable vector bundles*. *Proceedings of the London Mathematical Society* 50.1 (1985), pp. 1–26. DOI: 10.1112/plms/s3-50.1.1. MR: 765366. Zbl: 0529.53018 (cit. on p. 3).
- [DS11] S. K. Donaldson and E. P. Segal. *Gauge theory in higher dimensions, II. Surveys in differential geometry. Volume XVI. Geometry of special holonomy and related topics*. Vol. 16. 2011, pp. 1–41. arXiv: 0902.3239. MR: 2893675. Zbl: 1256.53038 (cit. on p. 1).
- [DT98] S. K. Donaldson and R. P. Thomas. *Gauge theory in higher dimensions. The geometric universe (Oxford, 1996)*. Oxford, 1998, pp. 31–47. MR: MR1634503. Zbl: 0926.58003. URL: <http://www.ma.ic.ac.uk/~rpwt/skd.pdf> (cit. on p. 1).
- [Har10] R. Hartshorne. *Deformation theory*. Vol. 257. Graduate Texts in Mathematics. 2010, pp. viii+234. DOI: 10.1007/978-1-4419-1596-2. MR: 2583634. Zbl: 1186.14004 (cit. on p. 10).
- [HHN15] M. Haskins, H.-J. Hein, and J. Nordström. *Asymptotically cylindrical Calabi–Yau manifolds*. *Journal of Differential Geometry* 101.2 (2015), pp. 213–265. DOI: 10.4310/jdg/1442364651. MR: 3399097. Zbl: 1332.32028 (cit. on p. 2).
- [HL10] D. Huybrechts and M. Lehn. *The geometry of moduli spaces of sheaves*. Second. Cambridge Mathematical Library. 2010, pp. xviii+325. DOI: 10.1017/CB09780511711985. MR: 2665168. Zbl: 1206.14027 (cit. on pp. 8–10).
- [HW15] A. Haydys and T. Walpuski. *A compactness theorem for the Seiberg–Witten equation with multiple spinors in dimension three*. *Geometric and Functional Analysis* 25.6 (2015), pp. 1799–1821. DOI: 10.1007/s00039-015-0346-3. arXiv: 1406.5683. MR: 3432158. Zbl: 1334.53039 (cit. on p. 1).
- [IP99] V. A. Iskovskih and Yu. G. Prokhorov. *Fano varieties. Algebraic geometry, V*. Vol. 47. Encyclopaedia Math. Sci. 1999, pp. 1–247. MR: 1668579 (cit. on p. 6).

- [Isk78] V. A. Iskovskih. *Fano threefolds. II. Izv. Akad. Nauk SSSR Ser. Mat.* 42.3 (1978), pp. 506–549. MR: 503430 (cit. on p. 6).
- [Joy96a] D. D. Joyce. *Compact Riemannian 7–manifolds with holonomy  $G_2$ . I. Journal of Differential Geometry* 43.2 (1996), pp. 291–328. DOI: 10.4310/jdg/1214458109. MR: MR1424428. Zbl: 0861.53022 (cit. on p. 4).
- [Joy96b] D. D. Joyce. *Compact Riemannian 7–manifolds with holonomy  $G_2$ . II. Journal of Differential Geometry* 43.2 (1996), pp. 329–375. DOI: 10.4310/jdg/1214458110. MR: MR1424428. Zbl: 0861.53023 (cit. on p. 4).
- [KL11] A. Kovalev and N.-H. Lee.  *$K_3$  surfaces with non-symplectic involution and compact irreducible  $G_2$ –manifolds. Mathematical Proceedings of the Cambridge Philosophical Society* 151.2 (2011), pp. 193–218. DOI: 10.1017/S030500411100003X. MR: 2823130. Zbl: 1228.53064 (cit. on pp. 2, 3).
- [Kov03] A. Kovalev. *Twisted connected sums and special Riemannian holonomy. Journal für die Reine und Angewandte Mathematik* 565 (2003), pp. 125–160. DOI: 10.1515/crll.2003.097. MR: MR2024648. Zbl: 1043.53041 (cit. on pp. 2–4).
- [Kov09] A. Kovalev. *Coassociative  $K_3$  fibrations of compact  $G_2$ –manifolds.* 2009. arXiv: math/0511150v2 (cit. on p. 5).
- [Kul90] S.A. Kuleshov. *An existence theorem for exceptional bundles on  $K_3$  surfaces. Math. USSR, Izv.* 34.2 (1990), pp. 373–388. DOI: 10.1070/IM1990v034n02ABEH001316. MR: 998301. Zbl: 0706.14009 (cit. on p. 7).
- [Mar78] M. Maruyama. *Moduli of stable sheaves. II. J. Math. Kyoto Univ.* 18.3 (1978), pp. 557–614. MR: 509499. Zbl: 0395.14006 (cit. on p. 9).
- [MM81] S. Mori and S. Mukai. *Classification of Fano 3–folds with  $B_2 \geq 2$ . Manuscripta Math.* 36.2 (1981), pp. 147–162. DOI: 10.1007/BF01170131. MR: 641971. Zbl: 0478.14033 (cit. on pp. 4, 6).
- [MNS17] G. Menet, J. Nordström, and H. N. Sá Earp. *Construction of  $G_2$ –instantons via twisted connected sums.* 2017. arXiv: 1510.03836 (cit. on p. 4).
- [Moř67] B. G. Mořezon. *Algebraic homology classes on algebraic varieties. Izv. Akad. Nauk SSSR Ser. Mat.* 31 (1967), pp. 225–268. MR: 0213351. Zbl: 0162.52503 (cit. on pp. 1, 5).
- [Nik79] V. V. Nikulin. *Integer symmetric bilinear forms and some of their geometric applications. Izv. Akad. Nauk SSSR Ser. Mat.* 43.1 (1979), pp. 111–177, 238. MR: 525944. Zbl: 0408.10011 (cit. on p. 7).
- [Sá 15] H. N. Sá Earp.  *$G_2$ –instantons over asymptotically cylindrical manifolds. Geometry and Topology* 19.1 (2015), pp. 61–111. DOI: 10.2140/gt.2015.19.61. MR: 3318748. Zbl: 1312.53044 (cit. on p. 2).

- [Sim94] C. T. Simpson. *Moduli of representations of the fundamental group of a smooth projective variety. I.* *Inst. Hautes Études Sci. Publ. Math.* 79 (1994), pp. 47–129. MR: 1307297. Zbl: 0891.14005. URL: [http://www.numdam.org/item?id=PMIHES\\_1994\\_\\_79\\_\\_47\\_0](http://www.numdam.org/item?id=PMIHES_1994__79__47_0) (cit. on p. 10).
- [Šok79] V. V. Šokurov. *Smoothness of a general anticanonical divisor on a Fano 3–fold.* *Izv. Akad. Nauk SSSR Ser. Mat.* 43.2 (1979), pp. 430–441. MR: 534602. Zbl: 0429.14012 (cit. on p. 4).
- [SW15] H. N. Sá Earp and T. Walpuski.  *$G_2$ –instantons over twisted connected sums.* *Geometry and Topology* 19.3 (2015), pp. 1263–1285. DOI: 10.2140/gt.2015.19.1263. arXiv: 1310.7933. MR: 3352236. Zbl: 06441803 (cit. on pp. 1–4).
- [Tho00] R. P. Thomas. *A holomorphic Casson invariant for Calabi–Yau 3–folds, and bundles on  $K3$  fibrations.* *Journal of Differential Geometry* 54.2 (2000), pp. 367–438. MR: 1818182. Zbl: 1034.14015. URL: <http://projecteuclid.org/euclid.jdg/1214341649> (cit. on pp. 9, 10).
- [Voio7] C. Voisin. *Hodge theory and complex algebraic geometry. II.* English. Vol. 77. Cambridge Studies in Advanced Mathematics. Translated from the French by Leila Schneps. 2007, pp. x+351. MR: 2449178. Zbl: 1129.14020 (cit. on p. 5).
- [Wal13a] T. Walpuski.  *$G_2$ –instantons on generalised Kummer constructions.* *Geometry and Topology* 17.4 (2013), pp. 2345–2388. DOI: 10.2140/gt.2013.17.2345. arXiv: 1109.6609. MR: 3110581. Zbl: 1278.53051 (cit. on p. 4).
- [Wal13b] T. Walpuski. *Gauge theory on  $G_2$ –manifolds.* Imperial College London, 2013. URL: <https://spiral.imperial.ac.uk/bitstream/10044/1/14365/1/Walpuski-T-2013-PhD-Thesis.pdf> (cit. on p. 1).
- [Wal17] T. Walpuski.  *$G_2$ –instantons, associative submanifolds, and Fueter sections.* *Communications in Analysis and Geometry* 25.4 (2017), pp. 847–893. DOI: 10.4310/CAG.2017.v25.n4.a4. arXiv: 1205.5350. MR: 3731643. Zbl: 06823232 (cit. on p. 1).