

Deformation theory of the blown-up Seiberg–Witten equation in dimension three

Aleksander Doan Thomas Walpuski

2018-09-26

Abstract

Associated with every quaternionic representation of a compact, connected Lie group there is a Seiberg–Witten equation in dimension three. The moduli spaces of solutions to these equations are typically non-compact. We construct Kuranishi models around boundary points of a partially compactified moduli space. The Haydys correspondence identifies such boundary points with Fueter sections—solutions of a non-linear Dirac equation—of the bundle of hyperkähler quotients associated with the quaternionic representation. We discuss when such a Fueter section can be deformed to a solution of the Seiberg–Witten equation.

1 Introduction

Associated with every quaternionic representation of a compact, connected Lie group there is a system of partial differential equations generalizing the classical Seiberg–Witten equations in dimension three and four; see, for example, Taubes [Tau99], Pidstrigach [Pido4], Haydys [Hay08], Salamon [Sal13, Section 6], and Nakajima [Nak16, Section 6(i)]. In fact, almost every equation studied in mathematical gauge theory arises in this way. In the present paper we focus on the 3-dimensional theory. A key difficulty in studying Seiberg–Witten equations arises from the non-compactness issue caused by a lack of a priori bounds on the spinor. This phenomenon has been studied in special cases by Taubes [Tau13a; Tau13b; Tau16], and Haydys and Walpuski [HW15]. To focus on the issue of the spinor becoming very large, one passes to a blown-up Seiberg–Witten equation. The lack of a priori bounds then manifests itself as the equation becoming degenerate elliptic when the norm of the spinor tends to infinity. However, the Haydys correspondence allows us to reinterpret the limiting equation as a non-linear version of the Dirac equation, known as the Fueter equation [Sal13; Hay14]. This suggests that, although formally the blown-up Seiberg–Witten equation appears to be degenerate, one should be able to develop an elliptic deformation theory around points at infinity of the moduli space. This is what is achieved in the current paper; the main result being [Theorem 2.29](#) below.

Our second result, [Theorem 2.31](#), asserts that, under a transversality assumption, Fueter sections cause wall-crossing for the signed count of solutions to the Seiberg–Witten equation—a

new phenomenon which has no analog in classical Seiberg–Witten theory. In [DW18] we analyze this wall-crossing phenomenon for the Seiberg–Witten equation with two spinors in detail.

Donaldson and Segal [DS11] proposed that there should be a similar wall-crossing phenomenon for the signed count of G_2 –instantons over a G_2 –manifold. The number of G_2 –instantons jumps due to the appearance of Fueter sections supported on 3–dimensional associative submanifolds of the G_2 –manifold. This is the basis of the conjectural relationship between Seiberg–Witten equations on 3–manifolds and enumerative theories for associative submanifolds and G_2 –instantons. Donaldson and Segal’s prediction was partially confirmed in [Wal17a]; our Theorem 2.31 can be understood as a Seiberg–Witten analog of this result.

Acknowledgements. This material is based upon work supported by the National Science Foundation under Grant No. 1754967 and the Simons Collaboration Grant on “Special Holonomy in Geometry, Analysis and Physics”.

2 Main results

For the reader’s convenience, before stating our main results, we begin by reviewing the necessary background on Seiberg–Witten equations associated with quaternionic representations.

2.1 Hyperkähler quotients of quaternionic vector spaces

Definition 2.1. A quaternionic Hermitian vector space is a real vector space S together with a linear map $\gamma: \text{Im } \mathbf{H} \rightarrow \text{End}(S)$ and an inner product $\langle \cdot, \cdot \rangle$ such that γ makes S into a left module over the quaternions $\mathbf{H} = \mathbf{R}\langle 1, i, j, k \rangle$, and i, j, k act by isometries. The unitary symplectic group $\text{Sp}(S)$ is the subgroup of $\text{GL}(S)$ preserving γ and $\langle \cdot, \cdot \rangle$.

Let G be a compact, connected Lie group.

Definition 2.2. A quaternionic representation of G is a Lie group homomorphism $\rho: G \rightarrow \text{Sp}(S)$ for some quaternionic Hermitian vector space S .

Suppose that a quaternionic representation $\rho: G \rightarrow \text{Sp}(S)$ has been fixed. By slight abuse of notation, we also denote the induced Lie algebra representation by $\rho: \mathfrak{g} \rightarrow \mathfrak{sp}(V)$. We combine ρ and γ into the map $\bar{\gamma}: \mathfrak{g} \otimes \text{Im } \mathbf{H} \rightarrow \text{End}(S)$ defined by

$$\bar{\gamma}(\xi \otimes v)\Phi := \rho(\xi)\gamma(v)\Phi.$$

The map $\bar{\gamma}$ takes values in symmetric endomorphisms of S . Denote the adjoint of $\bar{\gamma}$ by $\bar{\gamma}^*: \text{End}(S) \rightarrow (\mathfrak{g} \otimes \text{Im } \mathbf{H})^*$.

Proposition 2.3. The map $\mu: S \rightarrow (\mathfrak{g} \otimes \text{Im } \mathbf{H})^*$ defined by

$$(2.4) \quad \mu(\Phi) := \frac{1}{2}\bar{\gamma}^*(\Phi\Phi^*)$$

with $\Phi^* := \langle \Phi, \cdot \rangle$ is a **hyperkähler moment map**, that is, it is G -equivariant, and

$$\langle (d\mu)_\Phi \phi, \xi \otimes v \rangle = \langle \gamma(v) \rho(\xi) \Phi, \phi \rangle$$

for all $\xi \in \mathfrak{g}$ and $v \in \text{Im } \mathbf{H}$.

This is a straightforward calculation. Nevertheless, it leads to an important conclusion: there is a hyperkähler orbifold naturally associated with the quaternionic representation.

Definition 2.5. We call $\Phi \in S$ **regular** if $(d\mu)_\Phi : T_\Phi S \rightarrow (\mathfrak{g} \otimes \text{Im } \mathbf{H})^*$ is surjective. Denote by S^{reg} the open cone of regular elements of S .

By Hitchin, Karlhede, Lindström, and Roček [HKLR87, Section 3(D)],

$$X := S^{\text{reg}} // G := (\mu^{-1}(0) \cap S^{\text{reg}}) / G$$

is a hyperkähler orbifold; see also Proposition 4.2. For psychological convenience, we want to make the assumption that X is, in fact, a hyperkähler manifold. It will be important later that X is a cone; that is, it carries a free \mathbf{R}^+ -action.

The following summarizes the algebraic data required to write a Seiberg–Witten equation.

Definition 2.6. A set of **algebraic data** consists of:

1. a quaternionic Hermitian vector space $(S, \gamma, \langle \cdot, \cdot \rangle)$,
2. a compact Lie group H and a closed, connected, normal subgroup $G \triangleleft H$ such that G acts freely on $\mu^{-1}(0) \cap S^{\text{reg}}$,
3. an Ad-invariant inner product on $\text{Lie}(H)$, and
4. a quaternionic representation $\rho : H \rightarrow \text{Sp}(S)$.

Definition 2.7. Given a set of algebraic data as in Definition 2.6, the group $K := H/G$ is called the **flavor symmetry group**.

The groups G and K play different roles: G is the structure group of the equation, whereas K consists of any additional symmetries, which can be used to twist the setup or remain as symmetries of the theory. On first reading, the reader should feel free to assume for simplicity that $H = G \times K$, or even that K is trivial.

2.2 The Seiberg–Witten equation

Let M be a closed, connected 3-manifold.

Definition 2.8. A set of **geometric data** on M compatible with a set of algebraic data as in Definition 2.6 consists of:

1. a Riemannian metric g ,
2. a spin structure \mathfrak{s} ,
3. a principal H -bundle $Q \rightarrow M$,¹ and
4. a connection B on the principal K -bundle

$$R := Q \times_H K.$$

Suppose that a set of geometric data as in Definition 2.8 has been fixed. Left-multiplication by unit quaternions defines an action $\theta: \mathrm{Sp}(1) \rightarrow \mathrm{O}(S)$ such that

$$\theta(q)\gamma(v)\Phi = \gamma(\mathrm{Ad}(q)v)\theta(q)\Phi$$

for all $q \in \mathrm{Sp}(1) = \{q \in \mathbf{H} : |q| = 1\}$, $v \in \mathrm{Im} \mathbf{H}$, and $\Phi \in S$. This can be used to construct various bundles and operations as follows.

Definition 2.9. The **spinor bundle** is the vector bundle

$$\mathfrak{S} := (\mathfrak{s} \times Q) \times_{\mathrm{Sp}(1) \times H} S.$$

Since $T^*M \cong \mathfrak{s} \times_{\mathrm{Sp}(1)} \mathrm{Im} \mathbf{H}$, it inherits a **Clifford multiplication** $\gamma: T^*M \rightarrow \mathrm{End}(\mathfrak{S})$.

Definition 2.10. Denote by $\mathcal{A}(Q)$ the space of connections on Q . Set

$$\mathcal{A}_B(Q) := \{A \in \mathcal{A}(Q) : A \text{ induces } B \text{ on } R\}.$$

$\mathcal{A}_B(Q)$ is an affine space modeled on $\Omega^1(M, \mathfrak{g}_P)$ with \mathfrak{g}_P denoting the **adjoint bundle** associated with $\mathrm{Lie}(G)$, that is,

$$\mathfrak{g}_P := Q \times_{\mathrm{Ad}} \mathrm{Lie}(G).²$$

Definition 2.11. Every $A \in \mathcal{A}_B(Q)$ defines a covariant derivative $\nabla_A: \Gamma(\mathfrak{S}) \rightarrow \Omega^1(M, \mathfrak{S})$. The **Dirac operator** associated with A is the linear map $\not{D}_A: \Gamma(\mathfrak{S}) \rightarrow \Gamma(\mathfrak{S})$ defined by

$$\not{D}_A\Phi := \gamma(\nabla_A\Phi).$$

Definition 2.12. The hyperkähler moment map $\mu: S \rightarrow (\mathrm{Im} \mathbf{H} \otimes \mathfrak{g})^*$ induces a map

$$\mu: \mathfrak{S} \rightarrow \Lambda^2 T^*M \otimes \mathfrak{g}_P$$

since $(T^*M)^* \cong \Lambda^2 T^*M$.

¹The following observation is due to Haydys [Hay14, Section 3.1] and becomes important when formulating the Seiberg–Witten equation in dimension four. Suppose there is a homomorphism $\mathbb{Z}_2 \rightarrow Z(H)$ such that the non-unit in \mathbb{Z}_2 acts through ρ as -1 . Set $\hat{H} := (\mathrm{Sp}(1) \times H)/\mathbb{Z}_2$. All of the constructions in Section 2.2 go through with $\mathfrak{s} \times Q$ replaced by a \hat{H} -principal bundle \hat{Q} . In the classical Seiberg–Witten theory, this corresponds to endowing the manifold with a spin^c structure rather than a spin structure and a $U(1)$ -bundle.

² If $H = G \times K$, then the G -bundle P alluded to in this notation does exist. In general, it does not exist but traces of it do—e.g., its adjoint bundle and its gauge group.

Denoting by

$$\varpi: \mathfrak{g}_Q \rightarrow \mathfrak{g}_P$$

the projection induced by $\text{Lie}(H) \rightarrow \text{Lie}(G)$, we are finally in a position to state the equation we wish to study.

Definition 2.13. The **Seiberg–Witten equation** associated with the chosen geometric data is the following system of differential equations for $(\Phi, A) \in \Gamma(\mathfrak{S}) \times \mathcal{A}_B(Q)$:

$$(2.14) \quad \begin{aligned} \not{D}_A \Phi &= 0 \quad \text{and} \\ \varpi F_A &= \mu(\Phi). \end{aligned}$$

Most of the well-known equations of mathematical gauge theory on 3- and 4-manifolds can be obtained as a Seiberg–Witten equation.³

Example 2.15. $S = \mathbf{H}$ and $\rho: \text{U}(1) \rightarrow \mathbf{H}$ acting by right-multiplication with $e^{i\theta}$ leads to the **classical Seiberg–Witten equation** in dimension three.

For further examples, we refer the reader to [Appendix A](#).

The Seiberg–Witten equation is invariant with respect to gauge transformations which preserve the flavor bundle R .

Definition 2.16. The **group of restricted gauge transformations** is

$$\mathcal{G}(P) := \{u \in \mathcal{G}(Q) : u \text{ acts trivially on } R\}.$$

$\mathcal{G}(P)$ is an infinite dimensional Lie group with Lie algebra $\Omega^0(M, \mathfrak{g}_P)$; it acts on $\Gamma(\mathfrak{S}) \times \mathcal{A}_B(Q)$, and preserves the space of solutions of (2.14).

The main object of our study is the space of solutions to (2.14) modulo restricted gauge transformations. This space depends on the geometric data chosen as in [Definition 2.8](#). The topological part of the data, the bundles \mathfrak{s} and H , will be fixed. The remaining parameters of the equations, the metric g and the connection B , will be allowed to vary.

Definition 2.17. Let $\text{Met}(M)$ be the space of Riemannian metrics on M . The **parameter space** is

$$\mathcal{P} := \text{Met}(M) \times \mathcal{A}(R).$$

Definition 2.18. For $\mathfrak{p} = (g, B) \in \mathcal{P}$, the **Seiberg–Witten moduli space** is

$$\mathfrak{M}_{\text{SW}}(\mathfrak{p}) := \left\{ [(\Phi, A)] \in \frac{\Gamma(\mathfrak{S}) \times \mathcal{A}_B(Q)}{\mathcal{G}(P)} : (\Phi, A) \text{ satisfies (2.14) with respect to } g \text{ and } B \right\}.$$

The **universal Seiberg–Witten moduli space** is

$$\mathfrak{M}_{\text{SW}} := \left\{ (\mathfrak{p}, [(\Phi, A)]) \in \mathcal{P} \times \frac{\Gamma(\mathfrak{S}) \times \mathcal{A}(Q)}{\mathcal{G}(P)} : [(\Phi, A)] \in \mathfrak{M}_{\text{SW}}(\mathfrak{p}) \right\}.$$

³In fact, if we allow the Lie groups and the representations to be infinite-dimensional, we can also recover (special cases of) the G_2 - and $\text{Spin}(7)$ -instanton equations [[Hay12](#), Section 4.2].

The Seiberg–Witten moduli spaces are endowed with the quotient topology induced from the C^∞ -topology on the spaces of connections and sections. As we will explain in Section 3, if c_0 is a solution of (2.14) for some $\mathbf{p}_0 \in \mathcal{P}$, then the deformation theory of (2.14) at (\mathbf{p}_0, c_0) is controlled by a differential graded Lie algebra (DGLA). Associated with this DGLA is a formally-self adjoint elliptic operator $L_{\mathbf{p},c}$, which can be understood as a gauge fixed and co-gauge fixed linearization of (2.14). These operators equip \mathfrak{M}_{SW} with a real line bundle $\det L$ such that for each $(\mathbf{p}, [c]) \in \mathfrak{M}_{\text{SW}}$ we have

$$(\det L)_{(\mathbf{p}, [c])} \cong \det \ker L_{\mathbf{p},c} \otimes (\text{coker } L_{\mathbf{p},c})^*.$$

The fact that the operators $L_{\mathbf{p},c}$ are Fredholm allows us to construct finite dimensional models of \mathfrak{M}_{SW} by standard methods.

Proposition 2.19. *If c_0 is a solution of (2.14) for $\mathbf{p}_0 \in \mathcal{P}$ and c_0 is irreducible,⁴ then there is a **Kuranishi model** for a neighborhood of $(\mathbf{p}_0, [c_0]) \in \mathfrak{M}_{\text{SW}}$; that is: there are an open neighborhood of U of $\mathbf{p}_0 \in \mathcal{P}$, finite dimensional vector spaces I and O of the same dimension, an open neighborhood \mathcal{J} of $0 \in I$, a smooth map*

$$\text{ob}: U \times \mathcal{J} \rightarrow O,$$

an open neighborhood V of $(\mathbf{p}_0, [c_0]) \in \mathfrak{M}_{\text{SW}}$, and a homeomorphism

$$\mathfrak{x}: \text{ob}^{-1}(0) \rightarrow V \subset \mathfrak{M}_{\text{SW}},$$

which maps $(\mathbf{p}_0, 0)$ to $(\mathbf{p}_0, [c_0])$ and commutes with the projections to \mathcal{P} . Moreover, for each $(\mathbf{p}, c) \in \text{im } \mathfrak{x}$, there is an exact sequence

$$0 \rightarrow \ker L_{\mathbf{p},c} \rightarrow I \xrightarrow{d_I \text{ob}} O \rightarrow \text{coker } L_{\mathbf{p},c} \rightarrow 0$$

such that the induced maps $\det L_{\mathbf{p},c} \rightarrow \det(I) \otimes \det(O)^$ define an isomorphism of line bundles $\det L \cong \mathfrak{x}_*(\det(I) \otimes \det(O)^*)$ on $\text{im } \mathfrak{x} \subset \mathfrak{M}_{\text{SW}}$.*

2.3 The blown-up equation and the Haydys correspondence

Unless $\mu^{-1}(0) = \{0\}$, the projection map $\mathfrak{M}_{\text{SW}} \rightarrow \mathcal{P}$ is not expected to be proper. This potential non-compactness phenomenon is related to the lack of a priori bounds on Φ for (Φ, A) a solution of (2.14). With this in mind, we blow-up the equation (2.14); cf. [KM07, Section 2.5; HW15, Equation (1.4)].

Definition 2.20. The **blown-up Seiberg–Witten equation** is the following differential equation for $(\varepsilon, \Phi, A) \in [0, \infty) \times \Gamma(\mathfrak{S}) \times \mathcal{A}_B(Q)$:

$$(2.21) \quad \begin{aligned} \mathcal{D}_A \Phi &= 0, \\ \varepsilon^2 \omega F_A &= \mu(\Phi), \quad \text{and} \\ \|\Phi\|_{L^2} &= 1. \end{aligned}$$

⁴We say that c_0 is irreducible if $\Gamma_{c_0} := \{u \in \mathcal{G}(P) : uc_0 = c_0\} = \{\text{id}\}$, see Definition 3.3. There is a natural generalization of Proposition 2.19 to the case when c_0 is reducible. Then Γ_{c_0} acts on U and O and ob can be chosen to be Γ_{c_0} -equivariant, cf. [DK90, Section 4.2.5]. However, in this paper we focus on neighborhoods of infinity of the moduli space, and as we will see those contain only irreducible solutions.

Set

$$\mathfrak{S}^{\text{reg}} := (\mathfrak{s} \times Q) \times_{\text{Sp}(1) \times H} S^{\text{reg}}.$$

Definition 2.22. The partially compactified Seiberg–Witten moduli space is

$$\overline{\mathfrak{M}}_{\text{SW}}(g, B) := \left\{ (\varepsilon, [(\Phi, A)]) \in [0, \infty) \times \frac{\Gamma(\mathfrak{S}) \times \mathcal{A}_B(Q)}{\mathcal{G}(P)} : \begin{array}{l} (\varepsilon, \Phi, A) \text{ satisfies (2.21)} \\ \text{with respect to } g \text{ and } B; \\ \text{if } \varepsilon = 0, \text{ then } \Phi \in \Gamma(\mathfrak{S}^{\text{reg}}) \end{array} \right\}.$$

Likewise, the universal partially compactified Seiberg–Witten moduli space is

$$\overline{\mathfrak{M}}_{\text{SW}} := \left\{ (\mathfrak{p}, \varepsilon, [(\Phi, A)]) \in \mathcal{P} \times [0, \infty) \times \frac{\Gamma(\mathfrak{S}) \times \mathcal{A}(Q)}{\mathcal{G}(P)} : (\varepsilon, [(\Phi, A)]) \in \overline{\mathfrak{M}}_{\text{SW}}(\mathfrak{p}) \right\}.$$

The partially compactified Seiberg–Witten moduli spaces are also naturally topological spaces. The formal boundary of $\overline{\mathfrak{M}}_{\text{SW}}$ is

$$\partial \mathfrak{M}_{\text{SW}} := \left\{ (\mathfrak{p}, 0, [(\Phi, A)]) \in \overline{\mathfrak{M}}_{\text{SW}} \right\},$$

and the map

$$\overline{\mathfrak{M}}_{\text{SW}} \setminus \partial \mathfrak{M}_{\text{SW}} \rightarrow \mathfrak{M}_{\text{SW}}, \quad (\mathfrak{p}, \varepsilon, [(\Phi, A)]) \mapsto (\mathfrak{p}, [(\varepsilon^{-1}\Phi, A)])$$

is a homeomorphism. This justifies the term “partially compactified”.

Warning 2.23. The space $\overline{\mathfrak{M}}_{\text{SW}}(g, B)$ need not be compact. From work of Taubes [Tau13a] on Example A.2 with $G = \text{SO}(3)$ and work of Haydys and Walpuski [HW15] on Example A.3 with $k = 1$, we expect that the actual compactification will also contain singular solutions of (2.21) with $\varepsilon = 0$; see [DW18]. In fact, $\partial \mathfrak{M}_{\text{SW}}$ need not be compact [Wal17b]. Precisely understanding the full compactifications is one of the central challenges in this subject.

For $\varepsilon = 0$, (2.21) appears to be degenerate. However, since $\Phi \in \Gamma(\mathfrak{S}^{\text{reg}})$, this equation can be understood as an elliptic PDE as follows.

Definition 2.24. The bundle of hyperkähler quotients $\pi: \mathfrak{X} \rightarrow M$ is

$$\mathfrak{X} := (\mathfrak{s} \times R) \times_{\text{Sp}(1) \times K} X.$$

Its vertical tangent bundle is

$$V\mathfrak{X} := (\mathfrak{s} \times R) \times_{\text{Sp}(1) \times K} TX,$$

and $\gamma: \text{Im } \mathbf{H} \rightarrow \text{End}(S)$ induces a Clifford multiplication $\gamma: \pi^*TM \rightarrow \text{End}(V\mathfrak{X})$.

Definition 2.25. Using $B \in \mathcal{A}(R)$ we can assign to each $s \in \Gamma(\mathfrak{X})$ its covariant derivative $\nabla_B s \in \Omega^1(M, s^*V\mathfrak{X})$. A section $s \in \Gamma(\mathfrak{X})$ is called a **Fueter section** if it satisfies the **Fueter equation**

$$(2.26) \quad \mathfrak{F}(s) = \mathfrak{F}_B(s) := \gamma(\nabla_B s) = 0 \in \Gamma(s^*V\mathfrak{X}).$$

The map $s \mapsto \mathfrak{F}(s)$ is called the **Fueter operator**.⁵

⁵In the following, we will suppress the subscript B from the notation.

An elementary but important calculation shows that a pair $(\Phi, A) \in \Gamma(\mathfrak{S}^{\text{reg}}) \times \mathcal{A}_B(Q)$ satisfies $\mathbb{D}_A \Phi = 0$ and $\mu(\Phi) = 0$ if and only if the projection $s := p \circ \Phi \in \Gamma(\mathfrak{X})$ satisfies $\mathfrak{F}(s) = 0$. This is part of the Haydys correspondence, which will be discussed in more detail in Section 4.

The linearized Fueter operator $(d\mathfrak{F})_s : \Gamma(s^*V\mathfrak{X}) \rightarrow \Gamma(s^*V\mathfrak{X})$ is a formally self-adjoint elliptic differential operator of order one. In particular, it is Fredholm of index zero. However, the space of solutions to $\mathfrak{F}(s) = 0$, if non-empty, is never zero-dimensional. The reason is that the hyperkähler quotient $X = S^{\text{reg}} // G$ carries a free \mathbf{R}^+ -action inherited from the vector space structure on S . This induces a fiber-preserving action of \mathbf{R}^+ on \mathfrak{X} . One easily verifies that, for $\lambda \in \mathbf{R}^+$ and $s \in \Gamma(\mathfrak{X})$,

$$(2.27) \quad \mathfrak{F}(\lambda s) = \lambda \mathfrak{F}(s).$$

As a result, \mathbf{R}^+ acts freely on the space of solutions to (2.26) which shows that Fueter sections come in one-parameter families. At the infinitesimal level, this shows that every Fueter section is obstructed.

Definition 2.28. The radial vector field $\hat{v} \in \Gamma(\mathfrak{X}, V\mathfrak{X})$ is the vector field generating the \mathbf{R}^+ -action on \mathfrak{X} .

Differentiating (2.27) shows that if s is a Fueter section, then $\hat{v} \circ s \in \Gamma(s^*V\mathfrak{X})$ is a non-zero element of $\ker(d\mathfrak{F})_s$.

2.4 Kuranishi models for $\overline{\mathfrak{M}}_{\text{SW}}$

The main result of this article is the construction of Kuranishi models for $\overline{\mathfrak{M}}_{\text{SW}}$ centered at points of $\partial\mathfrak{M}_{\text{SW}}$.

Theorem 2.29. Let $\mathbf{p}_0 = (g_0, B_0) \in \mathcal{P}$ and $\mathbf{c}_0 = (\Phi_0, A_0) \in \Gamma(\mathfrak{S}^{\text{reg}}) \times \mathcal{A}_B(Q)$ be such that $(\mathbf{p}_0, 0, [\mathbf{c}_0]) \in \partial\mathfrak{M}_{\text{SW}}$. Denote by $s_0 = p \circ \Phi_0 \in \Gamma(\mathfrak{X})$ the corresponding Fueter section of \mathfrak{X} . Set

$$I_\partial := \ker(d\mathfrak{F})_{s_0} \cap (\hat{v} \circ s)^{\perp} \quad \text{and} \quad O := \text{coker}(d\mathfrak{F})_{s_0}.$$

Let $r \in \mathbf{N}$.

There exist an open neighborhood \mathcal{J}_∂ of $0 \in I_\partial$, a constant $\varepsilon_0 > 0$, an open neighborhood $U \subset \mathcal{P}$ of \mathbf{p}_0 , a C^{2r-1} map

$$\text{ob} : U \times [0, \varepsilon_0] \times \mathcal{J}_\partial \rightarrow O,$$

an open neighborhood V of $(\mathbf{p}_0, 0, [\mathbf{c}_0]) \in \overline{\mathfrak{M}}_{\text{SW}}$, and a homeomorphism

$$\mathfrak{x} : \text{ob}^{-1}(0) \rightarrow V \subset \overline{\mathfrak{M}}_{\text{SW}}$$

such that the following hold:

1. There are smooth functions

$$\text{ob}_\partial, \widehat{\text{ob}}_1, \dots, \widehat{\text{ob}}_r : U \times \mathcal{J}_\partial \rightarrow O$$

such that for all $m, n \in \mathbf{N}_0$ with $m + n \leq 2r$ we have

$$\left\| \nabla_{U \times \mathcal{J}_\partial}^m \partial_\varepsilon^n \left(\text{ob} - \text{ob}_\partial - \sum_{i=1}^r \varepsilon^{2i} \widehat{\text{ob}}_i \right) \right\|_{C^0} = O(\varepsilon^{2r-n+2}).$$

2. The map \mathfrak{x} commutes with the projection to $\mathcal{P} \times [0, \infty)$ and satisfies

$$\mathfrak{x}(\mathbf{p}_0, 0, 0) = (\mathbf{p}_0, 0, [c_0]).$$

3. For each $(\mathbf{p}, c) \in \text{im } \mathfrak{x} \cap \mathfrak{M}_{\text{SW}}$, the solution c is irreducible; moreover, it is unobstructed if d_{Job} is surjective.

Remark 2.30. The neighborhoods \mathcal{S}_∂ and U may depend on the choice of r .

The difficulty in proving this theorem arises from the fact that the (gauge fixed and co-gauge fixed) linearization of (2.21) appears to become degenerate as ε approaches zero. The Haydys correspondence, however, indicates that one can reinterpret (2.21) at $\varepsilon = 0$ as the Fueter equation; in particular, as a non-degenerate elliptic PDE. One can think of Theorem 2.29 as a gluing theorem for the Kuranishi model described in Proposition 2.19 with a Kuranishi model for the moduli space of Fueter sections divided by the \mathbf{R}^+ -action.

2.5 Wall-crossing

The main application of the work in this article—and our motivation for it—is to understand wall-crossing phenomena for signed counts of solutions to Seiberg–Witten equations arising from the non-compactness phenomenon mentioned in Section 2.3. In the generic situation of Theorem 2.29, one expects to have $\ker(d\tilde{\mathfrak{F}})_{s_0} = \mathbf{R}\langle \hat{v} \circ s_0 \rangle$. In this case, if $\{\mathbf{p}_t = (g_t, B_t) : t \in (-T, T)\}$ is a 1-parameter family in \mathcal{P} , then (for $T \ll 1$) one can find a 1-parameter family $\{(s_t) \in \Gamma(\mathfrak{X}) : t \in (-T, T)\}$ of sections of \mathfrak{X} and $\lambda : (-T, T) \rightarrow \mathbf{R}$ with $\lambda(0) = 0$ such that

$$\tilde{\mathfrak{F}}_t(s_t) = \lambda(t) \cdot \hat{v} \circ s_t.$$

Theorem 2.31. *In the situation above and assuming $\dot{\lambda}(0) \neq 0$, for each $r \in \mathbf{N}$, there exist $\varepsilon_0 > 0$ and C^{2r-1} maps $t : [0, \varepsilon_0) \rightarrow (-T, T)$ and $c : [0, \varepsilon_0) \rightarrow \Gamma(\mathfrak{S}^{\text{reg}}) \times \mathcal{A}(Q)$ such that an open neighborhood V of $(0, 0, [c_0])$ in the parametrized Seiberg–Witten moduli space*

$$\left\{ (t, \varepsilon, [(\Phi, A)]) \in (-T, T) \times [0, \infty) \times \frac{\Gamma(\mathfrak{S}) \times \mathcal{A}(Q)}{\mathcal{G}(P)} : (\varepsilon, [(\Phi, A)]) \in \overline{\mathfrak{M}}_{\text{SW}}(\mathbf{p}_t) \right\}$$

is given by

$$V = \{(t(\varepsilon), \varepsilon, [c(\varepsilon)]) : \varepsilon \in [0, \varepsilon_0)\}.$$

If $c(\varepsilon) = (\Phi(\varepsilon), A(\varepsilon))$, then there is $\phi \in \Gamma(\mathfrak{S})$ such that

$$\Phi(\varepsilon) = \Phi_0 + \varepsilon^2 \phi + O(\varepsilon^4),$$

and with

$$\delta := \langle \phi, \mathbb{D}_{A_0} \phi \rangle_{L^2}$$

we have

$$t(\varepsilon) = \frac{\delta}{\dot{\lambda}(0)} \varepsilon^4 + O(\varepsilon^6).$$

For $\varepsilon \in (0, \varepsilon_0)$, $c(\varepsilon)$ is irreducible; moreover, if $\delta \neq 0$, then $c(\varepsilon)$ is unobstructed.

Remark 2.32. In the situation of [Theorem 2.31](#), there is no obstruction to solving the Seiberg–Witten equation to order ε^2 —in fact, a solution can be found rather explicitly. The obstruction to solving to order ε^4 is precisely δ .

If \mathfrak{M}_{SW} is oriented (that is: $\det L \rightarrow \mathfrak{M}_{\text{SW}}$ is trivialized) around $(\mathbf{p}_0, [c_0])$, then identifying $\ker(d\mathfrak{F})_{s_0} = \text{coker}(d\mathfrak{F})_{s_0} = \mathbf{R}\langle \hat{v} \circ s \rangle$ determines a sign $\sigma = \pm 1$. If $\delta \neq 0$, then contribution of $[c(\varepsilon)]$ should be counted with sign $-\sigma \cdot \text{sign}(\delta)$; as is discussed in [Section 3.4](#). However, $\text{sign}(\delta/\dot{\lambda}(0))$ also determines whether the solution $c(\varepsilon)$ appears for $t < 0$ or $t > 0$. Thus, the overall contributions from $\text{sign}(\delta)$ cancel.




Figure 1: Two examples of wall-crossing.

This is illustrated in [Figure 1](#), which depicts two examples of wall-crossing. More precisely, it shows the projection of $\bigcup_{t \in (-T, T)} \overline{\mathfrak{M}}_{\text{SW}}(\mathbf{p}_t)$ on the (t, ε) -plane. In both cases we assume $\dot{\lambda}(0) > 0$ and $\sigma = +1$. [Figure 1a](#) represents the case $\delta > 0$, in which a solution $c(\varepsilon)$ with sign $\text{sign}(c(\varepsilon)) = -\sigma \cdot \text{sign}(\delta) = -1$ is born at $t = 0$. [Figure 1b](#) represents the case $\delta < 0$, in which $\text{sign}(c(\varepsilon)) = +1$ and the solution dies at $t = 0$. In both cases, as we cross from $t < 0$ to $t > 0$ the signed count of solutions to the Seiberg–Witten equation changes by -1 .

3 Deformation theory of the Seiberg–Witten equation

We begin with the deformation theory of the blown-up Seiberg–Witten equation away from $\varepsilon = 0$, that is, with the deformation theory of the Seiberg–Witten equation itself. All of this material is standard, but it will set the stage for what is to come.

3.1 The Seiberg–Witten DGLA

The deformation theory of the Seiberg–Witten equation is controlled by the following differential graded Lie algebra (DGLA).

Definition 3.1. Denote by L^\bullet the graded real vector space given by

$$\begin{aligned} L^0 &:= \Omega^0(M, \mathfrak{g}_P), \\ L^1 &:= \Gamma(\mathfrak{S}) \oplus \Omega^1(M, \mathfrak{g}_P), \\ L^2 &:= \Gamma(\mathfrak{S}) \oplus \Omega^2(M, \mathfrak{g}_P), \quad \text{and} \\ L^3 &:= \Omega^3(M, \mathfrak{g}_P). \end{aligned}$$

Denote by $\llbracket \cdot, \cdot \rrbracket : L^\bullet \otimes L^\bullet \rightarrow L^\bullet$ the graded skew-symmetric bilinear map defined by

$$\begin{aligned} \llbracket a, b \rrbracket &:= [a \wedge a] && \text{for } a, b \in \Omega^\bullet(M, \mathfrak{g}_P), \\ \llbracket \xi, \phi \rrbracket &:= \rho(\xi)\phi && \text{for } \xi \in \Omega^0(M, \mathfrak{g}_P) \text{ and } \phi \in \Gamma(\mathfrak{S}) \text{ in degree 1 or 2,} \\ \llbracket a, \phi \rrbracket &:= -\bar{\gamma}(a)\phi && \text{for } a \in \Omega^1(M, \mathfrak{g}_P) \text{ and } \phi \in \Gamma(\mathfrak{S}) \text{ in degree 1,} \\ \llbracket \phi, \psi \rrbracket &:= -2\mu(\phi, \psi) && \text{for } \phi, \psi \in \Gamma(\mathfrak{S}) \text{ in degree 1, and} \\ \llbracket \phi, \psi \rrbracket &:= -*\rho^*(\phi\psi^*) && \text{for } \phi \in \Gamma(\mathfrak{S}) \text{ in degree 1 and } \psi \in \Gamma(\mathfrak{S}) \text{ in degree 2.} \end{aligned}$$

Given $\mathfrak{c} = (\Phi, A) \in \Gamma(\mathfrak{S}) \times \mathcal{A}_B(Q)$, define the degree one linear map $\delta^\bullet = \delta_\mathfrak{c}^\bullet : L^\bullet \rightarrow L^{\bullet+1}$ by

$$\begin{aligned} \delta_\mathfrak{c}^0(\xi) &:= \begin{pmatrix} -\rho(\xi)\Phi \\ d_A\xi \end{pmatrix}, \\ \delta_\mathfrak{c}^1(\phi, a) &:= \begin{pmatrix} -\mathcal{D}_A\phi - \bar{\gamma}(a)\Phi \\ -2\mu(\Phi, \phi) + d_Aa \end{pmatrix}, \quad \text{and} \\ \delta_\mathfrak{c}^2(\psi, b) &:= *\rho^*(\psi\Phi^*) + d_Ab. \end{aligned}$$

Proposition 3.2. *The algebraic structures defined in Definition 3.1 determine a DGLA which controls the deformation theory of the Seiberg–Witten equation; that is:*

1. $(L^\bullet, \llbracket \cdot, \cdot \rrbracket)$ is a graded Lie algebra.
2. If $\mathfrak{c} = (\Phi, A)$ is a solution of (2.14), then $(L^\bullet, \llbracket \cdot, \cdot \rrbracket, \delta_\mathfrak{c}^\bullet)$ is a DGLA.
3. Suppose that $\mathfrak{c} = (\Phi, A)$ is a solution of (2.14). For every $\hat{\mathfrak{c}} = (\phi, a) \in L^1$, $(\Phi + \phi, A + a)$ solves (2.14) if and only if it is a **Maurer–Cartan element**, that is, $\delta_\mathfrak{c}\hat{\mathfrak{c}} + \frac{1}{2}\llbracket \hat{\mathfrak{c}}, \hat{\mathfrak{c}} \rrbracket = 0$.

The verification of (1) and (2) is somewhat lengthy, and is deferred to Appendix C. Part (3), however, is straightforward.

Definition 3.3. Let $\mathfrak{c} \in \Gamma(\mathfrak{S}) \times \mathcal{A}_B(Q)$ be a solution of (2.14). We call

$$\Gamma_\mathfrak{c} := \{u \in \mathcal{G}(P) : uc = \mathfrak{c}\}$$

the group of **automorphisms** of \mathfrak{c} . Its Lie algebra is the cohomology group $H^0(L^\bullet, \delta_\mathfrak{c})$; $H^1(L^\bullet, \delta_\mathfrak{c})$ is the space of **infinitesimal deformations**, and $H^2(L^\bullet, \delta_\mathfrak{c})$ the space of **infinitesimal obstructions**. We say that \mathfrak{c} is **irreducible** if $\Gamma_\mathfrak{c} = 0$, and **unobstructed** if $H^2(L^\bullet, \delta_\mathfrak{c}) = 0$.

Remark 3.4. $H^3(L^\bullet, \delta_\mathfrak{c})$ has no immediate interpretation, but it is isomorphic to $H^0(L^\bullet, \delta_\mathfrak{c})$, since the complex $(L^\bullet, \delta_\mathfrak{c})$ is self-dual (up to signs). The latter also shows that $H^1(L^\bullet, \delta_\mathfrak{c})$ is isomorphic to $H^2(L^\bullet, \delta_\mathfrak{c})$.

3.2 The linearized Seiberg–Witten equation

The operators

$$\begin{aligned}\tilde{\delta}_c^0 &:= \delta_c^0: \Omega^0(M, \mathfrak{g}_P) \rightarrow \Gamma(\mathfrak{S}) \oplus \Omega^1(M, \mathfrak{g}_P), \\ \tilde{\delta}_c^1 &:= (\text{id}_{\mathfrak{S}} \oplus *) \circ \delta_c^1: \Gamma(\mathfrak{S}) \oplus \Omega^1(M, \mathfrak{g}_P) \rightarrow \Gamma(\mathfrak{S}) \oplus \Omega^1(M, \mathfrak{g}_P), \quad \text{and} \\ \tilde{\delta}_c^2 &:= - * \circ \delta_c^2 \circ (\text{id}_{\mathfrak{S}} \oplus *): \Gamma(\mathfrak{S}) \oplus \Omega^1(M, \mathfrak{g}_P) \rightarrow \Omega^0(M, \mathfrak{g}_P)\end{aligned}$$

satisfy

$$(\tilde{\delta}_c^0)^* = \delta_c^2 \quad \text{and} \quad (\delta_c^1)^* = \tilde{\delta}_c^1,$$

and $L_c: \Gamma(\mathfrak{S}) \oplus \Omega^1(M, \mathfrak{g}_P) \oplus \Omega^0(M, \mathfrak{g}_P) \rightarrow \Gamma(\mathfrak{S}) \oplus \Omega^1(M, \mathfrak{g}_P) \oplus \Omega^0(M, \mathfrak{g}_P)$ defined by

$$\begin{aligned}L_c &:= \begin{pmatrix} \tilde{\delta}_c^1 & \tilde{\delta}_c^0 \\ \tilde{\delta}_c^2 & 0 \end{pmatrix} \\ &= \begin{pmatrix} -\not{D}_A & 0 & 0 \\ 0 & *d_A & d_A \\ 0 & d_A^* & 0 \end{pmatrix} + \begin{pmatrix} 0 & -\bar{y}(\cdot)\Phi & -\rho(\cdot)\Phi \\ -2 * \mu(\Phi, \cdot) & 0 & 0 \\ -\rho^*(\cdot)\Phi^* & 0 & 0 \end{pmatrix}\end{aligned}$$

is formally self-adjoint and elliptic.

Definition 3.5. We call L_c the **linearization** of the Seiberg–Witten equation at c .

If c is a solution of (2.14), then Hodge theory identifies $H^1(L^\bullet, \delta_c) \oplus H^0(L^\bullet, \delta_c)$ with $\ker L_c$ and $H^2(L^\bullet, \delta_c) \oplus H^3(L^\bullet, \delta_c)$ with $\text{coker } L_c$. The fact that (L^\bullet, δ_c) is self-dual (up to signs) manifests itself as L_c being formally self-adjoint. After gauge fixing and co-gauge fixing, we can understand (2.14) as an elliptic PDE as follows.

Proposition 3.6. *Given*

$$c_0 = (\Phi_0, A_0) \in \Gamma(\mathfrak{S}) \times \mathcal{A}_B(Q),$$

define $Q: \Gamma(\mathfrak{S}) \oplus \Omega^1(M, \mathfrak{g}_P) \oplus \Omega^0(M, \mathfrak{g}_P) \rightarrow \Gamma(\mathfrak{S}) \oplus \Omega^1(M, \mathfrak{g}_P) \oplus \Omega^0(M, \mathfrak{g}_P)$ by

$$Q(\phi, a, \xi) := \begin{pmatrix} -\bar{y}(a)\phi \\ \frac{1}{2} * [a \wedge a] - * \mu(\phi) \\ 0 \end{pmatrix},$$

$e_{c_0} \in \Gamma(\mathfrak{S}) \oplus \Omega^1(M, \mathfrak{g}_P) \oplus \Omega^0(M, \mathfrak{g}_P)$ by

$$e_{c_0} := \begin{pmatrix} -\not{D}_{A_0} \Phi_0 \\ * \omega F_{A_0} - * \mu(\Phi_0) \\ 0 \end{pmatrix},$$

and set

$$\text{sw}_{c_0}(\hat{c}) := L_{c_0} \hat{c} + Q_{c_0}(\hat{c}) + e_{c_0}.$$

There is a constant $\sigma > 0$ depending on c_0 such that, for every $\hat{c} = (\phi, a, \xi) \in \Gamma(\mathfrak{S}) \oplus \Omega^1(M, \mathfrak{g}_P) \oplus \Omega^0(M, \mathfrak{g}_P)$ satisfying $\|(\phi, a)\|_{L^\infty} < \sigma$, the equation

$$\mathfrak{sw}_{c_0}(\hat{c}) = 0$$

holds if and only if $c_0 + (\phi, a)$ satisfies (2.14) and the gauge fixing condition

$$(3.7) \quad d_{A_0}^* a - \rho^*(\phi\Phi_0^*) = 0$$

as well as

$$d_{A_0}\xi = 0 \quad \text{and} \quad \rho(\xi)\Phi_0 = 0;$$

moreover, if c_0 is infinitesimally irreducible (that is: $H^0(L^\bullet, \delta_{c_0}) = 0$), then $\xi = 0$.

The proof requires a number of useful identities for μ which are summarized and proved in Appendix B.

Proof. Setting $\Phi := \Phi_0 + \phi$ and $A := A_0 + a$, the equation $\mathfrak{sw}_{c_0}(\hat{c}) = 0$ amounts to

$$\begin{aligned} \mathcal{D}_A\Phi + \rho(\xi)\Phi_0 &= 0, \\ \omega F_A + *d_{A_0}\xi &= \mu(\Phi), \quad \text{and} \\ d_{A_0}^* a - \rho^*(\phi\Phi_0^*) &= 0. \end{aligned}$$

Since

$$d_A\mu(\Phi) = - * \rho^*((\mathcal{D}_A\Phi)\Phi^*)$$

by (B.5), applying d_A to the second equation above and using the first equation we obtain

$$d_{A_0}^* d_{A_0}\xi + \rho^*((\rho(\xi)\Phi_0)\Phi_0^*) - *[a \wedge *d_{A_0}\xi] + \rho^*((\rho(\xi)\Phi_0)\phi^*) = 0.$$

Taking the L^2 inner product with ξ_0 , the component of ξ in the L^2 orthogonal complement of $\ker \delta_{c_0}$ and integrating by parts yields that

$$\|d_{A_0}\xi\|_{L^2}^2 + \|\rho(\xi)\Phi_0\|_{L^2}^2 = \langle *[a \wedge *d_{A_0}\xi], \xi_0 \rangle_{L^2} - \langle \rho(\xi)\Phi_0, \rho(\xi_0)\phi \rangle_{L^2}.$$

The right-hand side can be bounded by a constant $c > 0$ (depending on c_0) times

$$\|(a, \phi)\|_{L^\infty} \left(\|d_{A_0}^*\xi\|_{L^2}^2 + \|\rho(\xi)\Phi_0\|_{L^2}^2 \right).$$

Therefore, if $\|(a, \phi)\|_{L^\infty} < \sigma := 1/c$, then

$$d_{A_0}\xi = 0 \quad \text{and} \quad \rho(\xi)\Phi_0 = 0.$$

It follows that $\hat{c} + (\phi, a)$ satisfies (2.14).

Since $\xi \in H^0(L^\bullet, \delta_{c_0})$, it vanishes if c_0 is infinitesimally irreducible. \square

The following standard observation shows that imposing the gauge fixing condition (3.7) is mostly harmless, as long as we are only interested in small variations \hat{c} ; c.f. [DK90, Proposition 4.2.9].

Notation 3.8. In what follows we denote by $W^{k,p}\Gamma(\mathfrak{S})$ the space of sections of \mathfrak{S} of Sobolev class $W^{k,p}$. We use similar notations for spaces of connections, gauge transformations, and differential forms.

Proposition 3.9. Fix $k \in \mathbb{N}$ and $p \in (1, \infty)$ with $(k+1)p > 3$. Given

$$c_0 = (\Phi_0, A_0) \in W^{k+1,p}\Gamma(\mathfrak{S}) \times W^{k+2,p}\mathcal{A}_B(Q),$$

there is a constant $\sigma > 0$ such that if we set

$$\mathfrak{U}_{c_0, \sigma} := \left\{ \hat{c} \in B_\sigma(0) \subset W^{k+1,p}\Gamma(\mathfrak{S}) \times W^{k+2,p}\Omega^1(M, \mathfrak{g}_P) : d_{A_0}^* a - \rho^*(\phi\Phi_0^*) = 0 \right\},$$

then the map

$$\mathfrak{U}_{c_0, \sigma} / \Gamma_{c_0} \ni [\hat{c}] \mapsto [c_0 + \hat{c}] \in \frac{W^{k+1,p}\Gamma(\mathfrak{S}) \times W^{k+2,p}\mathcal{A}_B(Q)}{W^{k+3,p}\mathcal{G}(P)}$$

is a homeomorphism onto its image; moreover, $\Gamma_{c_0 + \hat{c}}$ is the stabilizer of \hat{c} in Γ_c .

For $\hat{c} = (\phi, a, \xi)$ and $(\Phi, A) = c = c_0 + (\phi, a)$, we have

$$(d\mathfrak{sw}_{c_0})_{\hat{c}} = \begin{pmatrix} -\mathcal{D}_A & 0 & 0 \\ 0 & *d_A & d_{A_0} \\ 0 & d_{A_0}^* & 0 \end{pmatrix} + \begin{pmatrix} 0 & -\bar{y}(\cdot)\Phi & -\rho(\cdot)\Phi_0 \\ -2 * \mu(\Phi, \cdot) & 0 & 0 \\ -\rho^*(\cdot)\Phi_0^* & 0 & 0 \end{pmatrix}.$$

In particular, $(d\mathfrak{sw}_{c_0})_0$ agrees with L_{c_0} . The following proposition explains the relation between $(d\mathfrak{sw}_{c_0})_{\hat{c}}$ and L_c for $c = (\Phi, A, 0) + \hat{c}$.

Proposition 3.10. In the situation of Proposition 3.9, if $\hat{c} \in \mathfrak{U}_{c_0, \sigma}$ and $c = c_0 + \hat{c}$, then there is a $\tau > 0$ and a smooth map $\phi_{c_0, c} : B_\tau(c) \rightarrow B_\sigma(0)$ which maps $\mathfrak{U}_{c, \tau}$ to $\mathfrak{U}_{c_0, \sigma}$, commutes with the projection to $(W^{k+1,p}\Gamma(\mathfrak{S}) \times W^{k+2,p}\mathcal{A}_B(Q)) / (W^{k+3,p}\mathcal{G}(P))$, and satisfies

$$(d\phi)_c^{-1} (d\mathfrak{sw}_{c_0})_{\hat{c}} (d\phi)_c = (d\mathfrak{sw}_c)_0 = L_c.$$

3.3 Construction of Kuranishi models

The method of the proof of Proposition 2.19 is quite standard, c.f. [DK90, Section 4.2]. Fix $k \in \mathbb{N}$ and $p \in (1, \infty)$ with $(k+1)p > 3$. Given $\mathfrak{p} = (g, B) \in \mathcal{P}$, set

$$\mathfrak{M}_{\text{SW}}^{k,p}(\mathfrak{p}) := \left\{ [(\Phi, A)] \in \frac{W^{k+1,p}\Gamma(\mathfrak{S}) \times W^{k+2,p}\mathcal{A}_B(Q)}{W^{k+3,p}\mathcal{G}(P)} : \begin{array}{l} (\Phi, A) \text{ satisfies (2.14)} \\ \text{with respect to } g \text{ and } B \end{array} \right\},$$

and define $\mathfrak{M}_{\text{SW}}^{k,p}$ accordingly. It is a consequence of elliptic regularity for L_c and Proposition 3.9, that the inclusion $\mathfrak{M}_{\text{SW}} \subset \mathfrak{M}_{\text{SW}}^{k,p}$ is a homeomorphism. This together with Proposition 3.6 and Proposition 3.9 implies that if $(\mathbf{p}_0, [\hat{c}_0]) \in \mathfrak{M}_{\text{SW}}$ is irreducible, then there is a constant $\sigma > 0$ and an open neighborhood U of $\mathbf{p} \in \mathcal{P}$ such that if $B_\sigma(0)$ denotes the open ball of radius σ centered at 0 in $W^{k+1,p}\Gamma(\mathfrak{S}) \oplus W^{k+2,p}\Omega^1(M, \mathfrak{g}_P) \oplus W^{k+2,p}\Omega^0(M, \mathfrak{g}_P)$, then

$$\{(\mathbf{p}, \hat{c}) \in U \times B_\sigma(0) : \text{sw}_{\mathbf{p}, c_0}(\hat{c}) = 0\} \ni (\mathbf{p}, [(\phi, a, \xi)]) \mapsto (\mathbf{p}, [c + (\phi, a)]) \in \mathfrak{M}_{\text{SW}}$$

is a homeomorphism onto its image. Here we use subscripts to denote the dependence of L_{c_0} , Q , e_{c_0} , and $\text{sw}_{\mathbf{p}, c_0}$ on the parameter $\mathbf{p} \in \mathcal{P}$. The proof of Proposition 2.19 is completed by applying the following result to $\text{sw}_{\mathbf{p}, c_0}$ with $I = \ker L_{\mathbf{p}_0, c_0}$ and $O = \text{coker } L_{\mathbf{p}_0, c_0}$.

Lemma 3.11. *Let X and Y be Banach spaces, let $U \subset X$ be a neighborhood of $0 \in X$, let P be a Banach manifold, and let $F : P \times U \rightarrow Y$ be a smooth map of the form*

$$F(p, x) = L(p, x) + Q(p, x) + e(p)$$

such that:

1. L is smooth, for each $p \in P$, $L_p := L(p, \cdot) : X \rightarrow Y$ is a Fredholm operator, and we have $\sup_{p \in P} \|L_p\|_{\mathcal{L}(X, Y)} < \infty$,

2. Q is smooth and there exists a $c_Q > 0$ such that, for all $x_1, x_2 \in X$ and all $p \in P$, we have

$$(3.12) \quad \|Q(x_1, p) - Q(x_2, p)\|_Y \leq c_Q (\|x_1\|_X + \|x_2\|_X) \|x_1 - x_2\|_X,$$

and

3. $e : P \rightarrow Y$ is smooth and there is a constant c_e such that $\|e\|_Y \leq c_e$.

Let $I \subset X$ be a finite dimensional subspace and let $\pi : X \rightarrow I$ be a projection onto I . Let $O \subset Y$ be a finite dimensional subspace, let $\Pi : Y \rightarrow O$ be a projection onto O , and denote by $\iota : O \rightarrow Y$ the inclusion. Suppose that, for all $p \in P$, the operator $\bar{L}_p : O \oplus X \rightarrow I \oplus Y$ defined by

$$\bar{L}_p := \begin{pmatrix} 0 & \pi \\ \iota & L_p \end{pmatrix}$$

is invertible, and suppose that $c_R := \sup_{p \in P} \|\bar{L}_p^{-1}\|_{\mathcal{L}(Y, X)} < \infty$.

If $c_e \ll_{c_R, c_Q} 1$, then there is an open neighborhood \mathcal{F} of $0 \in I$, an open subset $V \subset P \times U$ containing $P \times \{0\}$, and a smooth map

$$x : P \times \mathcal{F} \rightarrow X$$

such that, for each $(p, x_0) \in \mathcal{F} \times P$, $(p, x(p, x_0))$ is the unique solution $(p, x) \in V$ of

$$(3.13) \quad (\text{id}_Y - \Pi)F(p, x) = 0 \quad \text{and} \quad \pi x = x_0.$$

In particular, if we define $\text{ob}: P \times \mathcal{F} \rightarrow O$ by

$$\text{ob}(p, x_0) := \Pi F(p, x(p, x_0)),$$

then the map $\text{ob}^{-1}(0) \rightarrow F^{-1}(0) \cap V$ defined by

$$(p, x_0) \mapsto (p, x(p, x_0))$$

is a homeomorphism. Moreover, for every $(p, x_0) \in P \times \mathcal{F}$ and $x = x(p, x_0)$, we have an exact sequence

$$0 \rightarrow \ker \partial_x F(p, x) \rightarrow I \xrightarrow{\partial_{x_0} \text{ob}(p, x_0)} O \rightarrow \text{coker } \partial_x F(p, x) \rightarrow 0;$$

which induces an isomorphism $\det \partial_x F \cong \det I \otimes (\det O)^*$.

Proof sketch. This result is essentially a summary of the discussion in Guo and Wu [GW13, Section 5]; see also [DK90, Proposition 4.2.4]. The crucial point is that \bar{L}_p induces an inverse to $(\text{id}_Y - \Pi)L_p: \ker \pi \rightarrow \ker \Pi$; thus by the Inverse Function Theorem there are $\sigma, \tau > 0$ such that $U' := B_\sigma(0) \times B_\tau(0) \subset I \times \ker \pi$, and there is a smooth map $\Xi: P \times U' \rightarrow \ker \pi$ such that, for each $p \in P$ and $x \in B_\sigma(0)$:

1. $\Xi(p, x_0, 0) = 0$,
2. $\Xi(p, x_0, \cdot)$ is a diffeomorphism onto its image, and
3. for all $p \in P$ and $(x_0, x_1) \in U'$, we have

$$\tilde{F}(p, x_0, x_1) = F(p, x_0, \Xi(p, x_0, x_1)) = \begin{pmatrix} f(p, x_0, x_1) \\ G_p(x_0) \end{pmatrix} + \epsilon(p)$$

where $G_p: \ker \pi \rightarrow \ker \Pi$ is the linear isomorphism induced by \bar{L}_p and $f(p, 0, 0) = 0$.

If $c_\epsilon \ll 1$, then $G_p^{-1}(\text{id}_Y - \Pi)\epsilon(p) \in B_\tau(0)$ and we can take

$$\mathcal{F} = B_\sigma(0) \quad \text{and} \quad x(p, x_0) := (x_0, G_p^{-1}(\text{id}_Y - \Pi)\epsilon(p)).$$

We have

$$\ker \partial_x F \cong \ker \partial_x \tilde{F} \quad \text{and} \quad \text{coker } \partial_x F \cong \text{coker } \partial_x \tilde{F}.$$

However, $\partial_x \tilde{F}$ induces $G_p(x_0)$ from $\ker \pi$ to $\ker \Pi$. Therefore,

$$\ker \partial_x \tilde{F} \cong \ker \partial_{x_0} f \quad \text{and} \quad \text{coker } \partial_x \tilde{F} \cong \text{coker } \partial_{x_0} f.$$

Since

$$\text{ob}(p, x_0) = f(p, x_0, G_p^{-1}(\text{id}_Y - \Pi)\epsilon(p)),$$

it follows that

$$\ker \partial_x F \cong \ker \partial_{x_0} \text{ob} \quad \text{and} \quad \text{coker } \partial_x F \cong \text{coker } \partial_{x_0} \text{ob}. \quad \square$$

3.4 Orientations

For the purpose of counting solutions to (2.14) orientations play an important role. Suppose a trivialization $\tau: \det L \cong \mathbf{R}$ has been chosen. If $\mathbf{p} \in \mathcal{P}$ and $[c] \in \mathfrak{M}_{\text{SW}}(\mathbf{p})$ is irreducible and unobstructed, then $\det L_c = \det(0) \otimes \det(0)^* = \mathbf{R} \otimes \mathbf{R}^*$ is canonically trivial, and we define $\tau([c]) = +1$ if the isomorphism $\tau_{[c]}: \mathbf{R} \cong \mathbf{R}$ is orientation preserving and $\tau(c) = -1$ if it is orientation reversing. If $\mathbf{p}_0 \in \mathcal{P}$ is such that all $[c] \in \mathfrak{M}_{\text{SW}}(\mathbf{p}_0)$ are irreducible and unobstructed, and $\mathfrak{M}_{\text{SW}}(\mathbf{p}_0)$ is finite, then we can define

$$n_{\text{SW}}(\mathbf{p}_0) := \sum_{[c] \in \mathfrak{M}_{\text{SW}}(\mathbf{p}_0)} \tau([c]).$$

The following is a useful criterion to check whether $\det L$ can be trivialized.

Proposition 3.14. *Suppose that algebraic data as in Definition 2.6 and compatible geometric data as in Definition 2.8 have been fixed. Let $\rho_G: G \rightarrow \text{Sp}(S)$ be the restriction of the quaternionic representation $\rho: H \rightarrow \text{Sp}(S)$ to $G \triangleleft H$. Denote by $c_2 \in B\text{Sp}(S)$ the universal second Chern class. If $(B\rho_G)^* c_2 \in H^4(BG, \mathbf{Z})$ can be written as*

$$(3.15) \quad (B\rho_G)^* c_2 = 2x + \alpha_1 y_1^2 + \cdots + \alpha_k y_k^2$$

with $x \in H^4(BG, \mathbf{Z})$, $y_1, \dots, y_k \in H^2(BG, \mathbf{Z})$, and $\alpha_1, \dots, \alpha_k \in \mathbf{Z}$, then

$$\det L \rightarrow \mathcal{P} \times \frac{\Gamma(\mathfrak{S}) \times \mathcal{A}(Q)}{\mathcal{G}(P)}$$

is trivial.

Proof. The parameter space \mathcal{P} is contractible; hence, it is enough to fix an element $\mathbf{p} \in \mathcal{P}$ and prove that $\det L$ is trivial over the second factor. We need to show that if $(c_t)_{t \in [0,1]}$ is a path in $\Gamma(\mathfrak{S}) \times \mathcal{A}_B(Q)$ and $u \in \mathcal{G}(P)$ is such that $u c_1 = c_0$, then the spectral flow of $(L_{c_t})_{t \in [0,1]}$ is even. The mapping torus of $u: Q \rightarrow Q$ is a principal H -bundle \mathbf{Q} over $S^1 \times M$, and the path $(c_t)_{t \in [0,1]}$ induces a connection \mathbf{A} on \mathbf{Q} . Over $S^1 \times M$ we also have an adjoint bundle $\mathfrak{g}_{\mathbf{P}}$ and the spinor bundles \mathfrak{S}^+ and \mathfrak{S}^- associated with \mathbf{Q} via the quaternionic representation $\rho: H \rightarrow \text{Sp}(S)$. According to Atiyah–Singer–Patodi, the spectral flow of $(L_{c_t})_{t \in [0,1]}$ is the index of the operator $\mathbf{L} = \partial_t - L_{c_t}$ which can be identified with an operator

$$\mathbf{L}: \Gamma(\mathfrak{S}^+) \oplus \Omega^1(S^1 \times M, \mathfrak{g}_{\mathbf{P}}) \rightarrow \Gamma(\mathfrak{S}^-) \oplus \Omega^+(S^1 \times M, \mathfrak{g}_{\mathbf{P}}) \oplus \Omega^0(S^1 \times M, \mathfrak{g}_{\mathbf{P}}).$$

In our case, \mathbf{L} is homotopic through Fredholm operators to the sum of the Dirac operator $\mathcal{D}_{\mathbf{A}}^+: \Gamma(\mathfrak{S}^+) \rightarrow \Gamma(\mathfrak{S}^-)$ and the Atiyah–Hitchin–Singer operator $d_{\mathbf{A}}^+ \oplus d_{\mathbf{A}}^*$ for $\mathfrak{g}_{\mathbf{P}}$. The index of the Atiyah–Hitchin–Singer operator is $-2p_1(\mathfrak{g}_{\mathbf{P}})$ and thus even. To compute the index of the Dirac operator, observe that the vector bundle $\mathbf{V} := \mathbf{Q} \times_{\rho} S$ inherits from S the structure of a left-module over \mathbf{H} and that

$$\mathfrak{S}^{\pm} = \mathcal{S}^{\pm} \otimes_{\mathbf{H}} \mathbf{V},$$

where \mathcal{S}^\pm are the usual spinor bundles of $S^1 \times M$ with the spin structure induced from that on M and we use the structure of \mathcal{S}^\pm as a right-modules over \mathbf{H} . \mathfrak{S}^\pm is a real vector bundle: it is a real form of $\mathcal{S}^\pm \otimes_{\mathbf{C}} \mathbf{V}$. Therefore, the complexification of \mathcal{D}_A^\pm is the standard complex Dirac operator on \mathcal{S}^\pm twisted by \mathbf{V} . By the Atiyah–Singer Index Theorem,

$$\begin{aligned} \text{index } \mathcal{D}_A^+ &= \int_{S^1 \times M} \hat{A}(S^1 \times M) \text{ch}(\mathbf{V}) \\ &= \int_{S^1 \times M} \text{ch}_2(\mathbf{V}) = - \int_{S^1 \times M} c_2(\mathbf{V}). \end{aligned}$$

The classifying map $f_V: S^1 \times M \rightarrow B\text{Sp}(S)$ of \mathbf{V} is related to the classifying map $f_Q: S^1 \times M \rightarrow BG$ of \mathbf{Q} through

$$f_V = B\rho_G \circ f_Q,$$

and

$$c_2(\mathbf{V}) = f_V^* c_2 = f_Q^* (B\rho_G)^* c_2.$$

Since the intersection form of $S^1 \times M$ is even, the hypothesis implies that the right-hand side of the above index formula is even. \square

Remark 3.16. If G is simply-connected, then the condition (3.15) is satisfied if and only if the image of

$$(\rho_G)_*: \pi_3(G) \rightarrow \pi_3(\text{Sp}(S)) = \mathbf{Z}$$

is generated by an even integer. To see this, observe that BG is 3-connected; hence, by the Hurewicz theorem, $H_4(BG, \mathbf{Z}) = \pi_4(BG) \cong \pi_3(G)$ and $H_i(BG, \mathbf{Z}) = 0$ for $i = 1, 2, 3$. The same is true for $\text{Sp}(S)$, and we have a commutative diagram

$$\begin{array}{ccc} H_4(BG, \mathbf{Z}) & \xrightarrow{(B\rho_G)^*} & H_4(B\text{Sp}(S), \mathbf{Z}) \\ \downarrow \cong & & \downarrow \cong \\ \pi_3(G) & \xrightarrow{(\rho_G)_*} & \pi_3(\text{Sp}(S)). \end{array}$$

The group $H_4(BG, \mathbf{Z})$ is freely generated by some elements x_1, \dots, x_k . Let x^1, \dots, x^k be the dual basis of $H^4(BG, \mathbf{Z}) = \text{Hom}(H_4(BG, \mathbf{Z}), \mathbf{Z})$. Likewise, $H_4(B\text{Sp}(S), \mathbf{Z})$ is freely generated by the unique element z satisfying $\langle c_2, z \rangle = 1$. We have

$$(3.17) \quad (B\rho_G)^* c_2 = \sum_{i=1}^k \langle (B\rho_G)^* c_2, x_i \rangle x^i$$

and

$$\langle (B\rho_G)^* c_2, x_i \rangle = \langle c_2, (B\rho_G)_* x_i \rangle.$$

Therefore, the coefficients in the sum (3.17) are all even if and only if the image of $(B\rho_G)_*$ is generated by $2mz$ for some $m \in \mathbf{Z}$.

Example 3.18. The hypothesis of Proposition 3.14 holds when $S = \mathbf{H} \otimes_{\mathbf{C}} W$ for some complex Hermitian vector space W of dimension n and ρ_G is induced from a unitary representation $G \rightarrow \mathrm{U}(W)$; as is the case for the representations leading to the classical Seiberg–Witten and $\mathrm{U}(n)$ –monopole equations, see Example 2.15 and Example A.1. To see that $(B\rho_G)^*c_2$ is of the desired form, note that if E is a rank n Hermitian vector bundle, then the corresponding quaternionic Hermitian bundle obtained via the inclusion $\mathrm{U}(n) \rightarrow \mathrm{Sp}(n)$ is $\mathbf{H} \otimes_{\mathbf{C}} E = E \oplus \bar{E}$ and

$$c_2(\mathbf{H} \otimes_{\mathbf{C}} E) = c_2(E \oplus \bar{E}) = 2c_2(E) - c_1(E)^2.$$

Example 3.19. The hypothesis of Proposition 3.14 is also satisfied when $S = \mathbf{H} \otimes_{\mathbf{R}} W$ for a real Euclidean vector space W , and ρ_G is induced from an orthogonal representation $G \rightarrow \mathrm{SO}(W)$; as is the case for the equation for flat $G^{\mathbf{C}}$ –connections, see Example A.2. To see that $(B\rho_G)^*c_2$ is of the desired form, note that if E is a Euclidean vector bundle of rank n , then the associated quaternionic Hermitian vector bundle is $\mathbf{H} \otimes_{\mathbf{R}} E$ and

$$c_2(\mathbf{H} \otimes_{\mathbf{R}} E) = -2p_1(E).$$

If two quaternionic representations satisfy the hypothesis of Proposition 3.14, then so does their direct sum. Therefore, the previous two examples together show that $\det L$ is trivial for the ADHM Seiberg–Witten equation described in Example A.3.

Example 3.20. In general, $\det L$ need not be orientable. If $S = \mathbf{H}$ and $G = H = \mathrm{Sp}(1)$ acts on S by right multiplication, then it is easy to see that the gauge transformation of the trivial bundle $Q = S^3 \times \mathrm{SU}(2)$ induced by $S^3 \cong \mathrm{SU}(2)$ gives rise to an odd spectral flow.

4 The Haydys correspondence

In order to discuss the deformation theory *on* the boundary of $\overline{\mathfrak{M}}_{\mathrm{SW}}$, it will be helpful to review the correspondence, discovered by Haydys [Hay12, Section 4.1], between Fueter sections of \mathfrak{X} and solutions $(\Phi, A) \in \Gamma(\mathfrak{S}^{\mathrm{reg}} \times \mathcal{A}_B(Q))$ of

$$(4.1) \quad \not{D}_A \Phi = 0 \quad \text{and} \quad \mu(\Phi) = 0.$$

For what follows it will be important to recall some details of hyperkähler reduction construction.

Proposition 4.2 (Hitchin, Karlhede, Lindström, and Roček [HKLR87, Section 3(D)]). *If $\rho: G \rightarrow \mathrm{Sp}(S)$ is a quaternionic representation, then the following hold:*

1. *The space*

$$X := S^{\mathrm{reg}} // G := (\mu^{-1}(0) \cap S^{\mathrm{reg}}) / G$$

is an orbifold (with discrete isotropy groups).

2. Denote by $p: \mu^{-1}(0) \cap S^{\text{reg}} \rightarrow X$ the canonical projection. Set

$$\begin{aligned} \mathfrak{S} &:= (\ker dp)^\perp \cap T(\mu^{-1}(0) \cap S^{\text{reg}}) \quad \text{and} \\ \mathfrak{R} &:= \mathfrak{S}^\perp \subset TS|_{\mu^{-1}(0) \cap S^{\text{reg}}}. \end{aligned}$$

For each $\Phi \in \mu^{-1}(0) \cap S^{\text{reg}}$, $(dp)_\Phi: \mathfrak{S}_\Phi \rightarrow T_{[\Phi]}X$ is an isomorphism, and

$$(4.3) \quad \mathfrak{R}_\Phi = \text{im}(\rho(\cdot)\Phi \oplus \bar{\gamma}(\cdot)\Phi: \mathfrak{g} \otimes \mathbf{H} \rightarrow S).$$

3. For each $\Phi \in \mu^{-1}(0) \cap S^{\text{reg}}$, γ preserves the splitting $S = \mathfrak{S}_\Phi \oplus \mathfrak{R}_\Phi$.

4. There exist a Riemannian metric g_X on X and a Clifford multiplication

$$\gamma_X: \text{Im } \mathbf{H} \rightarrow \text{End}(TX)$$

such that

$$p^*g_X = \langle \cdot, \cdot \rangle \quad \text{and} \quad p^*\gamma_X = \gamma.$$

5. γ_X is parallel with respect to g_X ; hence, X is a hyperkähler orbifold—which is called the **hyperkähler quotient** of S by G .

Remark 4.4. More generally, $\mu^{-1}(0)/G$ can be decomposed into a union of hyperkähler manifolds according to the conjugacy class of the stabilizers in G ; see Dancer and Swann [DS97, Theorem 2.1].

4.1 Lifting sections of \mathfrak{X}

Proposition 4.5. *Given a set of geometric data as in Definition 2.8, set*

$$X := S^{\text{reg}} // G \quad \text{and} \quad \mathfrak{X} := (\mathfrak{s} \times R) \times_{\text{Sp}(1) \times K} X.$$

Denote by $p: S^{\text{reg}} \cap \mu^{-1}(0) \rightarrow X$ the canonical projection.

1. If $s \in \Gamma(\mathfrak{X})$, then there exist a principal H -bundle Q together with an isomorphism $Q \times_H K \cong R$ and a section $\Phi \in \Gamma(\mathfrak{S}^{\text{reg}})$ of

$$\mathfrak{S}^{\text{reg}} := (\mathfrak{s} \times Q) \times_{\text{Sp}(1) \times H} S^{\text{reg}}$$

satisfying

$$\mu(\Phi) = 0 \quad \text{and} \quad s = p \circ \Phi.$$

Q and $Q \times_H K \cong R$ are unique up to isomorphism, and every two lifts Φ are related by a unique gauge transformation in $\mathcal{G}(P)$.

2. Suppose $B \in \mathcal{A}(R)$. If $\Phi \in \Gamma(\mathfrak{S}^{\text{reg}})$ satisfies $\mu(\Phi) = 0$, then there is a unique $A \in \mathcal{A}_B(Q)$ such that $\nabla_A \Phi \in \Omega^1(M, \mathfrak{S}_\Phi)$. In particular, for this connection

$$p_*(\mathbb{D}_A \Phi) = \mathfrak{F}(s).$$

3. The condition $p_*(\mathbb{D}_A \Phi) = \mathfrak{F}(s)$ characterizes $A \in \mathcal{A}_B(Q)$ uniquely.

Proof. Part (1) is proved by observing that the lift exists locally and that the obstruction to the local lifts patching defines a cocycle which determines Q ; see [Hay12] for details.

We prove (2). For an arbitrary connection $A_0 \in \mathcal{A}_B(Q)$ and for all $x \in M$, we have

$$(\nabla_{A_0} \Phi)(x) \in T_x^* M \otimes T_{\Phi(x)}(S^{\text{reg}} \cap \mu^{-1}(0)).$$

By Proposition 4.2(2) there exists a unique $a \in \Omega^1(M, \mathfrak{g}_P)$ such that

$$\nabla_{A_0+a} \Phi \in \Omega^1(M, \mathfrak{S}_\Phi).$$

The assertion in (2) now follows from the fact that for $s = p \circ \Phi$ we have $p_*(\nabla_{A_0} \Phi) = \nabla_B s$ and the definitions of \mathbb{D}_A and \mathfrak{F} .

We prove (3). If $a \in \Omega^1(M, \mathfrak{g}_P)$ and $A + a$ also satisfies this condition, then we must have

$$\bar{\gamma}(a)\Phi = 0.$$

This is impossible because $\Phi \in \Gamma(\mathfrak{S}^{\text{reg}})$, that is, $(d\mu)_\Phi$ is surjective; hence, its adjoint $\bar{\gamma}(\cdot)\Phi$ is injective. \square

Proposition 4.6. *Given a set of geometric data as in Definition 2.8, set*

$$R := Q \times_H K, \quad \mathfrak{X} := (\mathfrak{s} \times R) \times_{\text{Sp}(1) \times K} X, \quad \text{and} \quad \mathfrak{S}^{\text{reg}} := (\mathfrak{s} \times Q) \times_{\text{Sp}(1) \times H} S^{\text{reg}}.$$

The map

$$\begin{aligned} \Gamma(\mu^{-1}(0) \cap \mathfrak{S}^{\text{reg}}) / \mathcal{G}(P) &\rightarrow \Gamma(\mathfrak{X}) \\ [\Phi] &\mapsto p \circ \Phi \end{aligned}$$

is a homeomorphism onto its image.

Proof. Fix $\Phi_0 \in \Gamma(\mu^{-1}(0) \cap \mathfrak{S}^{\text{reg}})$ and set $s_0 := p \circ \Phi_0 \in \Gamma(\mathfrak{X})$. Given $0 < \sigma \ll 1$, for every $\Phi \in \Gamma(\mu^{-1}(0) \cap \mathfrak{S}^{\text{reg}})$ with $\|\Phi - \Phi_0\|_{L^\infty} < \sigma$, there is a unique $u \in \mathcal{G}(P)$ such that

$$u\Phi \perp \text{im}(\rho(\cdot)\Phi_0 : \Gamma(\mathfrak{g}_P) \rightarrow \Gamma(\mathfrak{S}));$$

moreover, for every $k \in \mathbb{N}$,

$$\|u\Phi - \Phi_0\|_{C^k} \lesssim_k \|\Phi - \Phi_0\|_{C^k}.$$

Thus, it suffices to show that the map

$$\left\{ \Phi \in \Gamma(\mu^{-1}(0) \cap \mathfrak{S}^{\text{reg}}) : \begin{array}{l} \|\Phi - \Phi_0\|_{L^\infty} < \sigma \text{ and} \\ \Phi \perp \text{im}(\rho(\cdot)\Phi_0 : \Gamma(\mathfrak{g}_P) \rightarrow \Gamma(\mathfrak{S})) \end{array} \right\} \rightarrow \Gamma(\mathfrak{X})$$

is a homeomorphism onto its image. This, however, is immediate from the Implicit Function Theorem and the fact that the tangent space at Φ_0 to the former space is $\Gamma(\mathfrak{H}_{\Phi_0})$ and the derivative of this map is the canonical isomorphism $\Gamma(\mathfrak{H}_{\Phi_0}) \cong \Gamma(s_0^*V\mathfrak{X})$ from Proposition 4.2(2). \square

In the situation of Proposition 4.5, we have $|\Phi| = |\hat{v} \circ s|$. The preceding results thus imply the following.

Corollary 4.7. *Let R be a principal K -bundle. Set $\mathfrak{X} := R \times_K X$ and*

$$\mathfrak{M}_F := \{(\mathbf{p}, s) \in \mathcal{P} \times \Gamma(\mathfrak{X}) : \mathfrak{F}(s) = 0 \text{ and } \|\hat{v} \circ s\|_{L^2} = 1\}.$$

The map

$$\bigsqcup_Q \partial \mathfrak{M}_{\text{SW}, Q} \rightarrow \mathfrak{M}_F$$

defined by

$$(\mathbf{p}, [(\Phi, A)]) \mapsto (\mathbf{p}, p \circ \Phi)$$

is a homeomorphism. Here, the disjoint union is taken over all isomorphism classes of principal H -bundles Q with isomorphisms $Q \times_H K \cong R$.

4.2 Lifting infinitesimal deformations

Proposition 4.8. *For $\Phi \in \Gamma(\mu^{-1}(0) \cap \mathfrak{S}^{\text{reg}})$, set $s := p \circ \Phi$ and let $A \in \mathcal{A}_B(Q)$ be as in Proposition 4.5. The isomorphism $p_* : \Gamma(\mathfrak{H}_\Phi) \rightarrow \Gamma(s^*V\mathfrak{X})$ identifies $\pi_{\mathfrak{S}} \nabla_A : \Omega^0(M, \mathfrak{H}_\Phi) \rightarrow \Omega^1(M, \mathfrak{H}_\Phi)$ with $\nabla_B : \Omega^0(M, s^*V\mathfrak{X}) \rightarrow \Omega^1(M, s^*V\mathfrak{X})$.*

Proof. If (Φ_t) is a one-parameter family of local sections of $\mu^{-1}(0) \cap \mathfrak{S}^{\text{reg}}$ with

$$(\partial_t \Phi_t)|_{t=0} = \phi,$$

A_t are as in Proposition 4.5, and $a = (\partial_t A_t)|_{t=0}$, then we have

$$\partial_t \left(\pi_{\mathfrak{S}_{\Phi_t}} \nabla_{A_t} \Phi_t \right) \Big|_{t=0} = (\partial_t \pi_{\mathfrak{S}_{\Phi_t}}) \Big|_{t=0} \nabla_{A_0} \Phi_0 + \pi_{\mathfrak{S}_{\Phi_0}}(\rho(a)\Phi_0) + \pi_{\mathfrak{S}_{\Phi_0}}(\nabla_{A_0} \phi).$$

The first term vanishes because $\nabla_{A_0} \Phi_0 \in \Gamma(\mathfrak{H}_{\Phi_0})$, and the second term vanishes because of Proposition 4.2(2). \square

If $\Phi \in \Gamma(\mu^{-1}(0) \cap \mathfrak{S}^{\text{reg}})$, then the induced splitting $\mathfrak{S} = \mathfrak{H}_\Phi \oplus \mathfrak{N}_\Phi$ given by Proposition 4.2(2) need not be parallel for A as in Proposition 4.5.

Definition 4.9. The second fundamental forms of the splitting $\mathfrak{H}_\Phi \oplus \mathfrak{R}_\Phi$ are defined by

$$\begin{aligned}\Pi &:= \pi_{\mathfrak{R}} \nabla_A \in \Omega^1(M, \text{Hom}(\mathfrak{H}_\Phi, \mathfrak{R}_\Phi)) \quad \text{and} \\ \Pi^* &:= -\pi_{\mathfrak{H}} \nabla_A \in \Omega^1(M, \text{Hom}(\mathfrak{R}_\Phi, \mathfrak{H}_\Phi)).\end{aligned}$$

We decompose the Dirac operator \mathcal{D}_A according to $\mathfrak{S} = \mathfrak{H}_\Phi \oplus \mathfrak{R}_\Phi$ as

$$(4.10) \quad \mathcal{D}_A = \begin{pmatrix} \mathcal{D}_{\mathfrak{H}} & -\gamma \Pi^* \\ \gamma \Pi & \mathcal{D}_{\mathfrak{R}} \end{pmatrix}$$

with

$$\begin{aligned}\mathcal{D}_{\mathfrak{H}} &:= \gamma(\pi_{\mathfrak{H}} \nabla_A): \Gamma(\mathfrak{H}_\Phi) \rightarrow \Gamma(\mathfrak{H}_\Phi) \quad \text{and} \\ \mathcal{D}_{\mathfrak{R}} &:= \gamma(\pi_{\mathfrak{R}} \nabla_A): \Gamma(\mathfrak{R}_\Phi) \rightarrow \Gamma(\mathfrak{R}_\Phi).\end{aligned}$$

The following result helps to better understand the off-diagonal terms in (4.10).

Proposition 4.11. Suppose $\Phi \in \Gamma(\mu^{-1}(0) \cap \mathfrak{S}^{\text{reg}})$ and $\mathcal{D}_A \Phi = 0$. Writing $\phi \in \Gamma(\mathfrak{R}_\Phi)$ as

$$\phi = \rho(\xi)\Phi + \bar{\gamma}(a)\Phi$$

for $\xi \in \Gamma(\mathfrak{g}_P)$ and $a \in \Omega^1(M, \mathfrak{g}_P)$, we have

$$-\gamma \Pi^* \phi = 2 \sum_{i=1}^3 \pi_{\mathfrak{H}} \left(\rho(a(e_i)) \nabla_{e_i}^A \Phi \right).$$

Here (e_1, e_2, e_3) is a local orthonormal frame.

Proof. Since $\nabla \Phi \in \Omega^1(M, \mathfrak{H}_\Phi)$ and $\mathcal{D}_A \Phi = 0$, we have

$$\begin{aligned}-\gamma \Pi^* (\rho(\xi)\Phi + \bar{\gamma}(a)\Phi) &= \sum_{i=1}^3 \gamma(e^i) \pi_{\mathfrak{H}} \left(\rho(\xi) \nabla_{e_i}^A \Phi + \bar{\gamma}(a) \nabla_{e_i}^A \Phi \right) \\ &= \sum_{i=1}^3 \pi_{\mathfrak{H}} \left((\gamma(e^i) \bar{\gamma}(a) + \bar{\gamma}(a) \gamma(e^i)) \nabla_{e_i}^A \Phi \right) \\ &= 2 \sum_{i=1}^3 \pi_{\mathfrak{H}} (\rho(a(e_i)) \nabla_{e_i}^A \Phi). \quad \square\end{aligned}$$

Proposition 4.12. The isomorphism $p_*: \Gamma(\mathfrak{H}_\Phi) \rightarrow \Gamma(s^*V\mathfrak{X})$ identifies the linearized Fueter operator $(d\mathfrak{F})_s: \Gamma(s^*V\mathfrak{X}) \rightarrow \Gamma(s^*V\mathfrak{X})$ with $\mathcal{D}_{\mathfrak{H}}: \Gamma(\mathfrak{H}_\Phi) \rightarrow \Gamma(\mathfrak{H}_\Phi)$.

Proof. The linearized Fueter operator is given by

$$(d\mathfrak{F})_s \hat{s} = \gamma(\nabla_B \hat{s})$$

The assertion thus follows from Proposition 4.2(4) and Proposition 4.8. □

5 Deformation theory of Fueter sections

Proposition 5.1. *Let $s_0 \in \Gamma(\mathfrak{X})$ be a Fueter section with respect to $\mathfrak{p}_0 = (g_0, B_0) \in \mathcal{P}$. Denote by $c_0 \in \Gamma(\mathfrak{S}^{\text{reg}}) \times \mathcal{A}(P)$ a lift of s_0 . There exist an open neighbourhood U of $\mathfrak{p}_0 \in \mathcal{P}$, an open neighborhood*

$$\mathcal{F}_\partial \subset I_\partial := \ker(d\mathfrak{F})_{s_0} \cap (\hat{v} \circ s)^{\perp}$$

of 0, a smooth map

$$\text{ob}_\partial: U \times \mathcal{F}_\partial \rightarrow \text{coker}(d\mathfrak{F})_{s_0},$$

an open neighborhood V of $([\mathfrak{p}_0, c_0]) \in \partial\mathfrak{M}_{\text{SW}}$, and a homeomorphism

$$\mathfrak{x}_\partial: \text{ob}_\partial^{-1}(0) \rightarrow V \subset \partial\mathfrak{M}_{\text{SW}}$$

which maps $(\mathfrak{p}_0, 0)$ to $(\mathfrak{p}_0, 0, [c_0])$ and commutes with the projections to \mathcal{P} .

Since $\partial\mathfrak{M}_{\text{SW}} \cong \mathfrak{M}_F$ through the Haydys correspondence, this has a straightforward proof using Lemma 3.11, which makes no reference to the Seiberg–Witten equation. However, this is not the approach we take because our principal goal is to compare the deformation theory of Fueter sections with that of solutions of the Seiberg–Witten equation.

Fix $k \in \mathbf{N}$ and $p \in (1, \infty)$ with $(k+1)p > 3$. Let

$$\partial\mathfrak{M}_{\text{SW}}^{k,p} = \left\{ (\mathfrak{p}, [(\Phi, A)]) \in \mathcal{P} \times \frac{W^{k+1,p}\Gamma(\mathfrak{S}) \times W^{k,p}\mathcal{A}(Q)}{W^{k+1,p}\mathcal{G}(P)} : \begin{array}{l} A \text{ induces } B, \\ (\Phi, A) \text{ satisfies (4.1),} \\ \text{and } \|\Phi\|_{L^2} = 1 \end{array} \right\}.$$

By the Haydys correspondence $\partial\mathfrak{M}_{\text{SW}}^{k,p}$ is homeomorphic to $\mathfrak{M}_F^{k,p}$, the universal moduli space of normalized $W^{k+1,p}$ Fueter sections of \mathfrak{X} . Consequently, for $\ell \in \mathbf{N}$ and $q \in (1, \infty)$ with $\ell \geq k$ and $q \geq p$, the inclusions $\partial\mathfrak{M}_{\text{SW}}^{\ell,q} \subset \partial\mathfrak{M}_{\text{SW}}^{k,p} \subset \partial\mathfrak{M}_{\text{SW}}$ are homeomorphisms; see also Proposition 5.11.

Proposition 5.2. *Assume the situation of Proposition 5.1. For $\mathfrak{p} \in \mathcal{P}$, set*

$$X_0 := W^{k+1,p}\Gamma(\mathfrak{S}) \oplus W^{k,p}\Omega^1(M, \mathfrak{g}_P) \oplus W^{k,p}\Omega^0(M, \mathfrak{g}_P)$$

$$\text{and } Y := W^{k,p}\Gamma(\mathfrak{S}) \oplus W^{k+1,p}\Omega^1(M, \mathfrak{g}_P) \oplus W^{k+1,p}\Omega^0(M, \mathfrak{g}_P) \oplus \mathbf{R},$$

and define a linear map $L_{\mathfrak{p},0}: X_0 \rightarrow Y$, a quadratic map $Q_{\mathfrak{p},0}: X_0 \rightarrow Y$, and $e_{\mathfrak{p},0} \in Y$ by

$$L_{\mathfrak{p},0} := \begin{pmatrix} -\mathcal{D}_{A_0} & -\bar{\gamma}(\cdot)\Phi_0 & -\rho(\cdot)\Phi_0 \\ -2 * \mu(\Phi_0, \cdot) & 0 & 0 \\ -\rho^*(\cdot)\Phi_0^* & 0 & 0 \\ 2\langle \cdot, \Phi_0 \rangle_{L^2} & 0 & 0 \end{pmatrix},$$

$$Q_{\mathfrak{p},0}(\phi, a, \xi) := \begin{pmatrix} -\bar{\gamma}(a)\phi \\ - * \mu(\phi) \\ 0 \\ \|\phi\|_{L^2}^2 \end{pmatrix}, \quad \text{and } e_{\mathfrak{p},0} := \begin{pmatrix} -\mathcal{D}_{A_0}\Phi_0 \\ -\mu(\Phi_0) \\ 0 \\ \|\Phi_0\|_{L^2}^2 - 1 \end{pmatrix},$$

respectively.⁶

There exist a neighborhood U of $\mathbf{p}_0 \in \mathcal{P}$ and $\sigma > 0$, such that, for every $\mathbf{p} \in U$ and $\hat{c} = (\phi, a, \xi) \in B_\sigma(0) \subset X_0$, we have

$$(5.3) \quad L_{\mathbf{p},0}\hat{c} + Q_{\mathbf{p},0}(\hat{c}) + \epsilon_{\mathbf{p},0} = 0$$

if and only if $\xi = 0$ and $(\Phi, A) = (\Phi_0 + \phi, A_0 + a)$ satisfies

$$(5.4) \quad \mathcal{D}_A\Phi = 0 \quad \text{and} \quad \mu(\Phi) = 0$$

as well as

$$\|\Phi\|_{L^2} = 1 \quad \text{and} \quad \rho^*(\Phi\Phi_0^*) = 0.$$

Remark 5.5. The above proposition engages in the following abuse of notation. If $A_0 \in \mathcal{A}_B(Q)$ and $B' \in \mathcal{A}(R)$, then $b = B' - B \in \Omega^1(M, \mathfrak{g}_R)$. Since $\text{Lie}(K) = \text{Lie}(G)^\perp \subset \text{Lie}(H)$ we have a map $\Omega^1(M, \mathfrak{g}_R) \rightarrow \Omega^1(M, \mathfrak{g}_Q)$ and can identify $A_0 \in \mathcal{A}_B(Q)$ with “ A_0 ” = $A_0 + b \in \mathcal{A}_{B'}(Q)$.

Together with (the argument from the proof of) Proposition 4.6 we obtain the following.

Corollary 5.6. *Assume the situation of Proposition 5.1. With $U \subset \mathcal{P}$ and $\sigma > 0$ as in Proposition 5.2, the map*

$$\{(\mathbf{p}, \hat{c}) \in U \times B_\sigma(0) \text{ satisfying (5.3)}\} \rightarrow \partial\mathfrak{M}_{\text{SW}}$$

defined by

$$(\mathbf{p}, \phi, a, \xi) \mapsto (\mathbf{p}, [(\Phi_0 + \phi, A_0 + a)])$$

is a homeomorphism onto a neighborhood of $[c_0]$.

Proof of Proposition 5.2. If $\hat{c} = (\phi, a, \xi)$ satisfies (5.3), then $\Phi = \Phi_0 + \phi$ and $A = A_0 + a$ satisfy

$$\mathcal{D}_A\Phi + \rho(\xi)\Phi_0 = 0, \quad \mu(\Phi) = 0, \quad \text{and} \quad \rho^*(\phi\Phi_0^*) = 0.$$

Hence, by Proposition B.4,

$$0 = d_A\mu(\Phi) = -\rho(\mathcal{D}_A\Phi\Phi_0^*) = \rho^*(\rho(\xi)\Phi_0(\Phi_0 + \phi)) = R_{\Phi_0}^* R_{\Phi_0}\xi + O(|\xi||\phi|)$$

with

$$R_{\Phi_0} := \rho(\cdot)\Phi_0.$$

Since Φ_0 is regular, R_{Φ_0} is injective, and it follows that $\xi = 0$ if $|\phi| \lesssim \sigma \ll 1$ and \mathbf{p} is sufficiently close to \mathbf{p}_0 . □

⁶The term $\epsilon_{\mathbf{p},0}$ vanishes for $\mathbf{p} = \mathbf{p}_0$.

Proof of Proposition 5.1. Denote by $\iota: \operatorname{coker}(d\tilde{\mathcal{F}})_{s_0} \cong \operatorname{coker} \bar{\mathcal{D}}_{\mathfrak{S}} \rightarrow \Gamma(\mathfrak{S})$ the inclusion of the L^2 orthogonal complement of $\operatorname{im} \bar{\mathcal{D}}_{\mathfrak{S}}$. Denote by $\pi_0: \Gamma(\mathfrak{S}) \rightarrow I_{\partial}$ the L^2 orthogonal projection onto $I_{\partial} = \ker(d\tilde{\mathcal{F}})_{s_0} \cap (\hat{\nu} \circ s)^\perp \subset \ker \bar{\mathcal{D}}_{\mathfrak{S}} \cong \ker(d\tilde{\mathcal{F}})_{s_0}$. Define

$$\bar{\bar{\mathcal{D}}}_{\mathfrak{S}}: \operatorname{coker}(d\tilde{\mathcal{F}})_{s_0} \oplus \Gamma(\mathfrak{S}) \rightarrow I_{\partial} \oplus \mathbf{R} \oplus \Gamma(\mathfrak{S})$$

by

$$\bar{\bar{\mathcal{D}}}_{\mathfrak{S}} := \begin{pmatrix} 0 & \pi_0 \\ 0 & -2\langle \cdot, \Phi_0 \rangle_{L^2} \\ \iota & \bar{\mathcal{D}}_{\mathfrak{S}} \end{pmatrix}.$$

Set

$$(5.7) \quad \begin{aligned} \bar{X}_0 &:= \operatorname{coker}(d\tilde{\mathcal{F}})_{s_0} \oplus W^{k+1,p}\Gamma(\mathfrak{S}) \\ &\quad \oplus W^{k+1,p}\Gamma(\mathfrak{N}) \\ &\quad \oplus W^{k,p}\Omega^1(M, \mathfrak{g}_P) \oplus W^{k,p}\Omega^0(M, \mathfrak{g}_P) \quad \text{and} \\ \bar{Y} &:= I_{\partial} \oplus \mathbf{R} \oplus W^{k,p}\Gamma(\mathfrak{S}) \\ &\quad \oplus W^{k,p}\Gamma(\mathfrak{N}) \\ &\quad \oplus W^{k+1,p}\Omega^1(M, \mathfrak{g}_P) \oplus W^{k+1,p}\Omega^0(M, \mathfrak{g}_P). \end{aligned}$$

Define the operator $\bar{L}_{\mathbf{p},0}: \bar{X}_0 \rightarrow \bar{Y}$ by

$$(5.8) \quad \bar{L}_{\mathbf{p},0} := \begin{pmatrix} -\bar{\bar{\mathcal{D}}}_{\mathfrak{S}} & \gamma\Pi^* & 0 \\ -\gamma\Pi & -\bar{\mathcal{D}}_{\mathfrak{N}} & -\mathfrak{a} & 0 \\ 0 & -\mathfrak{a}^* & 0 & 0 \end{pmatrix}$$

with $\mathfrak{a}: \Omega^1(M, \mathfrak{g}_P) \oplus \Omega^0(M, \mathfrak{g}_P) \rightarrow \Gamma(\mathfrak{N})$ defined by

$$\mathfrak{a}(a, \xi) := \bar{\gamma}(a)\Phi_0 + \rho(\xi)\Phi.$$

The operator $\bar{\bar{\mathcal{D}}}_{\mathfrak{S}}$ is invertible because

$$\begin{pmatrix} \pi_0 \\ -2\langle \cdot, \Phi_0 \rangle_{L^2} \end{pmatrix}$$

is essentially the L^2 orthogonal projection onto $\ker \bar{\mathcal{D}}_{\mathfrak{S}}$. It can be verified by a direct computation that $\bar{L}_{\mathbf{p},0}$ is invertible and its inverse is given by

$$(5.9) \quad \begin{pmatrix} -\bar{\bar{\mathcal{D}}}_{\mathfrak{S}}^{-1} & 0 & -\bar{\bar{\mathcal{D}}}_{\mathfrak{S}}^{-1}\gamma\Pi^*(\mathfrak{a}^*)^{-1} \\ 0 & 0 & -(\mathfrak{a}^*)^{-1} \\ \mathfrak{a}^{-1}\gamma\Pi\bar{\bar{\mathcal{D}}}_{\mathfrak{S}}^{-1} & -\mathfrak{a}^{-1} & \mathfrak{a}^{-1}\bar{\mathcal{D}}_{\mathfrak{N}}(\mathfrak{a}^*)^{-1} + \mathfrak{a}^{-1}\gamma\Pi\bar{\bar{\mathcal{D}}}_{\mathfrak{S}}^{-1}\gamma\Pi^*(\mathfrak{a}^*)^{-1} \end{pmatrix}.$$

After possibly shrinking U , we can assume that $\bar{L}_{\mathbf{p},0}$ is invertible for every $\mathbf{p} \in U$.

Since $Q_{\mathbf{p},0}$ is a quadratic map and

$$(5.10) \quad \begin{aligned} \|Q_{\mathbf{p},0}(\phi, a, \xi)\|_Y &= \|\bar{\gamma}(a)\phi\|_{W^{k,p}} + \|\mu(\phi)\|_{W^{k+1,p}} + \|\phi\|_{L^2}^2 \\ &\lesssim \|a\|_{W^{k,p}} \|\phi\|_{W^{k+1,p}} + \|\phi\|_{W^{k+1,p}}^2, \end{aligned}$$

$Q_{\mathbf{p},0}$ satisfies (3.12); hence, we can apply Lemma 3.11 to complete the proof. \square

In the following regularity result, we decorate X_0 and Y with superscripts indicating the choice of the differentiability and integrability parameters k and p .

Proposition 5.11. *Assume the situation of Proposition 5.1. For each $k, \ell \in \mathbf{N}$ and $p, q \in (1, \infty)$ with $(k+1)p > 3$, $\ell \geq k$, and $q \geq p$, there are constants $c, \sigma > 0$ and an open neighborhood U of \mathbf{p}_0 in \mathcal{P} such that if $\mathbf{p} \in U$ and $\hat{c} \in B_\sigma(0) \subset X_0^{k,p}$ is solution of*

$$L_{\mathbf{p},0}\hat{c} + Q_{\mathbf{p},0}(\hat{c}) + \epsilon_{\mathbf{p},0} = 0,$$

then $\hat{c} \in X_0^{\ell,q}$ and $\|\hat{c}\|_{X_0^{\ell,q}} \leq c\|\hat{c}\|_{X_0^{k,p}}$.

Proof. Provided U is a sufficiently small neighborhood of \mathbf{p}_0 and $0 < \sigma \ll 1$, it follows from Banach's Fixed Point Theorem that $(0, \hat{c})$ is the unique solution in $B_\sigma(0) \subset \bar{X}^{k,p}$ of

$$\bar{L}_{\mathbf{p},0}(0, \hat{c}) + Q_{\mathbf{p},0}(\hat{c}) + \epsilon_{\mathbf{p},0} = \begin{pmatrix} \pi\hat{c} \\ 0 \end{pmatrix},$$

and that there exists a $(o, \hat{\mathfrak{d}}) \in B_\sigma(0) \subset \bar{X}^{\ell,q}$ such that

$$\bar{L}_{\mathbf{p},0}(o, \hat{\mathfrak{d}}) + Q_{\mathbf{p},0}(\hat{\mathfrak{d}}) + \epsilon_{\mathbf{p},0} = \begin{pmatrix} \pi\hat{\mathfrak{d}} \\ 0 \end{pmatrix}.$$

Since $\bar{X}^{\ell,q} \subset \bar{X}^{k,p}$ and $\|(o, \hat{\mathfrak{d}})\|_{\bar{X}^{k,p}} \leq \|(o, \hat{\mathfrak{d}})\|_{\bar{X}^{\ell,q}} \leq \sigma$, it follows that $(o, \hat{\mathfrak{d}}) = (0, \hat{c})$ and thus $\hat{c} \in \bar{X}^{\ell,q}$ and $\|\hat{c}\|_{X^{\ell,q}} \leq \sigma$. From this it follows easily that $\|\hat{c}\|_{X^{\ell,q}} \leq c\|\hat{c}\|_{X^{k,p}}$. \square

6 Deformation theory around $\varepsilon = 0$

In this section we will prove Theorem 2.29, whose hypotheses we will assume throughout.

Fix $k \in \mathbf{N}$ and $p \in (1, \infty)$ with $(k+1)p > 3$. Let

$$\mathfrak{M}_{\text{SW}}^{k,p} = \left\{ (\mathbf{p}, \varepsilon, [(\Phi, A)]) \in \mathcal{P} \times \mathbf{R}^+ \times \frac{W^{k+1,p}\Gamma(\mathfrak{S}) \times W^{k+2,p}\mathcal{A}(P)}{W^{k+3,p}\mathcal{G}(P)} : (\varepsilon, \Phi, A) \text{ satisfies (2.21)} \right\}.$$

For $\ell \in \mathbf{N}$ and $q \in (1, \infty)$ with $\ell \geq k$ and $q \geq p$, the inclusions $\mathfrak{M}_{\text{SW}}^{\ell,q} \subset \mathfrak{M}_{\text{SW}}^{k,p} \subset \mathfrak{M}_{\text{SW}}$ are homeomorphisms; see also Proposition 6.12.

6.1 Reduction to a slice

Proposition 6.1. *Let $c_0 = (\Phi_0, A_0) \in \Gamma(\mathfrak{S}^{\text{reg}}) \times \mathcal{A}(P)$ and $\mathfrak{p}_0 \in \mathcal{P}$. For $\mathfrak{p} \in \mathcal{P}$, set*

$$X_\varepsilon := W^{k+1,p}\Gamma(\mathfrak{S}) \oplus W^{k+2,p}\Omega^1(M, \mathfrak{g}_P) \oplus W^{k+2,p}\Omega^0(M, \mathfrak{g}_P)$$

and

$$\|(\phi, a, \xi)\|_{X_\varepsilon} := \|\phi\|_{W^{k+1,p}} + \|(a, \xi)\|_{W^{k,p}} + \varepsilon\|\nabla^{k+1}(a, \xi)\|_{L^p} + \varepsilon^2\|\nabla^{k+2}(a, \xi)\|_{L^p}.$$

There exist a neighborhood U of $\mathfrak{p}_0 \in \mathcal{P}$ and constants $\sigma, \varepsilon_0, c > 0$ such that the following holds. If $\mathfrak{p} \in U$, $\hat{c} = (\phi, a) \in X_\varepsilon$, and $\varepsilon \in (0, \varepsilon_0]$ are such that

$$\|\hat{c}\|_{X_\varepsilon} < \sigma,$$

then there exists a $W^{k+3,p}$ gauge transformation g such that $(\tilde{\phi}, \tilde{a}) = g(c_0 + \hat{c}) - c_0$ satisfies

$$\|(\tilde{\phi}, \tilde{a})\|_{X_\varepsilon} < c\sigma,$$

and

$$(6.2) \quad \varepsilon^2 d_{A_0 B}^* \tilde{a} - \rho^*(\tilde{\phi} \Phi_0^*) = 0.$$

Proof. To construct g , note that for $g = e^\xi$ with $\xi \in W^{k+3,p}\Omega^0(M, \mathfrak{g}_P)$ we have

$$\tilde{\phi} = \rho(\xi)\Phi_0 + \rho(\xi)\phi + \mathfrak{m}(\xi) \quad \text{and} \quad \tilde{a} = a - d_{A_0}\xi - [a, \xi] + \mathfrak{n}(\xi).$$

Here \mathfrak{n} and \mathfrak{m} denote expressions which are algebraic and at least quadratic in ξ . The gauge fixing condition (6.2) can thus be written as

$$\mathfrak{l}_\varepsilon \xi + \mathfrak{d}_\varepsilon \xi + \mathfrak{q}_\varepsilon(\xi) + \mathfrak{e}_\varepsilon = 0.$$

with

$$\begin{aligned} \mathfrak{l}_\varepsilon &:= \varepsilon^2 \Delta_{A_0 B} + R_{\Phi_0}^* R_{\Phi_0}, & \mathfrak{d}_\varepsilon &:= \varepsilon^2 d_{A_0 B}^*[a, \cdot] + \rho^*(\rho(\cdot)\phi\Phi_0^*), \\ \mathfrak{q}_\varepsilon(\xi) &:= \varepsilon^2 d_{A_0 B}^* \mathfrak{n}(\xi) + \rho^*(\mathfrak{m}(\xi)\Phi_0^*), & \mathfrak{e}_\varepsilon &:= -\varepsilon^2 d_{A_0 B}^* a - \rho^*(\phi\Phi_0). \end{aligned}$$

Denote by G_ε the Banach space $W^{k+3,p}\Omega^0(M, \mathfrak{g}_P)$ equipped with the norm

$$(6.3) \quad \|\xi\|_{G_\varepsilon} := \|\xi\|_{W^{k+1,p}} + \varepsilon\|\nabla^{k+2}\xi\|_{L^p} + \varepsilon^2\|\nabla^{k+3}\xi\|_{L^p}.$$

Since Φ_0 is regular, the operator $R_{\Phi_0}^* R_{\Phi_0}$ is positive definite; hence, for $\varepsilon \ll 1$, the operator

$$\mathfrak{l}_\varepsilon : G_\varepsilon \rightarrow W^{k+1,p}\Omega^0(M, \mathfrak{g}_P)$$

is invertible and $\|\mathfrak{l}_\varepsilon^{-1}\|_{\mathcal{L}(G_\varepsilon, W^{k+1,p})}$ is bounded independent of ε . Since

$$\|\mathfrak{d}_\varepsilon\|_{\mathcal{L}(G_\varepsilon, W^{k+1,p})} \lesssim \sigma \ll 1,$$

$\mathfrak{l}_\varepsilon + \mathfrak{d}_\varepsilon: G_\varepsilon \rightarrow W^{k+1,p}\Omega^0(M, \mathfrak{g}_P)$ will also be invertible with inverse bounded independent of ε and σ . Since the non-linearity $\mathfrak{q}_\varepsilon: G_\varepsilon \rightarrow W^{k+1,p}\Omega^0(M, \mathfrak{g}_P)$ satisfies (3.12) and $\|\mathfrak{e}_\varepsilon\| \lesssim \sigma \ll 1$, it follows from Banach's Fixed Point Theorem that, for a suitable $c > 0$, there exists a unique solution $\xi \in B_{c\sigma}(0) \subset G_\varepsilon$ to (6.3). This proves the existence of the desired gauge transformation, and local uniqueness. Global uniqueness follows by an argument by contradiction, cf. [DK90, Proposition 4.2.9]. \square

Proposition 6.4. *Let $\mathfrak{c}_0 = (\Phi_0, A_0)$ be a lift of a Fueter section $s_0 \in \Gamma(\mathfrak{X})$ for $\mathfrak{p}_0 \in \mathcal{P}$. Fix $\varepsilon > 0$ and $\mathfrak{p} \in \mathcal{P}$. Define a linear map $L_{\mathfrak{p},\varepsilon}: X_\varepsilon \rightarrow Y$ and a quadratic map $Q_{\mathfrak{p},\varepsilon}: X_0 \rightarrow Y$ by*

$$L_{\mathfrak{p},\varepsilon} := \begin{pmatrix} -\mathcal{D}_{A_0} & -\gamma(\cdot)\Phi_0 & -\rho(\cdot)\Phi_0 \\ -2 * \mu(\Phi_0, \cdot) & * \varepsilon^2 d_{A_0} & \varepsilon^2 d_{A_0} \\ -\rho^*(\cdot \Phi_0^*) & \varepsilon^2 d_{A_0}^* & 0 \\ 2\langle \Phi_0, \cdot \rangle_{L^2} & 0 & 0 \end{pmatrix} \quad \text{and}$$

$$Q_{\mathfrak{p},\varepsilon}(\phi, a, \xi) := \begin{pmatrix} -\bar{\gamma}(a)\phi \\ \frac{1}{2}\varepsilon^2 * [a \wedge a] - * \mu(\phi) \\ 0 \\ \|\phi\|_{L^2}^2 \end{pmatrix},$$

respectively. With $\mathfrak{e}_{\mathfrak{p},0}$ as in Proposition 5.2 set

$$\mathfrak{e}_{\mathfrak{p},\varepsilon} := \mathfrak{e}_{\mathfrak{p},0} + \varepsilon^2(0, * \omega F_{A_0}, 0).$$

There exist a neighborhood U of $\mathfrak{p}_0 \in \mathcal{P}$ and $\sigma > 0$ such that $\hat{\mathfrak{c}} = (\phi, a, \xi) \in B_\sigma(0) \subset X_\varepsilon$ satisfies

$$(6.5) \quad L_{\mathfrak{p},\varepsilon} \hat{\mathfrak{c}} + Q_{\mathfrak{p},\varepsilon}(\hat{\mathfrak{c}}) + \mathfrak{e}_{\mathfrak{p},\varepsilon} = 0$$

if and only if $\xi = 0$, $(A, \Phi) = (A_0 + a, \Phi_0 + \phi)$ satisfies

$$\mathcal{D}_A \Phi = 0, \quad \varepsilon^2 \omega F_A = \mu(\Phi), \quad \text{and} \quad \|\Phi\|_{L^2} = 1,$$

and

$$(6.6) \quad \varepsilon^2 d_{A_0}^* a - \rho^*(\phi \Phi_0^*) = 0.$$

Proof. We only need to show that ξ vanishes, but this follows from the same argument as in the proof of Proposition 5.2 because $d_A F_A = 0$. \square

Corollary 6.7. *There exist $\varepsilon, \sigma > 0$ such the map*

$$\{(\mathfrak{p}, \varepsilon, \phi, a, \xi) \in \mathcal{P} \times U \times (0, \varepsilon_0) \times B_\sigma(0) \text{ satisfying (6.5)}\} \rightarrow \mathfrak{M}_{\text{SW}}$$

defined by

$$(\mathfrak{p}, \varepsilon, \phi, a, \xi) \mapsto (\mathfrak{p}, \varepsilon, [(\Phi_0 + \phi, A_0 + a)])$$

is a homeomorphism onto the intersection of \mathfrak{M}_{SW} with a neighborhood of $([\mathfrak{c}_0], \mathfrak{p}_0, 0)$ in $\overline{\mathfrak{M}}_{\text{SW}}$.

6.2 Inverting $\bar{L}_{\mathbf{p},\varepsilon}$

Define the Banach space $(\bar{X}_\varepsilon, \|\cdot\|_{\bar{X}_\varepsilon})$ by

$$\bar{X}_\varepsilon := \text{coker}(d\bar{\mathfrak{F}})_{s_0} \oplus W^{k+1,p}\Gamma(\mathfrak{S}) \oplus W^{k+2,p}\Omega^1(M, \mathfrak{g}_P) \oplus W^{k+2,p}\Omega^0(M, \mathfrak{g}_P)$$

with norm

$$\|(o, \hat{c})\|_{\bar{X}_\varepsilon} := |o| + \|\hat{c}\|_{X_\varepsilon},$$

and the Banach space $(\bar{Y}, \|\cdot\|_{\bar{Y}})$ by

$$\bar{Y} := I_\partial \oplus \mathbf{R} \oplus W^{k,p}\Gamma(\mathfrak{S}) \oplus W^{k+1,p}\Omega^1(M, \mathfrak{g}_P) \oplus W^{k+1,p}\Omega^0(M, \mathfrak{g}_P)$$

with the obvious norm. Let $\bar{D}_{\mathfrak{S}}: \text{coker}(d\bar{\mathfrak{F}})_{s_0} \oplus W^{k+1,p}\Gamma(\mathfrak{S}) \rightarrow I_\partial \oplus \mathbf{R} \oplus W^{k,p}\Gamma(\mathfrak{S})$ be as in the Proof of Proposition 5.1. Define $\bar{L}_{\mathbf{p},\varepsilon}: \bar{X}_\varepsilon \rightarrow \bar{Y}$ by

$$(6.8) \quad \bar{L}_{\mathbf{p},\varepsilon} := \begin{pmatrix} -\bar{D}_{\mathfrak{S}} & \gamma\Pi^* & 0 \\ -\gamma\Pi & -\bar{D}_{\mathfrak{S}\mathfrak{t}} & -\alpha \\ 0 & -\alpha^* & \varepsilon^2\delta_{A_0} \end{pmatrix}$$

with

$$\delta_{A_0} := \begin{pmatrix} *d_{A_0} & d_{A_0} \\ d_{A_0}^* & 0 \end{pmatrix}.$$

Proposition 6.9. *There exist $\varepsilon_0, c > 0$, and a neighborhood U of $\mathbf{p}_0 \in \mathcal{P}$ such that, for all $\mathbf{p} \in U$ and $\varepsilon \in (0, \varepsilon_0]$, $\bar{L}_{\mathbf{p},\varepsilon}: \bar{X}_\varepsilon \rightarrow \bar{Y}$ is invertible, and $\|\bar{L}_{\mathbf{p},\varepsilon}^{-1}\| \leq c$.*

The proof of this result relies on the following two observations.

Proposition 6.10. *For $i = 1, 2, 3$, let V_i and W_i be Banach spaces, and set*

$$V := \bigoplus_{i=1}^3 V_i \quad \text{and} \quad W := \bigoplus_{i=1}^3 W_i.$$

Let $L: V \rightarrow W$ be a bounded linear operator of the form

$$L = \begin{pmatrix} D_1 & B_+ & 0 \\ B_- & D_2 & A_+ \\ 0 & A_- & D_3 \end{pmatrix}.$$

If the operators

$$\begin{aligned} D_1 &: V_1 \rightarrow W_1, \\ A_- &: V_2 \rightarrow W_3, \quad \text{and} \\ Z &:= A_+ - (D_2 - B_-D_1^{-1}B_+)A_-^{-1}D_3: V_3 \rightarrow W_2 \end{aligned}$$

are invertible, then there exists a bounded linear operator $R: W \rightarrow V$ such that

$$RL = \text{id}_W.$$

Moreover, the operator norm $\|R\|$ is bounded by a constant depending only on $\|L\|$, $\|D_1^{-1}\|$, $\|A_1^{-1}\|$, and $\|Z^{-1}\|$.

Proposition 6.11. *There exist $\varepsilon_0, c > 0$ such that for $\varepsilon \in (0, \varepsilon_0]$, the linear map*

$$\mathfrak{z}_\varepsilon := \mathfrak{a} + \varepsilon^2 \left(\mathcal{D}_{\mathfrak{N}} + \gamma \mathbb{I} \mathcal{D}_{\mathfrak{S}}^{-1} \gamma \mathbb{I}^* \right) (\mathfrak{a}^*)^{-1} \delta_{A_0} : W^{k+2,p} \Omega^1(M, \mathfrak{g}_P) \oplus W^{k+2,p} \Omega^0(M, \mathfrak{g}_P) \rightarrow W^{k,p} \Gamma(\mathfrak{N})$$

is invertible, and

$$\|\mathfrak{z}_\varepsilon^{-1}(a, \xi)\|_{W^{k,p}} + \varepsilon \|\nabla^{k+1} \mathfrak{z}_\varepsilon^{-1}(a, \xi)\|_{L^p} + \varepsilon^2 \|\nabla^{k+2} \mathfrak{z}_\varepsilon^{-1}(a, \xi)\|_{L^p} \leq c \|(a, \xi)\|_{W^{k,p}}.$$

Proof of Proposition 6.9. It suffices to prove the result for $\mathfrak{p} = \mathfrak{p}_0$, for then it follows for \mathfrak{p} close to \mathfrak{p}_0 .

Recall that

$$\begin{aligned} \bar{X}_\varepsilon &= \text{coker}(d\mathfrak{F})_{s_0} \oplus W^{k+1,p} \Gamma(\mathfrak{S}) \\ &\quad \oplus W^{k+1,p} \Gamma(\mathfrak{N}) \\ &\quad \oplus W^{k+2,p} \Omega^1(M, \mathfrak{g}_P) \oplus W^{k+2,p} \Omega^0(M, \mathfrak{g}_P), \\ \bar{Y} &= I_\partial \oplus \mathbf{R} \oplus W^{k,p} \Gamma(\mathfrak{S}) \\ &\quad \oplus W^{k,p} \Gamma(\mathfrak{N}) \\ &\quad \oplus W^{k+1,p} \Omega^1(M, \mathfrak{g}_P) \oplus W^{k+1,p} \Omega^0(M, \mathfrak{g}_P), \end{aligned}$$

and $\bar{L}_{\mathfrak{p}_0, \varepsilon}$ can be written as

$$\begin{pmatrix} -\bar{\mathcal{D}}_{\mathfrak{S}} & \gamma \mathbb{I}^* & 0 \\ -\gamma \mathbb{I} & -\bar{\mathcal{D}}_{\mathfrak{N}} & -\mathfrak{a} \\ 0 & -\mathfrak{a}^* & \varepsilon^2 \delta_{A_0} \end{pmatrix}$$

with

$$\delta_{A_0} = \begin{pmatrix} *d_{A_0} & d_{A_0} \\ d_{A_0}^* & 0 \end{pmatrix}.$$

The operators $\bar{\mathcal{D}}_{\mathfrak{S}}: \text{coker}(d\mathfrak{F})_{s_0} \oplus W^{k+1,p} \Gamma(\mathfrak{S}) \rightarrow I_\partial \oplus \mathbf{R} \oplus W^{k,p} \Gamma(\mathfrak{S})$ and $\mathfrak{a}^*: W^{k+1,p} \Gamma(\mathfrak{N}) \rightarrow W^{k+1,p} \Omega^1(M, \mathfrak{g}_P) \oplus W^{k+1,p} \Omega^0(M, \mathfrak{g}_P)$ both are invertible with uniformly bounded inverses, and by Proposition 6.11 the same holds for \mathfrak{z}_ε , provided $\varepsilon \ll 1$. Thus, according to Proposition 6.10, $\bar{L}_{\mathfrak{p}_0, \varepsilon}$ has a left inverse $R_\varepsilon: \bar{Y}_0 \rightarrow \bar{X}_\varepsilon$ whose norm can be bounded independent of ε .

To see that R_ε is also a right inverse, observe that $L_{\mathfrak{p}_0, \varepsilon}$ is a formally self-adjoint elliptic operator and, hence, $L_{\mathfrak{p}_0, \varepsilon}: X_\varepsilon \rightarrow Y$ is Fredholm of index zero. Consequently, $\bar{L}_{\mathfrak{p}_0, \varepsilon}$ is Fredholm of index zero. The existence of R_ε shows that $\ker \bar{L}_{\mathfrak{p}_0, \varepsilon} = 0$ and thus $\text{coker } \bar{L}_{\mathfrak{p}_0, \varepsilon} = 0$. By the Open Mapping Theorem, $\bar{L}_{\mathfrak{p}_0, \varepsilon}$ has an inverse $\bar{L}_{\mathfrak{p}_0, \varepsilon}^{-1}$. It must agree with R_ε since $R_\varepsilon = R_\varepsilon \bar{L}_{\mathfrak{p}_0, \varepsilon} \bar{L}_{\mathfrak{p}_0, \varepsilon}^{-1} = \bar{L}_{\mathfrak{p}_0, \varepsilon}^{-1}$. \square

Proof of Proposition 6.10. The left inverse of L can be found by Gauss elimination [Str16, Chapter 2]. The formula found in this way is rather unwieldy; fortunately, however, the precise formula is not needed.

Step 1. Set

$$E := (D_2 - B_- D_1^{-1} B_+) A_-^{-1} : W_3 \rightarrow W_2$$

The linear map $P : W \rightarrow V$ defined by

$$P := \begin{pmatrix} D_1^{-1} & 0 & 0 \\ 0 & 0 & A_-^{-1} \\ -Z^{-1} B_- D_1^{-1} & Z^{-1} & -Z^{-1} E \end{pmatrix}$$

satisfies

$$PL = \begin{pmatrix} \text{id}_{V_1} & D_1^{-1} B_+ & 0 \\ 0 & \text{id}_{V_2} & A_-^{-1} D_3 \\ 0 & 0 & \text{id}_{V_3} \end{pmatrix}.$$

Moreover, $\|P\|$ and $\|PL\|$ are bounded by a constant depending only $\|L\|$, $\|D_1^{-1}\|$, $\|A_-^{-1}\|$, and $\|Z^{-1}\|$.

This can be verified directly; alternatively, one can check that a sequence of row operations transforms the augmented matrix $(L \mid \text{id})$ as follows:

$$\begin{aligned} & \left(\begin{array}{ccc|ccc} D_1 & B_+ & 0 & \text{id}_{W_1} & 0 & 0 \\ B_- & D_2 & A_+ & 0 & \text{id}_{W_2} & 0 \\ 0 & A_- & D_3 & 0 & 0 & \text{id}_{W_3} \end{array} \right) \\ \rightsquigarrow & \left(\begin{array}{ccc|ccc} \text{id}_{V_1} & D_1^{-1} B_+ & 0 & D_1^{-1} & 0 & 0 \\ B_- & D_2 & A_+ & 0 & \text{id}_{W_2} & 0 \\ 0 & \text{id}_{V_2} & A_-^{-1} D_3 & 0 & 0 & A_-^{-1} \end{array} \right) \\ \rightsquigarrow & \left(\begin{array}{ccc|ccc} \text{id}_{V_1} & D_1^{-1} B_+ & 0 & D_1^{-1} & 0 & 0 \\ 0 & \text{id}_{V_2} & A_-^{-1} D_3 & 0 & 0 & A_-^{-1} \\ B_- & D_2 & A_+ & 0 & \text{id}_{W_2} & 0 \end{array} \right) \\ \rightsquigarrow & \left(\begin{array}{ccc|ccc} \text{id}_{V_1} & D_1^{-1} B_+ & 0 & D_1^{-1} & 0 & 0 \\ 0 & \text{id}_{V_2} & A_-^{-1} D_3 & 0 & 0 & A_-^{-1} \\ 0 & D_2 - B_- D_1^{-1} B_+ & A_+ & -B_- D_1^{-1} & \text{id}_{W_2} & 0 \end{array} \right) \\ \rightsquigarrow & \left(\begin{array}{ccc|ccc} \text{id}_{V_1} & D_1^{-1} B_+ & 0 & D_1^{-1} & 0 & 0 \\ 0 & \text{id}_{V_2} & A_-^{-1} D_3 & 0 & 0 & A_-^{-1} \\ 0 & 0 & Z & -B_- D_1^{-1} & \text{id}_{W_2} & -E \end{array} \right) \\ \rightsquigarrow & \left(\begin{array}{ccc|ccc} \text{id}_{V_1} & D_1^{-1} B_+ & 0 & D_1^{-1} & 0 & 0 \\ 0 & \text{id}_{V_2} & A_-^{-1} D_3 & 0 & 0 & A_-^{-1} \\ 0 & 0 & \text{id}_{V_3} & -Z^{-1} B_- D_1^{-1} & Z^{-1} & -Z^{-1} E \end{array} \right). \end{aligned}$$

Step 2. *The inverse of PL is*

$$(PL)^{-1} = \begin{pmatrix} \text{id}_{V_1} & -D_1^{-1}B_+ & D_1^{-1}B_+A_-^{-1}D_3 \\ 0 & \text{id}_{V_2} & -A_-^{-1}D_3 \\ 0 & 0 & \text{id}_{V_3} \end{pmatrix}.$$

Hence, $R := (PL)^{-1}P$ is the desired left inverse.

It can be verified directly that the above expression gives the inverse of PL . \square

Proof of Proposition 6.11. It suffices to show that the linear maps $\tilde{\mathfrak{z}}_\varepsilon := \alpha^* \mathfrak{z}_\varepsilon$ are uniformly invertible. A short computation using Proposition B.4 shows that

$$\tilde{\mathfrak{z}}_\varepsilon = \varepsilon^2 \delta_{A_0}^2 + \alpha^* \alpha + \varepsilon^2 \mathfrak{e}$$

where \mathfrak{e} is a zeroth order operator which factors through $W^{k+1,p} \rightarrow W^{k+1,p}$. Since Φ_0 is regular, $\alpha^* \alpha$ is positive definite and, hence, for $\varepsilon \ll 1$, $\alpha^* \alpha + \varepsilon^2 \delta_{A_0}^2$ is uniformly invertible. Since $\varepsilon \ll 1$, $\varepsilon^2 \mathfrak{e}$ is a small perturbation of order ε and thus $\tilde{\mathfrak{z}}_\varepsilon$ is uniformly invertible. \square

The above analysis yields the following regularity result, in which we decorate X_ε and Y with superscripts indicating the choice of the differentiability and integrability parameters k and p . The proof is almost identical to that of Proposition 5.11, and will be omitted.

Proposition 6.12. *For each $k, \ell \in \mathbf{N}$ and $p, q \in (1, \infty)$ with $(k+1)p > 3$, $\ell \geq k$, and $q \geq p$, there are constants $c, \sigma, \varepsilon_0 > 0$ and an open neighborhood U of \mathfrak{p}_0 in \mathcal{P} such that if $\varepsilon \in (0, \varepsilon_0]$, $\mathfrak{p} \in U$, and $\hat{c} \in B_\sigma(0) \subset X_\varepsilon^{k,p}$ is solution of*

$$L_{\mathfrak{p},\varepsilon} \hat{c} + Q_{\mathfrak{p},\varepsilon}(\hat{c}) + \mathfrak{e}_{\mathfrak{p},\varepsilon} = 0,$$

then $\hat{c} \in X_\varepsilon^{\ell,q}$ and $\|\hat{c}\|_{X_\varepsilon^{\ell,q}} \leq c \|\hat{c}\|_{X_\varepsilon^{k,p}}$.

6.3 Proof of Theorem 2.29

Since $Q_{\mathfrak{p},\varepsilon}$ is quadratic and

$$\begin{aligned} \|Q_{\mathfrak{p},\varepsilon}(\phi, a, \xi)\|_Y &\leq \|\bar{\gamma}(a)\phi\|_{W^{k,p}} + \varepsilon^2 \|[a \wedge a]\|_{W^{k+1,p}} + \|\mu(\phi)\|_{W^{k+1,p}} + \|\phi\|_{L^2}^2 \\ &\lesssim \|a\|_{W^{k,p}} \|\phi\|_{W^{k+1,p}} \\ &\quad + \left(\|a\|_{W^{k,p}} + \varepsilon \|\nabla^{k+1} a\|_{L^p} + \varepsilon^2 \|\nabla^{k+2} a\|_{L^p} \right)^2 + \|\phi\|_{W^{k+1,p}}^2, \end{aligned}$$

$Q_{\mathfrak{p},\varepsilon}$ satisfies (3.12), and because of Proposition 6.9 we can apply Lemma 3.11 to construct a smooth map $\text{ob}_\circ : U \times (0, \varepsilon_0) \times \mathcal{F}_\partial \rightarrow \text{coker}(d\mathfrak{F})_{s_0}$ and a map $\mathfrak{x}_\circ : \text{ob}^{-1}(0) \rightarrow \overline{\mathfrak{M}}_{\text{SW}}$ which is a homeomorphism onto the intersection of \mathfrak{M}_{SW} with a neighborhood of $[(A_0, \Phi_0)]$. (There is a slight caveat in the application of Lemma 3.11: the Banach space X_ε does depend on \mathfrak{p} and ε and Y depends on \mathfrak{p} . The dependence, however, is mostly harmless as different values of \mathfrak{p} and ε lead to naturally

isomorphic Banach spaces.) For what follows it will be important to know that maps ob_\circ and \mathfrak{x}_\circ are uniquely characterized as follows: for \mathbf{p} in the open neighborhood U of $\mathbf{p}_0 \in \mathcal{P}$, d in the open neighborhood \mathcal{J}_∂ of $0 \in I_\partial$, and $\varepsilon \in (0, \varepsilon_0)$, there is a unique solution $\bar{c} = \bar{c}(\mathbf{p}, \varepsilon, d) \in B_\sigma(0) \subset \bar{X}_\varepsilon$ of

$$(6.13) \quad \bar{L}_{\mathbf{p}, \varepsilon} \bar{c} + Q_{\mathbf{p}, \varepsilon}(\bar{c}) + \mathfrak{e}_{\mathbf{p}, \varepsilon} = d \in \mathcal{J}_\partial \subset \bar{Y};$$

$\text{ob}_\circ(\mathbf{p}, \varepsilon, d)$ is the component of $\bar{c}(\mathbf{p}, \varepsilon, d)$ in $\text{coker}(d\tilde{\mathfrak{Y}})_{s_0}$ and if $\text{ob}_\circ(\mathbf{p}, \varepsilon, d) = 0$ and \hat{c} denotes the component of $c(\mathbf{p}, \varepsilon, d)$ in X_ε , then $\mathfrak{x}_\circ(\mathbf{p}, \varepsilon, d) = c_0 + \hat{c}$. (Similar, setting $\varepsilon = 0$ yields ob_∂ and \mathfrak{x}_∂ .)

We define $\text{ob}: U \times [0, \varepsilon_0) \times \mathcal{J}_\partial \rightarrow \text{coker}(d\tilde{\mathfrak{Y}})_{s_0}$ by

$$\text{ob}(\cdot, \varepsilon, \cdot) = \begin{cases} \text{ob}_\circ(\cdot, \varepsilon, \cdot) & \text{for } \varepsilon \in (0, \varepsilon_0) \\ \text{ob}_\partial(\cdot, \cdot) & \text{for } \varepsilon = 0, \end{cases}$$

and $\mathfrak{x}: \text{ob}^{-1}(0) \rightarrow \bar{\mathfrak{M}}_{\text{SW}}$ by

$$\mathfrak{x}(\cdot, \varepsilon, \cdot) = \begin{cases} \mathfrak{x}_\circ(\cdot, \varepsilon, \cdot) & \text{for } \varepsilon \in (0, \varepsilon_0) \\ \mathfrak{x}_\partial(\cdot, \cdot) & \text{for } \varepsilon = 0. \end{cases}$$

In order to prove [Theorem 2.29](#) we need to understand the regularity of ob near $\varepsilon = 0$; in other words: we need to understand how ob_\circ and ob_∂ fit together.

Let $k \in \mathbb{N}$ and $p \in (1, \infty)$ be the differentiability and integrability parameters used in the definition of \bar{X}_ε . If necessary, shrink U and \mathcal{J}_∂ and decrease σ so that the proof of [Proposition 5.1](#) goes through and [Proposition 5.2](#) holds with differentiability parameter $k + 2r + 2$ and integrability parameter p . Observe that $\bar{X}_0^{k+2, p} \subset \bar{X}_\varepsilon$ and the norm of the inclusion can be bounded by a constant independent of ε .

Proposition 6.14. *For every $(\mathbf{p}, d) \in U \times \mathcal{J}_\partial$, there are $\bar{c}_0(\mathbf{p}, d) \in \bar{X}_0^{k+2r+2, p}$ and $\hat{c}_i(\mathbf{p}, d) \in \bar{X}_0^{k+2(r-i)+2, p}$ (for $i = 1, \dots, r$) depending smoothly on \mathbf{p} and d , such that, for $m, n \in \mathbb{N}$ with $m + n \leq 2r$,*

$$\tilde{c}(\mathbf{p}, \varepsilon, d) := \bar{c}_0 + \sum_{i=1}^r \varepsilon^{2i} \hat{c}_i$$

satisfies

$$(6.15) \quad \left\| \nabla_{U \times \mathcal{J}_\partial}^m \partial_\varepsilon^n (\bar{c}(\mathbf{p}, \varepsilon, d) - \tilde{c}(\mathbf{p}, \varepsilon, d)) \right\|_{\bar{X}_\varepsilon} = O(\varepsilon^{2k+2-n}).$$

Proof. We construct \tilde{c} by expanding (6.13) in ε^2 . To this end we write

$$\bar{L}_{\mathbf{p}, \varepsilon} = \bar{L}_{\mathbf{p}, 0} + \varepsilon^2 \ell_{\mathbf{p}}, \quad Q_{\mathbf{p}, \varepsilon} = Q_{\mathbf{p}, 0} + \varepsilon^2 q_{\mathbf{p}}, \quad \text{and} \quad \mathfrak{e}_{\mathbf{p}, \varepsilon} = \mathfrak{e}_{\mathbf{p}, 0} + \varepsilon^2 \hat{\mathfrak{e}}_{\mathbf{p}},$$

with

$$\ell_{\mathbf{p}} := \begin{pmatrix} 0 & & \\ & 0 & \\ & & \delta_{A_0} \end{pmatrix}, \quad q_{\mathbf{p}}(\phi, a, \xi) := \begin{pmatrix} 0 & & \\ & 0 & \\ \frac{1}{2} * [a \wedge a] & & \end{pmatrix}, \quad \text{and} \quad \hat{\mathfrak{e}}_{\mathbf{p}} := \begin{pmatrix} 0 & \\ 0 & \\ * \omega_{F_{A_0}} & \end{pmatrix}.$$

Observe that $\ell_{\mathbf{p}}: \bar{X}_0^{\ell, p} \rightarrow \bar{Y}^{\ell-2, p}$ is a bounded linear map and $q_{\mathbf{p}}: \bar{X}_0^{\ell, p} \rightarrow \bar{Y}^{\ell-2, p}$ is a bounded quadratic map.

Step 1. *Construction of \bar{c}_0 and \hat{c}_i .*

By Banach's Fixed Point Theorem, there is a unique solution $\bar{c}_0 \in B_\sigma(0) \subset \bar{X}_0^{k+2r+2,p}$ of

$$\bar{L}_{\mathbf{p},0}\bar{c}_0 + Q_{\mathbf{p},0}(\bar{c}_0) + \mathfrak{e}_{\mathbf{p},0} = d \in \mathcal{J}_\partial \subset \bar{Y}^{k+2r+2}.$$

Moreover, \bar{c}_0 actually lies in $B_{\sigma/2}(0) \subset \bar{X}_0^{k+2r+2,p}$ provided U and \mathcal{J}_∂ have been chosen sufficiently small. We have

$$\bar{L}_{\mathbf{p},\varepsilon}\bar{c}_0 + Q_{\mathbf{p},0}(\bar{c}_\varepsilon) + \mathfrak{e}_{\mathbf{p},\varepsilon} - d = \varepsilon^2 r_0(\mathbf{p}, d) \in \bar{Y}^{k+2(r-1)+2,p}.$$

with

$$r_0(\mathbf{p}, d) := \ell_{\mathbf{p}}\bar{c}_0 + q_{\mathbf{p}}(\bar{c}_0) + \hat{c}_{\mathbf{p}}.$$

Since $\sigma \ll 1$, the operator $\bar{L}_{\mathbf{p},0} + 2Q_{\mathbf{p},0}(\bar{c}_0, \cdot): \bar{X}_0^{k+2(r-i)+2,p} \rightarrow \bar{Y}_0^{k+2(r-i)+2,p}$ is invertible for $i = 1, \dots, r$.⁷ Recursively define $r_i(\mathbf{p}, d) \in \bar{Y}^{k+2(r-i)+2,p}$ by

$$\varepsilon^{2i+2} r_i := \bar{L}_{\mathbf{p},\varepsilon}\bar{c}_\varepsilon^i + Q_{\mathbf{p},0}(\bar{c}_\varepsilon^i) + \mathfrak{e}_{\mathbf{p},\varepsilon} - d$$

with

$$\tilde{c}(\varepsilon, \mathbf{p}, d) := \bar{c}_0 + \varepsilon^2 \hat{c}_1 + \dots + \varepsilon^{2i} \hat{c}_i,$$

and define $\hat{c}_{i+1} \in \bar{X}_0^{k+2(r-i-1)+2}$ to be the unique solution of

$$\bar{L}_{\mathbf{p},0}\hat{c}_{i+1} + 2Q_{\mathbf{p},0}(\bar{c}_0, \hat{c}_{i+1}) = r_i.$$

Clearly, $\bar{c}_0, \hat{c}_1, \dots, \hat{c}_r$ depend smoothly on \mathbf{p} and d .

Step 2. *We prove (6.15).*

We have

$$(6.16) \quad \bar{L}_{\mathbf{p},\varepsilon}\bar{c}_\varepsilon + Q_{\mathbf{p},\varepsilon}(\bar{c}_\varepsilon) - \bar{L}_{\mathbf{p},\varepsilon}\tilde{c} - Q_{\mathbf{p},\varepsilon}(\tilde{c}) = -\varepsilon^{2k+2} r$$

with $r = r_r$ as in the previous step. Both \bar{c} and \tilde{c} are small in \bar{X}_ε ; hence, it follows that

$$\|\bar{c} - \tilde{c}\|_{\bar{X}_\varepsilon} = O(\varepsilon^{2k+2}).$$

To obtain estimates for the derivatives of $\bar{c} - \tilde{c}$, we differentiate (6.16) and obtain an identity whose left-hand side is

$$\bar{L}_{\mathbf{p},0}\nabla^m \partial_\varepsilon^n (\bar{c} - \tilde{c}) + 2Q_{\mathbf{p},0}(\bar{c}, \nabla^m \partial_\varepsilon^n (\bar{c} - \tilde{c})) + 2Q_{\mathbf{p},0}(\bar{c} - \tilde{c}, \nabla^m \partial_\varepsilon^n \bar{c})$$

and whose right-hand side can be controlled in terms of the lower order derivatives of δ_ε^k . This gives the asserted estimates. \square

From Proposition 6.14 it follows that \mathfrak{x} is a homeomorphism onto its image and that the estimate in Theorem 2.29(1) holds with $\widehat{\text{ob}}_i$ denoting the component of \hat{c}_i in $\text{coker}(d\mathfrak{Y})_{s_0}$. This expansion implies that ob is C^{2r-1} up to $\varepsilon = 0$. \square

⁷Here we engage in the slight abuse of notation to use the same notation for a bilinear map and its associated quadratic form.

7 Proof of Theorem 2.31

The first part of Theorem 2.31 follows directly from Theorem 2.29, since in this situation

$$\text{ob}(\varepsilon, t) = \dot{\lambda}(0) \cdot t + O(t^2) + O(\varepsilon^2)$$

because $\text{ob}_\partial(t) = \dot{\lambda}(0) \cdot t + O(t^2)$. The second part requires a more detailed analysis to show that

$$\text{ob}(\varepsilon, t) = \dot{\lambda}(0) \cdot t - \delta\varepsilon^4 + O(t^2) + O(\varepsilon^6).$$

To establish the above expansion of ob , we solve

$$\bar{L}_\varepsilon(o_\varepsilon, \hat{c}) + Q_\varepsilon(\hat{c}) + \begin{pmatrix} 0 \\ 0 \\ \varepsilon^2 * \omega F_{A_0} \\ 0 \end{pmatrix} = 0$$

by formally expanding in ε^2 . Inspection of (5.9) shows that the obstruction to being able to solve $L_0\hat{c} = (\psi, b, \eta)$ is

$$-\pi(\psi + \gamma\Pi(\alpha^*)^{-1}(b, \eta))$$

where π denotes the L^2 -orthogonal projection onto $\ker \mathcal{D}_{\mathfrak{S}}$. In the case at hand, $\ker \mathcal{D}_{\mathfrak{S}} = \mathbf{R}\langle\Phi_0\rangle$, and we have

$$\begin{aligned} \langle\Phi_0, \gamma\Pi^*(\alpha^*)^{-1}(b, \eta)\rangle_{L^2} &= \sum_{i=1}^3 \langle\Phi_0, \gamma(e_i)\nabla_{e_i}(\alpha^*)^{-1}(b, \eta)\rangle_{L^2} \\ &= \sum_{i=1}^3 \langle\gamma(e_i)\nabla_{e_i}\Phi_0, (\alpha^*)^{-1}(b, \eta)\rangle_{L^2} = 0 \end{aligned}$$

since $\alpha: \Omega^1(M, \mathfrak{g}_P) \oplus \Omega^0(M, \mathfrak{g}_P) \rightarrow \Gamma(\mathfrak{A})$ and thus $(\alpha^*)^{-1}$ also maps to $\Gamma(\mathfrak{A})$. Thus the obstruction reduces to

$$-\langle\Phi_0, \psi\rangle_{L^2}.$$

By (5.9), the solution to $L_0(\phi, a, \xi) = (0, * \omega F_{A_0}, 0)$ is

$$(7.1) \quad \begin{aligned} \phi &= -\mathcal{D}_{\mathfrak{S}}^{-1} \gamma\Pi^* \chi - \chi, \quad \text{and} \\ (a, \xi) &= \alpha^{-1} \mathcal{D}_{\mathfrak{A}} \chi + \alpha^{-1} \gamma\Pi \mathcal{D}_{\mathfrak{S}}^{-1} \gamma\Pi^* \chi \end{aligned}$$

with

$$(7.2) \quad \chi := (\alpha^*)^{-1} * \omega F_{A_0}.$$

Setting $\hat{c}_0 := \varepsilon^2(\phi, a, \xi)$, we have

$$\varepsilon^4 \hat{\mathfrak{d}}_1 := \bar{L}_\varepsilon(0, \hat{c}_0) + Q_\varepsilon(\hat{c}_0) + (0, 0, \varepsilon^2 * \omega F_{A_0}, 0) = O(\varepsilon^4).$$

The component of $\hat{\mathfrak{d}}_1$ in $\Gamma(\mathfrak{S})$ is

$$-\bar{\gamma}(a)\phi.$$

Using $\bar{\gamma}(a)\Phi_0 \in \Gamma(\mathfrak{R})$ and $\rho(\mathfrak{g}_P)\Phi \perp \chi$, we find that the obstruction to being able to solve $L_0(\phi_1, a_1, \xi_1) = \hat{\mathfrak{d}}_1$ is

$$\begin{aligned} \mathfrak{v} &:= \langle \Phi_0, \bar{\gamma}(a)\phi \rangle_{L^2} = \langle \bar{\gamma}(a)\Phi_0, \phi \rangle_{L^2} \\ &= -\langle \bar{\gamma}(a)\Phi_0, \chi \rangle_{L^2} \\ &= -\langle \mathfrak{a}(a, \xi), \chi \rangle_{L^2} \\ &= -\langle \mathcal{D}_{\mathfrak{R}}\chi + \gamma\Pi\mathcal{D}_{\mathfrak{S}}^{-1}\gamma\Pi^*\chi, \chi \rangle_{L^2} \\ &= -\langle \mathcal{D}_{\mathfrak{R}}\chi, \chi \rangle_{L^2} + \langle \mathcal{D}_{\mathfrak{S}}^{-1}\gamma\Pi^*\chi, \gamma\Pi^*\chi \rangle_{L^2}. \end{aligned}$$

Comparing this with

$$\begin{aligned} \langle \mathcal{D}_{A_0}\phi, \phi \rangle_{L^2} &= \langle \mathcal{D}_{A_0}\mathcal{D}_{\mathfrak{S}}^{-1}\gamma\Pi^*\chi + \mathcal{D}_{A_0}\chi, \mathcal{D}_{\mathfrak{S}}^{-1}\gamma\Pi^*\chi + \chi \rangle_{L^2} \\ &= \langle (\mathcal{D}_{\mathfrak{S}} + \gamma\Pi)\mathcal{D}_{\mathfrak{S}}^{-1}\gamma\Pi^*\chi + (\mathcal{D}_{\mathfrak{R}} - \gamma\Pi^*)\chi, \mathcal{D}_{\mathfrak{S}}^{-1}\gamma\Pi^*\chi + \chi \rangle_{L^2} \\ &= \langle \gamma\Pi^*\chi, \mathcal{D}_{\mathfrak{S}}^{-1}\gamma\Pi^*\chi \rangle_{L^2} + \langle \gamma\Pi\mathcal{D}_{\mathfrak{S}}^{-1}\gamma\Pi^*\chi, \chi \rangle_{L^2} \\ &\quad + \langle \mathcal{D}_{\mathfrak{R}}\chi, \chi \rangle_{L^2} - \langle \gamma\Pi^*\chi, \mathcal{D}_{\mathfrak{S}}^{-1}\gamma\Pi^*\chi \rangle_{L^2} \\ &= -\langle \mathcal{D}_{\mathfrak{S}}^{-1}\gamma\Pi^*\chi, \gamma\Pi^*\chi \rangle_{L^2} + \langle \mathcal{D}_{\mathfrak{R}}\chi, \chi \rangle_{L^2} \\ &= -\mathfrak{v} \end{aligned}$$

completes the proof. \square

A Examples of Seiberg–Witten equations

Example A.1. Let $G = \mathrm{U}(n)$ and $S = \mathbf{H} \otimes_{\mathbf{C}} \mathbf{C}^n$, where the complex structure on \mathbf{H} is given by right-multiplication by i . Let $\rho: \mathrm{U}(n) \rightarrow \mathrm{Sp}(\mathbf{H} \otimes_{\mathbf{C}} \mathbf{C}^n)$ be induced from the standard representation of $\mathrm{U}(n)$. The corresponding Seiberg–Witten equation is the $\mathrm{U}(n)$ –**monopole equation** in dimension three. The closely related $\mathrm{PU}(2)$ –monopole equation on 4-manifolds plays a crucial role in Pidstrigach and Tyurin’s approach to proving Witten’s conjecture relating Donaldson and Seiberg–Witten invariants; see, e.g., [PT95; FL98; Teloo].

In this example as well as in Example 2.15, we have $\mu^{-1}(0) = \{0\}$.

Example A.2. Let G be a compact Lie group, $\mathfrak{g} = \mathrm{Lie}(G)$, and fix an Ad –invariant inner product on \mathfrak{g} . $S := \mathbf{H} \otimes_{\mathbf{R}} \mathfrak{g}$ is a quaternionic Hermitian vector space, and $\rho: G \rightarrow \mathrm{Sp}(S)$ induced by the adjoint action is a quaternionic representation. The moment map $\mu: \mathbf{H} \otimes_{\mathbf{R}} \mathfrak{g} \rightarrow (\mathrm{Im} \mathbf{H} \otimes \mathfrak{g})^*$ is given by

$$\begin{aligned} \mu(\xi) &= \frac{1}{2}[\xi, \xi] \\ &= ([\xi_2, \xi_3] + [\xi_0, \xi_1]) \otimes i + ([\xi_3, \xi_1] + [\xi_0, \xi_2]) \otimes j + ([\xi_1, \xi_2] + [\xi_0, \xi_3]) \otimes k \end{aligned}$$

for $\xi = \xi_0 \otimes 1 + \xi_1 \otimes i + \xi_2 \otimes j + \xi_3 \otimes k \in \mathbf{H} \otimes_{\mathbf{R}} \mathfrak{g}$. Set $H := \mathrm{Sp}(1) \times G$ and extend the above quaternionic representation of G to H by declaring that $q \in \mathrm{Sp}(1)$ acts by right-multiplication with q^* .

Taking Q to be the product of the chosen spin structure \mathfrak{s} with a principal G -bundle, and choosing B such that it induces the spin connection on \mathfrak{s} , (2.14) becomes

$$\begin{aligned} d_A^* a &= 0, \\ *d_A a + d_A \xi &= 0, \quad \text{and} \\ F_A &= \frac{1}{2}[a \wedge a] + *[\xi, a]. \end{aligned}$$

for $\xi \in \Gamma(\mathfrak{g}_P)$, $a \in \Omega^1(M, \mathfrak{g}_P)$ and $A \in \mathcal{A}(P)$. If M is closed, then integration by parts shows that every solution of this equation satisfies $d_A \xi = 0$ and $[\xi, a] = 0$; hence, $A + ia$ defines a **flat $G^{\mathbf{C}}$ -connection**. Here $G^{\mathbf{C}}$ denotes the complexification of G .

In the above situation, we have $\mu^{-1}(0)/G \cong (\mathbf{H} \otimes \mathfrak{t})/W$ where \mathfrak{t} is the Lie algebra of a maximal torus $T \subset G$ and $W = N_G(T)/T$ is the Weyl group of G . However, since each $\xi \in \mu^{-1}(0)$ has stabilizer conjugate to T , we have $\mu^{-1}(0) \cap S^{\mathrm{reg}} = \emptyset$, and the hyperkähler quotient $S^{\mathrm{reg}} // G$ is empty.

Example A.3. The motivating example for us is the (r, k) **ADHM Seiberg–Witten equation**, which we expect to play an important role in gauge theory on G_2 -manifolds,⁸ and which arises from

$$S = \mathrm{Hom}_{\mathbf{C}}(\mathbf{C}^r, \mathbf{H} \otimes_{\mathbf{C}} \mathbf{C}^k) \oplus \mathbf{H}^* \otimes_{\mathbf{R}} \mathfrak{u}(k)$$

with

$$G = \mathrm{U}(k) \triangleleft H = \mathrm{SU}(r) \times \mathrm{Sp}(1) \times \mathrm{U}(k)$$

where $\mathrm{SU}(r)$ acts on \mathbf{C}^r in the obvious way, $\mathrm{U}(k)$ acts on \mathbf{C}^k in the obvious way and on $\mathfrak{u}(k)$ by the adjoint representation, and $\mathrm{Sp}(1)$ acts on the first copy of \mathbf{H} trivially and on the second copy by right-multiplication with the conjugate. According to Atiyah, Hitchin, Drinfeld, and Manin [AHDM78], if $r \geq 2$, then $S^{\mathrm{reg}} // G$ is the moduli space of framed $\mathrm{SU}(r)$ ASD instantons of charge k on \mathbf{R}^4 , and $\mu^{-1}(0)/G$ is its Uhlenbeck compactification. If $r = 1$, then $\mu^{-1}(0) \cap S^{\mathrm{reg}} = \emptyset$, and $\mu^{-1}(0)/G = \mathrm{Sym}^k \mathbf{H} := \mathbf{H}^k / S_k$ by Nakajima [Nak99, Example 3.14].

B Useful identities involving μ

This appendix summarizes and proves a few useful identities regarding μ , some of which are used in this article.

Proposition B.1. *For $\xi \in \Omega^0(M, \mathfrak{g}_P)$, $a \in \Omega^1(M, \mathfrak{g}_P)$, and $\phi, \psi \in \Gamma(\mathfrak{S})$, we have*

$$(B.2) \quad [\xi, \mu(\phi, \psi)] = \mu(\phi, \rho(\xi)\psi) + \mu(\psi, \rho(\xi)\phi),$$

⁸More precisely, we expect solutions of the (r, k) ADHM Seiberg–Witten equation to play a role in counter-acting the bubbling phenomenon along associative submanifolds discussed in [DS11; Wal17a]; see also [Hay17].

and for $a \in \Omega^1(M, \mathfrak{g}_P)$ and $\phi, \psi \in \Gamma(\Xi)$, we have

$$(B.3) \quad 2[a \wedge \mu(\phi, \psi)] = - * \rho^* ((\bar{\gamma}(a)\phi)\psi^*) - * \rho^* ((\bar{\gamma}(a)\psi)\phi^*).$$

Proof. For all $a \in \Omega^1(M, \mathfrak{g}_P)$, we have

$$\begin{aligned} 2\langle [\xi, \mu(\phi, \psi)], *a \rangle &= \langle \mu(\phi, \psi), - * [\xi, a] \rangle \\ &= \langle \phi, -\bar{\gamma}([\xi, a])\psi \rangle \\ &= -\langle \phi, \rho(\xi)\bar{\gamma}(a)\psi \rangle + \langle \phi, \bar{\gamma}(a)\rho(\xi)\psi \rangle \\ &= \langle \rho(\xi)\phi, \bar{\gamma}(a)\psi \rangle + \langle \phi, \bar{\gamma}(a)\rho(\xi)\psi \rangle \\ &= 2\langle \mu(\phi, \rho(\xi)\psi), *a \rangle + 2\langle \mu(\psi, \rho(\xi)\phi), *a \rangle. \end{aligned}$$

This proves the first identity. To prove the second identity, note that, for all $\eta \in \Omega^0(M, \mathfrak{g}_P)$, we have

$$\begin{aligned} 2\langle [a \wedge \mu(\phi, \psi)], *\eta \rangle &= \langle 2\mu(\phi), *[\eta, a] \rangle \\ &= \langle \phi, \bar{\gamma}([\eta, a])\psi \rangle \\ &= \langle \phi, \rho(\xi)\bar{\gamma}(a)\psi \rangle - \langle \phi, \bar{\gamma}(a)\rho(\xi)\psi \rangle \\ &= -\langle \xi, \rho^* ((\bar{\gamma}(a)\psi)\phi^*) \rangle - \langle \xi, \rho^* ((\bar{\gamma}(a)\phi)\psi^*) \rangle. \quad \square \end{aligned}$$

Proposition B.4. For all $A \in \mathcal{A}(Q)$ and $\phi, \psi \in \Gamma(\Xi)$ we have

$$(B.5) \quad d_A \mu(\phi, \psi) = - * \frac{1}{2} \rho^* ((\mathcal{D}_A \phi)\psi^* + (\mathcal{D}_A \psi)\phi^*)$$

and

$$(B.6) \quad \begin{aligned} d_A^* \mu(\phi, \psi) &= * \mu(\mathcal{D}_A \phi, \psi) + * \mu(\mathcal{D}_A \psi, \phi) \\ &\quad - \frac{1}{2} \rho^* ((\nabla_A \phi)\psi^*) - \frac{1}{2} \rho^* ((\nabla_A \psi)\phi^*). \end{aligned}$$

Proof. Fix a point $x \in M$, a positive local orthonormal frame (e_i) around x with $(\nabla e_i)(x) = 0$, and let ξ be a local section of \mathfrak{g}_P defined in a neighborhood of x satisfying $(\nabla \xi)(x) = 0$. We set $\nabla_i^A := \nabla_{e_i}^A$.

At the point $x \in M$, we compute with

$$\begin{aligned} \langle d_A \mu(\phi, \psi), *\xi \rangle &= -\langle d_A^* * \mu(\phi, \psi), \xi \rangle \\ &= \frac{1}{2} \sum_{i=1}^3 \nabla_i^A \langle \bar{\gamma}(\xi \otimes e^i)\phi, \psi \rangle \\ &= \frac{1}{2} (\langle \rho(\xi)\mathcal{D}_A \phi, \psi \rangle + \langle \phi, \rho(\xi)\mathcal{D}_A \psi \rangle) \\ &= -\frac{1}{2} \langle \xi, (\mathcal{D}_A \phi)\psi^* + (\mathcal{D}_A \psi)\phi^* \rangle. \end{aligned}$$

This proves the first identity. To prove the second identity, we compute

$$\begin{aligned}
\langle d_A^* \mu(\phi, \psi), \xi \rangle &= \langle * d_A * \mu(\phi, \psi), \xi \rangle \\
&= * \frac{1}{2} \sum_{i,j=1}^3 \nabla_i^A \langle \bar{\gamma}(\xi \otimes e^j) \phi, \psi \rangle e^i \wedge e^j \\
&= * \frac{1}{2} \sum_{i,j=1}^3 \left(\langle \bar{\gamma}(\xi \otimes e^j) \nabla_i^A \phi, \psi \rangle + \langle \bar{\gamma}(\xi \otimes e^j) \nabla_i^A \psi, \phi \rangle \right) e^i \wedge e^j \\
&= \frac{1}{2} \sum_{i,j,k=1}^3 \varepsilon_{ijk}^2 \left(\langle \rho(\xi) \gamma(e^k) \gamma(e^i) \nabla_i^A \phi, \psi \rangle \right. \\
&\quad \left. + \langle \rho(\xi) \gamma(e^k) \gamma(e^i) \nabla_i^A \psi, \phi \rangle \right) e^k \\
&= \frac{1}{2} \sum_{k=1}^3 \left(\langle \bar{\gamma}(\xi \otimes e^k) \mathbb{D}_A \phi, \psi \rangle + \langle \bar{\gamma}(\xi \otimes e^k) \mathbb{D}_A \psi, \phi \rangle \right. \\
&\quad \left. + \langle \rho(\xi) \nabla_k^A \phi, \psi \rangle + \langle \rho(\xi) \nabla_k^A \psi, \phi \rangle \right) e^k \\
&= \langle \xi, * \mu(\mathbb{D}_A \phi, \psi) \rangle + \langle \xi, * \mu(\mathbb{D}_A \psi, \phi) \rangle \\
&\quad + \frac{1}{2} \langle \rho(\xi) \nabla_A \phi, \psi \rangle + \frac{1}{2} \langle \rho(\xi) \nabla_A \psi, \phi \rangle. \quad \square
\end{aligned}$$

Proposition B.7. *If $(\varepsilon, \Phi, A) \in (0, \infty) \times \Gamma(\mathfrak{S}) \times \mathcal{A}_B(Q)$ is a solution of (2.21) and $R_\Phi(\xi) = \rho(\xi)\Phi$, then*

$$\begin{aligned}
(d_A^* d_A + d_A d_A^* + \varepsilon^{-2} R_\Phi^* R_\Phi) \mu(\Phi) &= \sum_{i,j=1}^3 \frac{1}{2} \rho^* \left(((F_{ij}^B + F_{ij}^S) \cdot \Phi) \Phi^* \right) e^{ij} \\
&\quad + \rho^* \left((\nabla_j^A \Phi) (\nabla_i^A \Phi)^* \right) e^{ij}.
\end{aligned}$$

Here (e_1, e_2, e_3) is local orthonormal frame, (e^1, e^2, e^3) is the dual coframe, $F_{ij}^B := F_B(e_i, e_j)$, $F_{ij}^S := F_S(e_i, e_j)$ with F_S denoting the curvature of the spin connection on \mathfrak{s} , and $e^{ij} := e^i \wedge e^j$.

Proof. We compute

$$\begin{aligned}
d_A \rho^* [(\nabla_A \Phi) \Phi^*] &= \sum_{i,j=1}^3 \rho^* [(\nabla_i^A \nabla_j^A \Phi) \Phi^*] e^{ij} + \rho^* [(\nabla_j^A \Phi) (\nabla_i^A \Phi)^*] e^{ij} \\
&= \sum_{i,j=1}^3 \frac{1}{2} \rho^* [(F_{ij}^A \cdot \Phi) \Phi^*] e^{ij} + \rho^* [(\nabla_j^A \Phi) (\nabla_i^A \Phi)^*] e^{ij}.
\end{aligned}$$

Since

$$\sum_{i,j=1}^3 \rho^* [\rho(\mu(\Phi)_{ij}) \Phi] e^{ij} = R_\Phi^* R_\Phi \mu(\Phi),$$

the result now follows from Proposition B.4. □

C Proof of Proposition 3.2

For the reader's convenience, we recall the definitions of the graded vector space L^\bullet ,

$$\begin{aligned} L^0 &:= \Omega^0(M, \mathfrak{g}_P), \\ L^1 &:= \Gamma(\mathfrak{S}) \oplus \Omega^1(M, \mathfrak{g}_P), \\ L^2 &:= \Gamma(\mathfrak{S}) \oplus \Omega^2(M, \mathfrak{g}_P), \quad \text{and} \\ L^3 &:= \Omega^3(M, \mathfrak{g}_P), \end{aligned}$$

the graded Lie bracket $[[\cdot, \cdot]]$,

$$\begin{aligned} [[a, b]] &:= [a \wedge b] && \text{for } a, b \in \Omega^\bullet(M, \mathfrak{g}_P), \\ [[\xi, \phi]] &:= \rho(\xi)\phi && \text{for } \xi \in \Omega^0(M, \mathfrak{g}_P) \text{ and } \phi \in \Gamma(\mathfrak{S}) \text{ in degree 1 or 2,} \\ [[a, \phi]] &:= -\bar{\gamma}(a)\phi && \text{for } a \in \Omega^1(M, \mathfrak{g}_P) \text{ and } \phi \in \Gamma(\mathfrak{S}) \text{ in degree 1,} \\ [[\phi, \psi]] &:= -2\mu(\phi, \psi) && \text{for } \phi, \psi \in \Gamma(\mathfrak{S}) \text{ in degree 1, and} \\ [[\phi, \psi]] &:= -*\rho^*(\phi\psi^*) && \text{for } \phi \in \Gamma(\mathfrak{S}) \text{ in degree 1 and } \psi \in \Gamma(\mathfrak{S}) \text{ in degree 2,} \end{aligned}$$

and the graded differential δ_c ,

$$\begin{aligned} \delta_c^0(\xi) &:= \begin{pmatrix} -\rho(\xi)\Phi \\ d_A\xi \end{pmatrix}, \\ \delta_c^1(\phi, a) &:= \begin{pmatrix} -\mathcal{D}_A\phi - \bar{\gamma}(a)\Phi \\ -2\mu(\Phi, \phi) + d_Aa \end{pmatrix}, \quad \text{and} \\ \delta_c^2(\psi, b) &:= *\rho^*(\psi\Phi^*) + d_Ab. \end{aligned}$$

We proceed in four steps.

Step 1. $(L^\bullet, [[\cdot, \cdot]])$ is a graded Lie algebra.

We need to verify the graded Jacobi identity, that is, for every three homogeneous elements $x, y, z \in L^\bullet$ we need to show that

$$J(x, y, z) := (-1)^{\deg x \cdot \deg z} [[x, [[y, z]]] + (-1)^{\deg y \cdot \deg x} [[y, [[z, x]]] + (-1)^{\deg z \cdot \deg y} [[z, [[x, y]]]$$

vanishes. Here $\deg x$ denotes the degree of x .

For degree reasons $J(x, y, z) = 0$, unless $\deg x + \deg y + \deg z \leq 3$. $(\Omega^\bullet(M, \mathfrak{g}_P), [\cdot \wedge \cdot])$ is a graded Lie algebra. Since $J(x, y, z)$ is invariant under permutations of x, y , and z , we can assume that $z \in \Gamma(\mathfrak{S})$ in degree 1 or 2. Hence, only the following five cases remain:

- For $\xi, \eta \in \Omega^0(M, \mathfrak{g}_P)$, and $\phi \in \Gamma(\mathfrak{S})$ in degree 1 or 2, we have

$$\begin{aligned} J(\xi, \eta, \phi) &= [[\xi, [[\eta, \phi]]] + [[\eta, [[\phi, \xi]]] + [[\phi, [[\xi, \eta]]] \\ &= \rho(\xi)\rho(\eta)\phi - \rho(\eta)\rho(\xi)\phi - \rho([\xi, \eta])\phi = 0. \end{aligned}$$

- For $\xi \in \Omega^0(M, \mathfrak{g}_P)$, and $\phi, \psi \in \Gamma(\Xi)$ in degree 1, we have

$$\begin{aligned} J(\xi, \phi, \psi) &= \llbracket \xi, \llbracket \phi, \psi \rrbracket \rrbracket + \llbracket \phi, \llbracket \psi, \xi \rrbracket \rrbracket - \llbracket \psi, \llbracket \xi, \phi \rrbracket \rrbracket \\ &= -2\llbracket \xi, \mu(\phi, \psi) \rrbracket + 2\mu(\phi, \rho(\xi)\psi) + 2\mu(\psi, \rho(\xi)\phi) = 0 \end{aligned}$$

by Proposition B.1.

- For $\xi \in \Omega^0(M, \mathfrak{g}_P)$, $\phi \in \Gamma(\Xi)$ in degree 1 and $\psi \in \Gamma(\Xi)$ in degree 2, we have

$$\begin{aligned} J(\xi, \phi, \psi) &= \llbracket \xi, \llbracket \phi, \psi \rrbracket \rrbracket + \llbracket \phi, \llbracket \psi, \xi \rrbracket \rrbracket + \llbracket \psi, \llbracket \xi, \phi \rrbracket \rrbracket \\ &= -(\llbracket \xi, * \rho^*(\phi\psi^*) \rrbracket - * \rho^*(\phi(\rho(\xi)\psi)^*) + * \rho^*(\psi(\rho(\xi)\phi)^*)) \\ &= - * \rho^*(\llbracket \rho(\xi), \phi\psi^* \rrbracket + \phi\psi^*\rho(\xi) - \rho(\xi)\phi\psi^*) = 0. \end{aligned}$$

- For $\xi \in \Omega^0(M, \mathfrak{g}_P)$, $a \in \Omega^1(M, \mathfrak{g}_P)$, and $\phi \in \Gamma(\Xi)$ in degree 1, we have

$$\begin{aligned} J(\xi, a, \phi) &= \llbracket \xi, \llbracket a, \phi \rrbracket \rrbracket + \llbracket a, \llbracket \phi, \xi \rrbracket \rrbracket - \llbracket \phi, \llbracket \xi, a \rrbracket \rrbracket \\ &= -\rho(\xi)\bar{\gamma}(a)\phi + \bar{\gamma}(a)\rho(\xi)\phi + \bar{\gamma}(\llbracket \xi, a \rrbracket)\phi = 0. \end{aligned}$$

- For $a \in \Omega^1(M, \mathfrak{g}_P)$ and $\phi, \psi \in \Gamma(\Xi)$ in degree 1, we have

$$\begin{aligned} J(a, \phi, \psi) &= -\llbracket a, \llbracket \phi, \psi \rrbracket \rrbracket - \llbracket \phi, \llbracket \psi, a \rrbracket \rrbracket - \llbracket \psi, \llbracket a, \phi \rrbracket \rrbracket \\ &= 2[a \wedge \mu(\phi, \psi)] + * \rho^*((\bar{\gamma}(a)\psi)\phi^*) + * \rho^*((\bar{\gamma}(a)\phi)\psi^*) = 0 \end{aligned}$$

by Proposition B.1.

Step 2. $(L^\bullet, \delta_c^\bullet)$ is a DGA.

We need to show that $\delta_c \circ \delta_c = 0$. Using Proposition B.1, we compute that

$$\begin{aligned} \delta_c^1 \circ \delta_c^0(\xi) &= \begin{pmatrix} \mathcal{D}_A \rho(\xi)\Phi - \bar{\gamma}(d_A \xi)\Phi \\ 2\mu(\Phi, \rho(\xi)\Phi) + d_A d_A \xi \end{pmatrix} \\ &= \begin{pmatrix} \rho(\xi)\mathcal{D}_A \Phi \\ [F_A - \mu(\Phi), \xi] \end{pmatrix} = 0, \end{aligned}$$

and, using Proposition B.4 and Proposition B.1, we compute that

$$\begin{aligned} \delta_c^2 \circ \delta_c^1(\phi, a) &= - * \rho^*((\mathcal{D}_A \phi)\Phi^*) - * \rho^*((\bar{\gamma}(a)\Phi)\Phi^*) - 2d_A \mu(\Phi, \phi) + d_A d_A a \\ &= \rho^*((\mathcal{D}_A \Phi)\phi) + [(F_A - \mu(\Phi)) \wedge a] = 0. \end{aligned}$$

Step 3. $(L^\bullet, \llbracket \cdot, \cdot \rrbracket, \delta_c^\bullet)$ is a DGLA.

We need to verify that δ_ζ^\bullet satisfies the graded Leibniz rule with respect to $[[\cdot, \cdot]]$, that is for every two homogeneous elements $x, y \in L^\bullet$ we need to show that

$$D(x, y) = \delta[[x, y]] - [[\delta x, y]] - (-1)^{\deg x} [[x, \delta y]]$$

vanishes.

For degree reasons, $D(x, y) = 0$ unless $\deg x + \deg y \leq 2$; hence, only the following eight cases remain:

- For $\xi, \eta \in \Omega^0(M, \mathfrak{g}_P)$, we have

$$D(\xi, \eta) = \begin{pmatrix} -\rho([\xi, \eta])\Phi \\ d_A[\xi, \eta] \end{pmatrix} - \left[\begin{pmatrix} -\rho(\xi)\Phi \\ d_A\xi \end{pmatrix}, \eta \right] - \left[\xi, \begin{pmatrix} -\rho(\eta)\Phi \\ d_A\eta \end{pmatrix} \right] = 0.$$

- For $\xi \in \Omega^0(M, \mathfrak{g}_P)$ and $\phi \in \Gamma(\mathfrak{S})$ in degree 1, we have

$$\begin{aligned} D(\xi, \phi) &= \begin{pmatrix} -\mathcal{D}_A\rho(\xi)\phi \\ -2\mu(\Phi, \rho(\xi)\phi) \end{pmatrix} - \left[\begin{pmatrix} -\rho(\xi)\Phi \\ d_A\xi \end{pmatrix}, \phi \right] - \left[\xi, \begin{pmatrix} -\mathcal{D}_A\phi \\ -2\mu(\Phi, \phi) \end{pmatrix} \right] \\ &= \begin{pmatrix} -\mathcal{D}_A\rho(\xi)\phi + \bar{\gamma}(d_A\xi)\phi + \rho(\xi)\mathcal{D}_A\phi \\ -2\mu(\Phi, \rho(\xi)\phi) - 2\mu(\rho(\xi)\Phi, \phi) + 2[\xi, \mu(\Phi, \phi)] \end{pmatrix} = 0 \end{aligned}$$

by Proposition B.1.

- For $\xi \in \Omega^0(M, \mathfrak{g}_P)$ and $a \in \Omega^1(M, \mathfrak{g}_P)$, we have

$$\begin{aligned} D(\xi, a) &= \begin{pmatrix} -\bar{\gamma}([\xi, a])\Phi \\ d_A[\xi, a] \end{pmatrix} - \left[\begin{pmatrix} -\rho(\xi)\Phi \\ d_A\xi \end{pmatrix}, a \right] - \left[\xi, \begin{pmatrix} -\bar{\gamma}(a)\Phi \\ d_Aa \end{pmatrix} \right] \\ &= \begin{pmatrix} -\bar{\gamma}([\xi, a])\Phi - \bar{\gamma}(a)\rho(\xi)\Phi + \rho(\xi)\bar{\gamma}(a)\Phi \\ d_A[\xi, a] - [d_A\xi \wedge a] - [\xi, d_Aa] \end{pmatrix} = 0. \end{aligned}$$

- For $\xi \in \Omega^0(M, \mathfrak{g}_P)$ and $\phi \in \Gamma(\mathfrak{S})$ in degree 2, we have

$$\begin{aligned} D(\xi, \phi) &= *\rho^*(\rho(\xi)\phi\Phi^*) - [[\rho(\xi)\Phi, \phi]] - [[d_A\xi, \phi]] - [[\xi, *\rho^*(\phi\Phi^*)]] \\ &= *\rho^*(\rho(\xi)\phi\Phi^*) - *\rho^*(\phi\Phi^*\rho(\xi)) - [\xi, *\rho^*(\phi\Phi^*)] = 0. \end{aligned}$$

- For $\xi \in \Omega^0(M, \mathfrak{g}_P)$ and $b \in \Omega^2(M, \mathfrak{g}_P)$, we have

$$D(\xi, b) = d_A[\xi, b] - [d_A\xi, b] - [\xi, d_Ab] = 0.$$

- For $\phi, \psi \in \Gamma(\mathfrak{S})$ in degree 1, we have

$$\begin{aligned} D(\phi, \psi) &= -2d_A\mu(\phi, \psi) - \left[\begin{pmatrix} -\mathcal{D}_A\phi \\ -2\mu(\Phi, \phi) \end{pmatrix}, \psi \right] + \left[\phi, \begin{pmatrix} -\mathcal{D}_A\psi \\ -2\mu(\Phi, \psi) \end{pmatrix} \right] \\ &= -2d_A\mu(\phi, \psi) - *\rho^*((\mathcal{D}_A\phi)\psi^*) - *\rho^*((\mathcal{D}_A\psi)\phi^*) = 0 \end{aligned}$$

by Proposition B.4.

- For $a \in \Omega^1(M, \mathfrak{g}_P)$ and $\phi \in \Gamma(\mathfrak{S})$ in degree 1, we have

$$\begin{aligned} D(a, \phi) &= * \rho^* ((\bar{\gamma}(a)\phi)\Phi^*) - \left[\left(\begin{array}{c} -\bar{\gamma}(a)\Phi \\ d_A a \end{array} \right), \phi \right] + \left[a, \left(\begin{array}{c} -\mathcal{D}_A \phi \\ -2\mu(\Phi, \phi) \end{array} \right) \right] \\ &= - * \rho^* ((\bar{\gamma}(a)\phi)\Phi^*) - * \rho^* (\bar{\gamma}(a)\Phi)\phi^* - 2[a \wedge \mu(\Phi, \phi)] = 0 \end{aligned}$$

by Proposition B.1.

- For $a, b \in \Omega^1(M, \mathfrak{g}_P)$, we have

$$D(a, b) = \left(\begin{array}{c} -\bar{\gamma}[a \wedge b]\Phi \\ d_A[a \wedge b] \end{array} \right) - \left[\left(\begin{array}{c} \bar{\gamma}(a)\Phi \\ d_A a \end{array} \right), b \right] + \left[a, \left(\begin{array}{c} \bar{\gamma}(b)\Phi \\ d_A b \end{array} \right) \right] = 0.$$

Step 4. For every $\hat{c} := (a, \phi) \in L^1$, $(A + a, \Phi + \phi)$ solves (2.14) if and only if $\delta_c \hat{c} + \frac{1}{2} \llbracket \hat{c}, \hat{c} \rrbracket = 0$.

For $\hat{c} := (a, \phi) \in L^1$, we have

$$\delta_c \hat{c} + \frac{1}{2} \llbracket \hat{c}, \hat{c} \rrbracket = \left(\begin{array}{c} -\mathcal{D}_A \phi - \bar{\gamma}(a)\Phi - \bar{\gamma}(a)\phi \\ -2\mu(\Phi, \phi) - \mu(\phi, \phi) + d_A a + \frac{1}{2}[a \wedge a] \end{array} \right),$$

which vanishes if and only if $(A + a, \Phi + a)$ solves (2.14). □

References

- [AHDM78] M. F. Atiyah, N. J. Hitchin, V. G. Drinfeld, and Yu. I. Manin. *Construction of instantons. Physics Letters. A* 65.3 (1978), pp. 185–187. DOI: 10.1016/0375-9601(78)90141-X. MR: 598562. Zbl: 0424.14004 (cit. on p. 38).
- [DK90] S. K. Donaldson and P. B. Kronheimer. *The geometry of four-manifolds*. Oxford Mathematical Monographs. Oxford Science Publications. New York, 1990, pp. x+440. MR: MR1079726. Zbl: 0904.57001 (cit. on pp. 6, 14, 16, 29).
- [DS11] S. K. Donaldson and E. P. Segal. *Gauge theory in higher dimensions, II. Surveys in differential geometry. Volume XVI. Geometry of special holonomy and related topics*. Vol. 16. 2011, pp. 1–41. arXiv: 0902.3239. MR: 2893675. Zbl: 1256.53038 (cit. on pp. 2, 38).
- [DS97] A. Dancer and A. Swann. *The geometry of singular quaternionic Kähler quotients. Internat. J. Math.* 8.5 (1997), pp. 595–610. DOI: 10.1142/S0129167X97000317. MR: 1468352. Zbl: 0886.53037 (cit. on p. 20).
- [DW18] A. Doan and T. Walpuski. *On the existence of harmonic Z_2 spinors. Journal of Differential Geometry* (2018). arXiv: 1710.06781. to appear (cit. on pp. 2, 7).

- [FL98] P. M. N. Feehan and T. G. Leness. *PU(2) monopoles and relations between four-manifold invariants*. *Topology and its Applications* 88.1-2 (1998). Symplectic, contact and low-dimensional topology (Athens, GA, 1996), pp. 111–145. DOI: 10.1016/S0166-8641(97)00201-0. MR: 1634566. Zbl: 0931.58012 (cit. on p. 37).
- [GW13] S. Guo and J. Wu. *Bifurcation theory of functional differential equations*. Vol. 184. Applied Mathematical Sciences. 2013, pp. x+289. DOI: 10.1007/978-1-4614-6992-6. MR: 3098815. Zbl: 1316.34003 (cit. on p. 16).
- [Hayo8] A. Haydys. *Nonlinear Dirac operator and quaternionic analysis*. *Communications in Mathematical Physics* 281.1 (2008), pp. 251–261. DOI: 10.1007/s00220-008-0445-1. arXiv: 0706.0389. MR: MR2403610. Zbl: 1230.30034 (cit. on p. 1).
- [Hay12] A. Haydys. *Gauge theory, calibrated geometry and harmonic spinors*. *Journal of the London Mathematical Society* 86.2 (2012), pp. 482–498. DOI: 10.1112/jlms/jds008. arXiv: 0902.3738. MR: 2980921. Zbl: 1256.81080 (cit. on pp. 5, 19, 21).
- [Hay14] A. Haydys. *Dirac operators in gauge theory. New ideas in low-dimensional topology, to appear*. 2014. arXiv: 1303.2971v2. MR: 3381325. Zbl: 1327.57028 (cit. on pp. 1, 4).
- [Hay17] A. Haydys. *G_2 instantons and the Seiberg–Witten monopoles*. 2017. arXiv: 1703.06329 (cit. on p. 38).
- [HKLR87] N. J. Hitchin, A. Karlhede, U. Lindström, and M. Roček. *Hyper-Kähler metrics and supersymmetry*. *Communications in Mathematical Physics* 108.4 (1987), pp. 535–589. DOI: 10.1007/BF01214418. MR: 877637. Zbl: 0632.53073 (cit. on pp. 3, 19).
- [HW15] A. Haydys and T. Walpuski. *A compactness theorem for the Seiberg–Witten equation with multiple spinors in dimension three*. *Geometric and Functional Analysis* 25.6 (2015), pp. 1799–1821. DOI: 10.1007/s00039-015-0346-3. arXiv: 1406.5683. MR: 3432158. Zbl: 1334.53039 (cit. on pp. 1, 6, 7).
- [KM07] P. B. Kronheimer and T. S. Mrowka. *Monopoles and three-manifolds*. Vol. 10. New Mathematical Monographs. Cambridge, 2007, pp. xii+796. DOI: 10.1017/CB09780511543111. MR: 2388043. Zbl: 1158.57002 (cit. on p. 6).
- [Nak16] H. Nakajima. *Towards a mathematical definition of Coulomb branches of 3-dimensional $\mathcal{N} = 4$ gauge theories, I*. *Advances in Theoretical and Mathematical Physics* 20.3 (2016), pp. 595–669. DOI: 10.4310/ATMP.2016.v20.n3.a4. arXiv: 1503.03676. MR: 3565863. Zbl: 06659459 (cit. on p. 1).
- [Nak99] H. Nakajima. *Lectures on Hilbert schemes of points on surfaces*. Vol. 18. University Lecture Series. Providence, RI, 1999, pp. xii+132. MR: 1711344. Zbl: 0949.14001 (cit. on p. 38).
- [Pido4] V. Ya. Pidstrigach. *Hyper-Kähler manifolds and Seiberg–Witten equations*. English. *Algebraic geometry. Methods, relations, and applications. Collected papers. Dedicated to the memory of Andrei Nikolaevich Tyurin*. 2004, pp. 249–262. MR: 2101297. Zbl: 1101.53026 (cit. on p. 1).

- [PT95] V. Ya. Pidstrigach and A.N. Tyurin. *Localisation of Donaldson invariants along the Seiberg–Witten classes*. 1995. arXiv: dg-ga/9507004 (cit. on p. 37).
- [Sal13] D.A. Salamon. *The three-dimensional Fueter equation and divergence-free frames*. *Abhandlungen aus dem Mathematischen Seminar der Universität Hamburg* 83.1 (2013), pp. 1–28. DOI: 10.1007/s12188-013-0075-1. MR: 3055820. Zbl: 1280.53073 (cit. on p. 1).
- [Str16] G. Strang. *Introduction to linear algebra. 5th edition*. English. 5th edition. 2016, pp. x + 574. Zbl: 1351.15002 (cit. on p. 32).
- [Tau13a] C. H. Taubes. *PSL(2; C) connections on 3-manifolds with L^2 bounds on curvature*. *Cambridge Journal of Mathematics* 1.2 (2013), pp. 239–397. DOI: 10.4310/CJM.2013.v1.n2.a2. arXiv: 1205.0514. MR: 3272050. Zbl: 1296.53051 (cit. on pp. 1, 7).
- [Tau13b] C. H. Taubes. *Compactness theorems for SL(2; C) generalizations of the 4-dimensional anti-self dual equations*. 2013. arXiv: 1307.6447 (cit. on p. 1).
- [Tau16] C. H. Taubes. *On the behavior of sequences of solutions to U(1) Seiberg–Witten systems in dimension 4*. 2016. arXiv: 1610.07163 (cit. on p. 1).
- [Tau99] C. H. Taubes. *Nonlinear generalizations of a 3-manifold’s Dirac operator*. *Trends in mathematical physics (Knoxville, TN, 1998)*. Vol. 13. AMS/IP Studies in Advanced Mathematics. Providence, RI, 1999, pp. 475–486. MR: 1708781. Zbl: 1049.58504 (cit. on p. 1).
- [Tel00] A. Teleman. *Moduli spaces of PU(2)–monopoles*. *Asian Journal of Mathematics* 4.2 (2000), pp. 391–435. DOI: 10.4310/AJM.2000.v4.n2.a10. MR: 1797591. Zbl: 0982.58008 (cit. on p. 37).
- [Wal17a] T. Walpuski. *G_2 –instantons, associative submanifolds, and Fueter sections*. *Communications in Analysis and Geometry* 25.4 (2017), pp. 847–893. DOI: 10.4310/CAG.2017.v25.n4.a4. arXiv: 1205.5350. MR: 3731643. Zbl: 06823232 (cit. on pp. 2, 38).
- [Wal17b] T. Walpuski. *A compactness theorem for Fueter sections*. *Commentarii Mathematici Helvetici* 92.4 (2017), pp. 751–776. arXiv: 1507.03258. MR: 3623252. Zbl: 1383.58009 (cit. on p. 7).