

Castelnuovo’s bound and rigidity in almost complex geometry

Aleksander Doan

Thomas Walpuski

2018-09-12

Abstract

This article is concerned with the question of whether an energy bound implies a genus bound for pseudo-holomorphic curves in almost complex manifolds. After reviewing what is known in dimensions other than 6, we establish a new result in this direction in dimension 6; in particular, for symplectic Calabi–Yau 3–folds. The proof relies on compactness and regularity theorems for pseudo-holomorphic currents.

1 Introduction

In 1889, Castelnuovo [Cas89] found a sharp upper bound for the genus of an irreducible degree d curve in \mathbf{P}^n ; see [ACGH85, Chapter III Section 2] for a proof in modern language. A corollary of this result is that for every projective variety there is an upper bound for the genus of an irreducible curve representing a given homology class. Our starting point is the question:

Are there analogues of Castelnuovo’s bound in almost complex geometry?

For plane curves Castelnuovo’s bound reduces to the degree-genus formula. The latter is a consequence of the adjunction formula, which generalizes to an inequality for almost complex 4–manifolds [MS12, Theorem 2.6.4]. The adjunction inequality directly implies the following well-known genus bound.

Proposition 1.1. *Suppose that (M, J) is an almost complex 4–manifold. If there exists a simple J –holomorphic map $u: \Sigma \rightarrow M$ representing $A \in H_2(M)$, then the genus $g(\Sigma)$ satisfies*

$$(1.2) \quad g(\Sigma) \leq \frac{1}{2} (A \cdot A - \langle c_1(M, J), A \rangle) + 1.$$

The following consequence of Gromov’s h-principle shows that in higher dimensions there cannot be a genus bound which holds for all almost complex structures.

Proposition 1.3 (cf. Li [Li05, Corollary 2.13]). *Let (M, ω) be a symplectic manifold of dimension $2n \geq 6$. For every $A \in H_2(M)$ with $\langle [\omega], A \rangle > 0$ and every $g \in \mathbf{N}$ there is an almost complex structure J compatible with ω and an embedded J -holomorphic curve C satisfying*

$$g(C) \geq g.$$

There are, however, genus bounds for *generic* almost complex structures. Here is a simple example, which follows easily from the index formula for J -holomorphic maps.

Proposition 1.4. *Let M be a manifold of dimension $2n$. Denote by \mathcal{F} the space of almost complex structures of class C^2 on M . There is a residual¹ subset $\mathcal{F}_\bullet \subset \mathcal{F}$ such that for every $J \in \mathcal{F}_\bullet$ the following holds: if there exists a simple J -holomorphic map $u: \Sigma \rightarrow M$ representing $A \in H_2(M)$, then*

$$(1.5) \quad \begin{cases} \langle c_1(M, J), A \rangle \geq 0 & \text{if } n = 3 \text{ and} \\ g(\Sigma) \leq \frac{\langle c_1(M, J), A \rangle}{n-3} + 1 & \text{if } n \geq 3. \end{cases}$$

Moreover, if M carries a symplectic form ω , then the same holds with \mathcal{F} replaced by the space $\mathcal{F}(\omega)$ of almost complex structures of class C^2 compatible with ω .

We give proofs of Proposition 1.3 and Proposition 1.4 in Appendix A.

The preceding discussion leaves open the case of almost complex manifolds of dimension six and homology classes satisfying $\langle c_1(M, J), A \rangle \geq 0$. We focus on the case

$$\langle c_1(M, J), A \rangle = 0,$$

that is: on classes for which the corresponding moduli space of J -holomorphic maps has expected dimension zero. This includes all homology classes in symplectic Calabi–Yau 3–folds. Our motivation for considering this case comes from our project to construct a symplectic analogue of the Pandharipande–Thomas invariants of projective Calabi–Yau 3–folds [DW17, Section 7]. Another motivation comes from the Gopakumar–Vafa conjecture. Bryan and Pandharipande [BP01] defined the Gopakumar–Vafa BPS invariants $n_A^g(M)$ of a symplectic Calabi–Yau 3–fold M in terms of its Gromov–Witten partition function. They conjectured that the BPS invariants $n_A^g(M)$ are integers and vanish for all but finitely many g [BP01, Conjecture 1.2]. The integrality conjecture has been proved by Ionel and Parker [IP18]. The finiteness conjecture remains open and is closely related to the question about the existence of genus bounds for symplectic Calabi–Yau 3–folds.

Motivated by Gromov–Witten theory, Bryan and Pandharipande introduced the notion of k -rigidity for almost complex structures; see Definition 2.10. Eftekhary [Eft16] proved that 4-rigidity is a generic property; see Theorem 2.13. Conjecturally, a generic almost complex structure is *super-rigid*, that is: ∞ -rigid. The main result of this article shows that k -rigidity implies a Castelnuovo bound.

¹Let X be a topological space. A subset $A \subset X$ is called **residual** if it is the intersection of countably many dense open subsets. Recall that a residual subset of a complete metric space is dense.

Theorem 1.6. *Let $k \in \mathbb{N} \cup \{\infty\}$. Let (M, J, g) be a compact almost Hermitian 6 -manifold with a k -rigid almost complex structure J . Suppose $A \in H_2(M)$ satisfies $\langle c_1(M, J), A \rangle = 0$ and has divisibility at most k . Given any $\Lambda > 0$, there are only finitely many simple J -holomorphic maps representing A and with energy at most Λ .*

Remark 1.7. Theorem 1.6 immediately implies a Castelnuovo bound for every fixed k -rigid almost complex structure J . Unlike (1.5), however, this bound may depend on J .

If J is tamed by a symplectic form ω , then imposing an upper bound for the energy is superfluous since the energy of any J -holomorphic map representing A is $\langle [\omega], A \rangle$.

Corollary 1.8. *Let (M, ω) be a compact symplectic Calabi–Yau 3 -fold. Suppose J is a super-rigid almost complex structure compatible with ω . Then for every $A \in H_2(M)$ there are only finitely many simple J -holomorphic maps representing A .*

In the situation of Theorem 1.6, Gromov’s compactness theorem [Gro85; PW93; Ye94; Hum97] shows that there are only finitely many J -holomorphic maps representing A from Riemann surfaces of fixed genus. It is thus not of much use for proving Theorem 1.6. Instead, we use the following compactness result for J -holomorphic cycles, that is: formal sums of J -holomorphic curves, with respect to geometric convergence; see Definition 4.1 and Definition 4.2.

Lemma 1.9. *Let M be a manifold and let $(J_n, g_n)_{n \in \mathbb{N}}$ be a sequence of almost Hermitian structures converging to an almost Hermitian structure (J, g) . Let $K \subset M$ be a compact subset and let $\Lambda > 0$. For each $n \in \mathbb{N}$ let C_n be a J_n -holomorphic cycle with support contained in K and of mass at most Λ . Then a subsequence of $(C_n)_{n \in \mathbb{N}}$ geometrically converges to a J -holomorphic cycle C .*

In dimension 4, this result was proved by Taubes [Tau96a]. The proof in higher dimensions relies on results in geometric measure theory; in particular, the recent work of De Lellis, Spadaro, and Spolaor [DSS17b; DSS18; DSS17a; DSS15] on the regularity of semi-calibrated currents.

Acknowledgements We thank Aleksey Zinger for insightful discussions, Tristan Rivière for answering our questions regarding [RT09], Costante Bellettini for pointing us towards the work of De Lellis, Spadaro, and Spolaor, and Simon Donaldson for reminding us of Gromov’s h-principle for symplectic immersions.

This material is based upon work supported by an Alfred P. Sloan fellowship, the National Science Foundation under Grant No. 1754967, and the Simons Collaboration Grant on “Special Holonomy in Geometry, Analysis and Physics”.

2 k -rigidity of J -holomorphic maps

We begin by briefly recalling the notion of k -rigidity as defined by Eftekhary. For a more detailed discussion we refer the reader to [Eft16, Section 2; Wen16, Section 2.1; DW18, Section 7]. Throughout, let (M, J, g) be an almost Hermitian $2n$ -manifold.

Definition 2.1. A J -holomorphic map $u: (\Sigma, j) \rightarrow (M, J)$ is a pair consisting of a closed, connected Riemann surface (Σ, j) and a smooth map $u: \Sigma \rightarrow M$ satisfying the non-linear Cauchy–Riemann equation

$$(2.2) \quad \bar{\partial}_J(u, j) := \frac{1}{2}(du + J(u) \circ du \circ j) = 0.$$

Definition 2.3. Let $u: (\Sigma, j) \rightarrow (M, J)$ be a J -holomorphic map. Let $\phi \in \text{Diff}(\Sigma)$ be a diffeomorphism. The **reparametrization** of u by ϕ is the J -holomorphic map $u \circ \phi^{-1}: (\Sigma, \phi_*j) \rightarrow (M, J)$.

Definition 2.4. Let $u: (\Sigma, j) \rightarrow (M, J)$ be a J -holomorphic map and let $\pi: (\tilde{\Sigma}, \tilde{j}) \rightarrow (\Sigma, j)$ be a holomorphic map of degree $\deg(\pi) \geq 2$. The composition $u \circ \pi: (\tilde{\Sigma}, \tilde{j}) \rightarrow (M, J)$ is said to be a **multiple cover** of u . A J -holomorphic map is **simple** if it is not constant and not a multiple cover.

Rigidity and k -rigidity are conditions on the infinitesimal deformation theory of J -holomorphic curves up to reparametrization. We will have to briefly review parts of this theory.

The index of a J -holomorphic map $u: (\Sigma, j) \rightarrow (M, J)$ is defined as

$$(2.5) \quad \text{index}(u) := (n - 3)\chi(\Sigma) + 2\langle [\Sigma], u^*c_1(M, J) \rangle.$$

This is the Fredholm index of the gauge-fixed linearization of (2.2). The restriction of this linearization to $\Gamma(u^*TM)$ is given by

$$(2.6) \quad \xi \mapsto \frac{1}{2}(\nabla\xi + J \circ (\nabla\xi) \circ j + (\nabla_\xi J) \circ du \circ j).$$

Here ∇ denotes any torsion-free connection on TM and also the induced connection on u^*TM . Since (u, j) is a J -holomorphic map, the right-hand side of (2.6) does not depend on the choice of ∇ ; see [MS12, Proposition 3.1.1].

Let $u: (\Sigma, j) \rightarrow (M, J)$ be a non-constant J -holomorphic map. There exists a unique complex subbundle

$$Tu \subset u^*TM$$

of rank one containing $du(T\Sigma)$; see [IS99, Section 1.3; Wen10, Section 3.3; DW18, Appendix A]. The generalized normal bundle of u is defined as

$$Nu := u^*TM/Tu.$$

If u is an immersion, then Nu is the usual normal bundle. If $\tilde{u} = u \circ \pi$ is a multiple cover of an immersion, then $N\tilde{u} = \pi^*Nu$. Define the normal Cauchy–Riemann operator

$$(2.7) \quad \mathfrak{d}_{u,J}^N: \Gamma(Nu) \rightarrow \Omega^{0,1}(Nu)$$

by the formula (2.6). The non-zero elements of the kernel of $\mathfrak{d}_{u,J}^N$ correspond to infinitesimal deformations of u which deform the image $u(\Sigma)$.

Definition 2.8. A non-constant J -holomorphic map u is **rigid** if $\ker \mathfrak{d}_{u,J}^N = 0$.

A multiple cover \tilde{u} of u may fail to be rigid, even if u itself is rigid.

Definition 2.9. Let $k \in \mathbb{N} \cup \{\infty\}$. A simple J -holomorphic map $u: (\Sigma, j) \rightarrow (M, J)$ is called **k -rigid** if it is rigid and all of its multiple covers of degree at most k are rigid.

It follows from [IS99, Lemma 1.5.1; Wen10, Theorem 3] that $\dim \ker \mathfrak{d}_{u,J}^N \geq \text{index}(u)$. Consequently, a k -rigid J -holomorphic map must have $\text{index}(u) \leq 0$.

Definition 2.10. Let $k \in \mathbb{N} \cup \{\infty\}$. An almost complex structure J is called **k -rigid** if the following hold:

1. Every simple J -holomorphic map of index zero is k -rigid.
2. Every simple J -holomorphic map has non-negative index.
3. Every simple J -holomorphic map of index zero is an embedding, and every two simple J -holomorphic maps of index zero either have disjoint images or are related by a reparametrization.

Remark 2.11. In dimension four, one should weaken (3) and require only that every simple J -holomorphic map of index zero is an immersion with transverse self-intersections, and that two such maps are either transverse to one another or are related by reparametrization. However, we will only be concerned with dimension (at least) six.

Definition 2.12. Let $k \in \mathbb{N} \cup \{\infty\}$. Let (M, ω) be a symplectic manifold. Denote by $\mathcal{F}(\omega)$ the space of almost complex structures on M compatible with ω . Denote by $\mathcal{R}_k(\omega)$ the subset of those almost complex structures $J \in \mathcal{F}(\omega)$ which are k -rigid.

Theorem 2.13 (Eftekhary [Eft16, Theorem 1.2]). *Let (M, ω) be a symplectic manifold. If $\dim M \geq 6$, then $\mathcal{R}_4(\omega) \subset \mathcal{F}(\omega)$ is a residual subset.*

Conjecture 2.14 (Bryan and Pandharipande [BP01, p. 290]). *Let (M, ω) be a symplectic manifold. If $\dim M \geq 6$, then $\mathcal{R}_\infty(\omega) \subset \mathcal{F}(\omega)$ is a residual subset.*

Wendl [Wen16] has made remarkable progress towards proving this conjecture. His work shows that Conjecture 2.14 holds provided generic real Cauchy Riemann operators satisfy an analytic condition known as Petri's condition; see also [DW18].

3 Real Cauchy–Riemann operators and almost complex structures

We will show that associated with every real Cauchy–Riemann operator defined on a vector bundle there is a natural almost complex structure on the total space of that bundle. This construction is inspired by [Tau96b, p. 825–826]; similar material can be found in [Wen16, Appendix B].

Definition 3.1. Let (Σ, j) be a Riemann surface. Let $\pi : E \rightarrow \Sigma$ be a Hermitian vector bundle over Σ . A first order linear differential operator $\mathfrak{d} : \Gamma(E) \rightarrow \Omega^{0,1}(\Sigma, E)$ is called a **real Cauchy–Riemann operator** if

$$(3.2) \quad \mathfrak{d}(fs) = (\bar{\partial}f)s + f\mathfrak{d}s$$

for all $f \in C^\infty(M, \mathbf{R})$. The **anti-linear part** of \mathfrak{d} is defined as

$$\mathfrak{n} = \mathfrak{n}_\mathfrak{d} := \frac{1}{2}(\mathfrak{d} + J\mathfrak{d}J) \in \Gamma(\text{Hom}(E, \overline{\text{Hom}}_{\mathbf{C}}(T\Sigma, E))).$$

Every real Cauchy–Riemann operator can be written as

$$\mathfrak{d} = \bar{\partial}_\nabla + \mathfrak{n}$$

where $\bar{\partial}_\nabla := \nabla^{0,1}$ is the Dolbeault operator associated with a Hermitian connection ∇ on E . Denote by $H_\nabla \subset TE$ the horizontal distribution of ∇ . It induces an isomorphism

$$(3.3) \quad TE = H_\nabla \oplus \pi^*E \cong \pi^*T\Sigma \oplus \pi^*E.$$

Definition 3.4. The **complex structure** J_∇ on E associated with ∇ is defined by pulling back the standard complex structure $j \oplus i$ on $\pi^*T\Sigma \oplus \pi^*E$ by the isomorphism (3.3).

It is well-known that a section $s \in \Gamma(E)$ satisfies $\bar{\partial}_\nabla s = 0$ if and only if the map $s : \Sigma \rightarrow E$ is J_∇ -holomorphic. The following proposition extends this to real Cauchy–Riemann operators.

Definition 3.5. Let $\mathfrak{d} = \bar{\partial}_\nabla + \mathfrak{n}$ be a real Cauchy–Riemann operator. Define $L_\mathfrak{n} : TE \rightarrow TE$ by

$$L_\mathfrak{n} = -2\mathfrak{n}(v)j\pi_*$$

at $v \in E$. The **almost complex structure** $J_\mathfrak{d}$ on E associated with \mathfrak{d} is defined by

$$J_\mathfrak{d} := J_\nabla + L_\mathfrak{n}.$$

Proposition 3.6. For every real Cauchy–Riemann operator $\mathfrak{d} : \Gamma(E) \rightarrow \Omega^{0,1}(E)$ the following hold:

1. $J_\mathfrak{d}$ is an almost complex structure.
2. The projection $\pi : E \rightarrow \Sigma$ is holomorphic with respect to $J_\mathfrak{d}$.
3. For every $x \in \Sigma$ the fiber $E_x = \pi^{-1}(x)$ is a $J_\mathfrak{d}$ -holomorphic submanifold of E .
4. A section $s \in \Gamma(E)$ satisfies $\mathfrak{d}s = 0$ if and only if $s : \Sigma \rightarrow E$ is a $J_\mathfrak{d}$ -holomorphic map.
5. There exists a symplectic form ω on the unit disc bundle $B_1(\Sigma) \subset N\Sigma$ which tames $J_\mathfrak{d}$.

Proof. With respect to (3.3) we have

$$(3.7) \quad J_{\mathfrak{b}} = \begin{pmatrix} j & 0 \\ -2\mathfrak{n}(v)j & i \end{pmatrix}$$

at $v \in E$. Since $\mathfrak{n}(v)$ is anti-linear,

$$\mathfrak{n}(v)j^2 + i\mathfrak{n}(v)j = 0.$$

Therefore,

$$J_{\mathfrak{b}}^2 = -\text{id};$$

that is, (1) holds.

Both (2) and (3) immediately follow from (3.7).

We prove (4). Let $s: \Sigma \rightarrow E$ be a section. The projection of ds to the first factor of (3.3) is $\pi_* \circ ds = \text{id}_{T\Sigma}$ and thus j -linear. The projection of $ds: T\Sigma \rightarrow s^*TE$ to the second factor is its covariant derivative $\nabla s: T\Sigma \rightarrow s^*E$. Therefore, the J_{∇} -antilinear part of ds is

$$\frac{1}{2}(ds + J_{\nabla} \circ ds \circ j) = (\nabla s)^{0,1} = \bar{\partial}_{\nabla} s.$$

The $J_{\mathfrak{b}}$ -antilinear part of ds is

$$\begin{aligned} \frac{1}{2}(ds + J_{\mathfrak{b}} \circ ds \circ j) &= \frac{1}{2}(ds + J_{\nabla} \circ ds \circ j + L_{\mathfrak{n}} \circ ds \circ j) \\ &= \bar{\partial}_{\nabla} s + L_{\mathfrak{n}} \circ ds \circ j \\ &= \bar{\partial}_{\nabla} s + \mathfrak{n}_{\mathfrak{b}} s = \mathfrak{d}s. \end{aligned}$$

Therefore, $ds: T\Sigma \rightarrow TE$ is $J_{\mathfrak{b}}$ -linear if and only if $\mathfrak{d}s = 0$.

The proof of (5) is standard; see, e.g., [Wen16, Lemma B.2]. Nevertheless, we include it here for completeness. Let ω_{Σ} be an area form on Σ . Let ω_E be any closed 2-form on $B_1(\Sigma)$ which is positive when restricted to the fibers of E ; that is, for all vertical tangent vectors v_E

$$(3.8) \quad \omega_E(v_E, J_{\nabla} v_E) \gtrsim |v_E|^2.$$

Such a form can be constructed by choosing local unitary trivializations of $E|_{U_i} \cong U_i \times \mathbb{C}^r$, denoting by λ_i the corresponding Liouville 1-forms on \mathbb{C}^r vanishing at zero, and setting

$$\omega_E = \text{d} \left(\sum_i \chi_i \circ \pi \cdot \lambda_i \right)$$

for a partition of unity (χ_i) . This form satisfies (3.8) on E . It remains to show that for $\tau \gg 1$ the closed 2-form $\omega = \tau\omega_{\Sigma} + \omega_E$ tames J_u on $B_1(\Sigma)$. For a tangent vector w to E at a point $(x, v) \in B_1(\Sigma)$ denote by w_H and w_E its horizontal and vertical parts in the decomposition (3.3). We have

$$\begin{aligned} \omega(w, J_{\mathfrak{b}} w) &= (\tau\omega_{\Sigma} + \omega_E)(w, (J_{\nabla} + L_{\mathfrak{n}})w) \\ &= \tau\omega_{\Sigma}(w_H, jw_H) + \omega_E(w_E, J_{\nabla} w_E) + \omega_E(w_E, L_{\mathfrak{n}} w_H). \end{aligned}$$

From $|L_n(v)| \lesssim |v| < 1$ it follows that

$$|\omega_E(w_E, L_n w_H)| \lesssim |w_E| |w_H|.$$

Since

$$\tau \omega_\Sigma(w_H, j w_H) + \omega_E(w_E, J \nabla w_E) \gtrsim \tau |w_H|^2 + |v_E|,$$

it follows that ω tames J_u provided $\tau \gg 1$. \square

The next two propositions are concerned with the following situation. Let $u: (\Sigma, j) \rightarrow (M, J)$ be a J -holomorphic embedding. Denote by $Nu \rightarrow \Sigma$ its normal bundle and by $\mathfrak{d}_{u,J}^N$ the normal Cauchy–Riemann introduced in (2.7). Write

$$(3.9) \quad J_u := J_{\mathfrak{d}_{u,J}^N}$$

for the almost complex structure on the total space of Nu associated with $\mathfrak{d}_{u,J}^N$.

Proposition 3.10. *For every $\lambda > 0$ define $\sigma_\lambda: Nu \rightarrow Nu$ by*

$$\sigma_\lambda(v) := \lambda v.$$

If $U \subset Nu$ is an open neighborhood of the zero section in Nu such that the exponential map $\exp: U \rightarrow M$ is an embedding, then

$$\sigma_\lambda^* \exp^* J \rightarrow J_u \quad \text{as } \lambda \rightarrow 0.$$

Proof. Denote by ∇ the connection on $Nu \rightarrow \Sigma$ induced by the Levi–Civita connection on M . Throughout this proof, we identify

$$TU = \pi^* T\Sigma \oplus \pi^* Nu$$

as in (3.3). The two almost complex structures J_∇ and $\exp^* J$ on $U \subset Nu$ agree along the zero section. The Taylor expansion of $\exp^* J$ is of the form

$$(3.11) \quad \exp^* J(x, v) = J_\nabla(x, 0) + \nabla_v J(x, 0) + O(|v|^2).$$

Set

$$L(x, v) := \nabla_n J(x, 0).$$

We write L as the matrix

$$L(x, v) = \begin{pmatrix} L_{11}(x, v) & L_{12}(x, v) \\ L_{21}(x, v) & L_{22}(x, v) \end{pmatrix}.$$

Here each L_{ij} is linear in v . The derivative $d\sigma_\lambda$ is given by

$$d\sigma_\lambda = \begin{pmatrix} \text{id} & \\ & \lambda \end{pmatrix}.$$

Therefore,

$$\begin{aligned} (\sigma_\lambda)^*L(x, v) &= \begin{pmatrix} \text{id} & \\ & \lambda^{-1} \end{pmatrix} \begin{pmatrix} L_{11}(x, \lambda v) & L_{12}(x, \lambda v) \\ L_{21}(x, \lambda v) & L_{22}(x, \lambda v) \end{pmatrix} \begin{pmatrix} \text{id} & \\ & \lambda \end{pmatrix} \\ &= \begin{pmatrix} \lambda L_{11}(x, v) & \lambda^2 L_{12}(x, v) \\ L_{21}(x, v) & \lambda L_{22}(x, v) \end{pmatrix}. \end{aligned}$$

As λ tend to zero, all but the bottom left entry tend to zero.

By construction, $\sigma_\lambda^*J_\nabla = J_\nabla$. As λ tends to zero, the rescalings of terms of second order and higher in (3.11) tend to zero. It remains to identify the term L_{21} . By definition,

$$L_{21}(x, v) = \pi_{Nu} \circ \nabla_v J(x, 0) \circ \pi_*.$$

Comparing (2.6), Definition 3.1, and Definition 3.5, we see that $L_{21} = L_u$. This finishes the proof. \square

Proposition 3.12. *If $\tilde{u}: (\tilde{\Sigma}, \tilde{j}) \rightarrow (Nu, J_u)$ is a simple J_u -holomorphic map whose image is not contained in the zero section, then:*

1. *the map $\varphi: (\tilde{\Sigma}, \tilde{j}) \rightarrow (\Sigma, j)$ given by $\varphi := \pi \circ \tilde{u}$ is non-constant and holomorphic, and*
2. *the J -holomorphic map $u \circ \varphi: (\tilde{\Sigma}, \tilde{j}) \rightarrow (M, J)$ is not rigid; in particular, the J -holomorphic map $u: (\Sigma, j) \rightarrow (M, J)$ is not k -rigid for $k = \deg(\varphi)$.*

Proof. By Proposition 3.6 (2), $\pi: Nu \rightarrow \Sigma$ is J_δ -holomorphic. Therefore, φ is holomorphic. The map φ is constant if and only if the image of \tilde{u} is contained in a fiber of π . This is impossible, because then \tilde{u} would be constant. This proves (1).

The normal bundle of the J -holomorphic map $u \circ \varphi$ is $N_{u \circ \varphi} = \varphi^*Nu$. The corresponding normal Cauchy–Riemann operator is

$$(3.13) \quad \mathfrak{d}_{u \circ \varphi, J}^N = \varphi^* \mathfrak{d}_{u, J}^N.$$

Since \tilde{u} takes values in Nu , for every $x \in \tilde{\Sigma}$ we have

$$\tilde{u}(x) \in Nu_{\pi(u(x))} = Nu_{\varphi(x)} = (\varphi^*Nu)_x.$$

This gives rise to the section $s \in \Gamma(\varphi^*Nu)$ defined by

$$s(x) := \tilde{u}(x) \in (\varphi^*Nu)_x.$$

This section is not the zero section, because the image of \tilde{u} is not contained in the zero section. The upcoming discussion will show that

$$\mathfrak{d}_{u \circ \varphi, J}^N s = 0.$$

In light of Proposition 3.6 (4), this will imply (2).

Denote by Z the finite set of critical values of φ . Set $\tilde{Z} := \varphi^{-1}(Z)$. The restriction $\tilde{u}: \tilde{\Sigma} \setminus \tilde{Z} \rightarrow Nu$ is a J_u -holomorphic embedding and $\varphi: \tilde{\Sigma} \setminus \tilde{Z} \rightarrow \Sigma \setminus Z$ is an unbranched holomorphic covering map. Hence, every $x \in \tilde{\Sigma} \setminus \tilde{Z}$ has an open neighborhood U such that $\tilde{u}|_U$ is an embedding and $\varphi|_U$ is biholomorphic. Therefore, $\pi|_{\tilde{u}(U)}$ maps $\tilde{u}(U)$ holomorphically to $\varphi(U)$. It follows that $\tilde{u}(U) \subset Nu$ is the graph of a J_u -holomorphic section $f: \varphi(U) \rightarrow Nu|_{\varphi(U)}$. By construction and by Proposition 3.6,

$$s|_U = \varphi^* f \quad \text{and} \quad \mathfrak{d}_{u,J}^N f = 0.$$

The relation (3.13) shows that $\mathfrak{d}_{u \circ \varphi, J}^N s = 0$ holds on U . Since $x \in \tilde{\Sigma} \setminus \tilde{Z}$ was arbitrary, it holds on all of $\tilde{\Sigma} \setminus \tilde{Z}$. In fact, since s is smooth, it holds on all of $\tilde{\Sigma}$. \square

4 J -holomorphic cycles and geometric convergence

In this section we introduce the notions of J -holomorphic cycles and geometric convergence. We then compare these with the notions of closed J -holomorphic integral currents and weak convergence. This comparison, combined with a classical compactness result in geometric measure theory, implies Lemma 1.9. Throughout, let (M, J, g) be an almost Hermitian manifold. Denote by

$$\sigma := g(J, \cdot)$$

the corresponding Hermitian form.

Definition 4.1. A J -holomorphic curve is a subset of M which is the image of a simple J -holomorphic map $u: (\Sigma, j) \rightarrow (M, J)$. A J -holomorphic cycle C is a formal linear combination

$$C = \sum_{i=1}^I m_i C_i$$

of J -holomorphic curves C_1, \dots, C_I with coefficients $m_1, \dots, m_I \in \mathbf{N}$. The **homology class** represented by C is

$$[C] := \sum_{i=1}^I m_i (u_i)_* [\Sigma_i].$$

The **support** of C is the subset

$$\text{supp}(C) := \bigcup_{i=1}^I C_i.$$

The **current** associated with C is defined by

$$\delta_C(\alpha) := \sum_{i=1}^I m_i \int_{\Sigma_i} u_i^* \alpha \quad \text{for} \quad \alpha \in \Omega_c^2(M).$$

The **mass** of C is

$$\mathbf{M}(C) := \sum_{i=1}^I m_i \text{area}(C_i) = \delta_C(\sigma).$$

We say that C is **smooth** if the J -holomorphic curves C_1, \dots, C_I are embedded and pairwise disjoint.

Definition 4.2 (Taubes [Tau98, Definition 3.1]). Let M be a manifold and let $(J_n, g_n)_{n \in \mathbf{N}}$ be a sequence of almost Hermitian structures converging to an almost Hermitian structure (J, g) . For every $n \in \mathbf{N}$ let C_n be a J_n -holomorphic cycle. We say that $(C_n)_{n \in \mathbf{N}}$ **geometrically converges** to a J -holomorphic cycle C if:

1. $(\delta_{C_n})_{n \in \mathbf{N}}$ weakly converges to δ_C ; that is:

$$\lim_{n \rightarrow \infty} \delta_{C_n}(\alpha) = \delta_C(\alpha) \quad \text{for all } \alpha \in \Omega_c^2(M)$$

and

2. $(\text{supp}(C_n))_{n \in \mathbf{N}}$ converges to $\text{supp}(C)$ in the Hausdorff distance; that is:

$$(4.3) \quad \lim_{n \rightarrow \infty} d_H(\text{supp}(C), \text{supp}(C_n)) \rightarrow 0.$$

Recall that the Hausdorff distance between two closed sets X and Y is defined by

$$d_H(X, Y) := \max \left\{ \sup_{x \in X} d(x, Y), \sup_{y \in Y} d(y, X) \right\}.$$

The following results compare J -holomorphic cycles and geometric convergence with closed integral currents on M which are calibrated by σ and weak convergence. We refer the reader to the lecture notes [Lan05] for the required background on geometric measure theory.

Proposition 4.4. *If δ_C is a closed integral current which is calibrated by σ , then there exist a J -holomorphic map $u: (\Sigma, j) \rightarrow (M, J)$, with a possibly disconnected domain Σ , and a locally constant function $k: \Sigma \rightarrow \mathbf{N}$ such that*

$$\delta_C(\alpha) = \int_{\Sigma} k \cdot u^* \alpha \quad \text{for all } \alpha \in \Omega_c^2(M).$$

In particular, there exists a J -holomorphic cycle C whose associated current is δ_C .

Proposition 4.5. *In the situation of Definition 4.2, if condition (1) holds and there exists a compact subset containing $\text{supp}(C_n)$ for every $n \in \mathbf{N}$, then condition (2) holds as well.*

We prove the second result first, since it is more elementary.

Proof of Proposition 4.5. This is contained in the proof of [Tau98, Proposition 3.3] and also a well-known fact in geometric measure theory. Let us explain the proof nevertheless. The salient point is the monotonicity formula for J -holomorphic curves; see, e.g., [PW93, Corollary 3.2; Tau98, Lemma 3.4]. It states that there are constants $c, r_0 > 0$ such that for every J -holomorphic curve C , every $x \in C$, and every $r \in [0, r_0]$

$$(4.6) \quad \mathbf{M}(\delta_C|_{B_r(x)}) = \delta_C|_{B_r(x)}(\sigma) \geq cr^2.$$

Moreover, it follows from the proof of the monotonicity formula that these constants can be chosen such that (4.6) holds for every almost Hermitian structures in the sequence $(J_n, g_n)_{n \in \mathbb{N}}$ as well as the limit (J, g) .

Condition (4.3) is equivalent to

$$(4.7) \quad \lim_{n \rightarrow \infty} \sup \{d(x, \text{supp}(C)) : x \in \text{supp}(C_n)\} = 0 \quad \text{and}$$

$$(4.8) \quad \lim_{n \rightarrow \infty} \sup \{d(x, \text{supp}(C_n)) : x \in \text{supp}(C)\} = 0.$$

If (4.7) fails, then after passing to a subsequence there exists an $\varepsilon > 0$ and a sequence of points (x_n) with

$$x_n \in \text{supp}(C_n) \quad \text{but} \quad d(x_n, \text{supp}(C)) \geq \varepsilon.$$

After passing to a further subsequence (x_n) converges to a limit $x \in M$ with $d(x, \text{supp}(C)) \geq \varepsilon$. Fix $0 < r \leq \min\{\varepsilon/2, r_0\}$. Let $\chi \in C^\infty(M, [0, 1])$ be supported in $B_{2r}(x)$ and equal to one in $B_r(x)$. By (4.6), for $n \gg 1$

$$c_0 r^2 \leq \mathbf{M}(\delta_{C_n}|_{B_r(x)}) \leq \delta_{C_n}(\chi\sigma).$$

This contradicts the weak convergence condition (1), because

$$\delta_C(\chi\sigma) = 0.$$

If (4.8) fails, then a slight variation of this argument derives another contradiction to (1). \square

Proof of Proposition 4.4. For symplectic 4-manifolds this was proved by Taubes [Tau96a, Proposition 6.1]. Taubes' argument and the work of Rivière and Tian [RT09] establish the result for general symplectic manifolds. The extension to almost Hermitian manifolds relies on the work of De Lellis, Spadaro, and Spolaor [DSS18; DSS17b; DSS17a; DSS15]. Their main result [DSS18, Theorem 0.2] implies that the singular set of δ_C is finite (since it is discrete and since δ_C is closed and of finite mass and, thus, has compact support). We need not just their main result but also the following intermediate result.

Definition 4.9. Given $k \in \mathbb{N}$, set

$$\tilde{D}^k := \{(z, w) \in \mathbb{C}^2 : z = w^k \text{ and } |z| < 1\}.$$

We consider $\tilde{D}^k \setminus \{0\}$ as oriented smooth manifold such that the map $(z, w) \mapsto z$ is an orientation-preserving local diffeomorphism. We equip it with the pull-back of the flat metric.

Definition 4.10. Let $k \in \mathbf{N}$ and $\alpha \in (0, 1)$. Let $f: \tilde{D}^k \rightarrow \mathbf{R}^{2n-2}$ be a continuous injective map which is of class $C^{3,\alpha}$ on $\tilde{D}^k \setminus \{0\}$ and satisfies $|df| \lesssim |z|^\alpha$. Define $\underline{f}: \tilde{D}^k \rightarrow \mathbf{R}^{2n}$ by

$$\underline{f}(z, w) := (z, f(w)).$$

Let $U \subset \mathbf{R}^{2n}$ be an open subset and let $\phi: U \rightarrow M$ be a chart. The **graph** of f with respect to ϕ is the integral current $G_{f,\phi}$ defined by

$$G_{f,\phi}(\alpha) := \int_{\tilde{D}^k \setminus \{0\}} \underline{f}^* \phi^* \alpha \quad \text{for } \alpha \in \Omega_c^2(M).$$

Lemma 4.11 (De Lellis, Spadaro, and Spolaor [DSS17a, Section 1]). *For every $x \in \text{supp}(\delta_C)$ there are: a neighborhood U of x , finite collections of maps f_1, \dots, f_m and charts ϕ_1, \dots, ϕ_m as in Definition 4.10, and weights $\ell_1, \dots, \ell_m \in \mathbf{N}$ such that*

$$\delta_C|_U = \sum_{i=1}^m \ell_i G_{f_i, \phi_i}.$$

Denote by $\mathring{\Sigma}$ the regular part of $\text{supp}(\delta_C)$. Since δ_C is calibrated, the tangent spaces to $\mathring{\Sigma}$ are J -invariant. Therefore, $\mathring{\Sigma}$ canonically is a Riemann surface. As mentioned earlier, the singular set $\text{sing}(\delta_C) := \text{supp}(C) \setminus \mathring{\Sigma}$ is finite. Lemma 4.11 shows that every $x \in \text{sing}(\delta_C)$ has a neighborhood U such that

$$\mathring{\Sigma} \cap U \cong \mathbf{C}^* \sqcup \dots \sqcup \mathbf{C}^*.$$

Thus, $\mathring{\Sigma}$ can be compactified to a Riemann surface Σ by adding finitely many points.

The Riemann surface Σ comes with a continuous map $u: \Sigma \rightarrow M$. Its restriction to $\mathring{\Sigma}$ is smooth and J -holomorphic. It follows from elliptic regularity that u is, in fact, smooth and J -holomorphic on all of Σ . The above discussion shows that

$$\delta_C(\alpha) = \int_{\Sigma} k \cdot u^* \alpha$$

for some locally constant function $k: \Sigma \rightarrow \mathbf{N}$. □

Proof of Lemma 1.9. The sequence of closed integral currents $(\delta_{C_n})_{n \in \mathbf{N}}$ has uniformly bounded mass. Therefore, there exists a subsequence which weakly converges to a closed integral current δ_C calibrated by σ ; see, e.g., [Fed69, Theorem 4.2.17; Sim83, Theorem 27.3; Lan05, Theorem 3.7]. By Proposition 4.4, δ_C is the current associated with a J -holomorphic cycle C . By Proposition 4.5, the sequence of pseudo-holomorphic cycles (C_n) geometrically converges to C . □

5 Proof of Theorem 1.6

Suppose that J is k -rigid and that $A \in H_2(M)$ satisfies $\langle c_1(M, J), A \rangle = 0$ and its divisibility is at most k . If the conclusion of the theorem fails, then there are infinitely many *distinct* J -holomorphic curves $C_n \subset M$ representing A and of energy at most Λ . By Lemma 1.9, after passing to a subsequence, the sequence (C_n) converges geometrically to a J -holomorphic cycle

$$C_\infty = \sum_{i=1}^I m_i C_\infty^i.$$

Proposition 5.1. *C_∞ is connected, smooth, and its multiplicity is at most the divisibility of A .*

Proof. By Definition 4.2 (1), $[C_\infty] = [A]$. Let $u_i : \Sigma_i \rightarrow M$ be a simple J -holomorphic map whose image is C_∞^i . The index formula (2.5) yields

$$\sum_{i=1}^I m_i \text{index}(u_i) = \sum_{i=1}^I 2m_i \langle c_1(M, J), [C_\infty^i] \rangle = 2 \langle c_1(M, J), [C_\infty] \rangle = 0.$$

Since J is k -rigid, by Definition 2.10 (2), there are no J -holomorphic curves of negative index. Thus, we have $\text{index}(u_i) \geq 0$ for every $i \in \{1, \dots, I\}$ and the above computation shows that

$$\text{index}(u_1) = \dots = \text{index}(u_I) = 0.$$

Therefore, by Definition 2.10 (3), the J -holomorphic curves $C_\infty^1, \dots, C_\infty^I$ are embedded and pairwise disjoint. This proves that C_∞ is smooth.

To see that C_∞ is connected, observe that if C_∞ were disconnected, then Definition 4.2 (2) would imply that C_n is disconnected for $n \gg 1$. However, C_n is a J -holomorphic curve and thus connected by definition.

Since $A = m_1 [C_\infty^1]$, it follows that m_1 is at most the divisibility of A . \square

In the following, we rescale the sequence (C_n) and extract a further limit \tilde{C}_∞ . The properties of \tilde{C}_∞ will give a contradiction to J being k -rigid.

Henceforth, we denote by C_∞^1 the J -holomorphic curve underlying the J -holomorphic cycle C_∞ . Since the curves C_n are all distinct, we can assume that they are all distinct from C_∞^1 . We can also assume that every C_n is contained in a sufficiently small tubular neighborhood of C_∞^1 . By slight abuse of notation, we regard C_n as an $\exp^* J$ -holomorphic curve in the normal bundle NC_∞^1 and C_∞^1 as the zero section in NC_∞^1 .

For every $\lambda > 0$ let σ_λ be as in Proposition 3.10. Choose (λ_n) such that such that the sets

$$\tilde{C}_n := \sigma_{\lambda_n}^{-1}(C_n)$$

satisfy

$$(5.2) \quad d_H(\tilde{C}_n, C_\infty^1) = 1/2.$$

Set

$$J_n := \sigma_{\lambda_n}^* \exp^* J.$$

By construction, the \tilde{C}_n are J_n -holomorphic. By Proposition 3.10, the sequence (J_n) converges to the almost complex structure J_u associated with the J -holomorphic map $u: C_\infty^1 \hookrightarrow M$. The sequence (\tilde{C}_n) is contained in the compact disc bundle $\bar{B}_{1/2}(C_\infty^1) \subset NC_\infty^1$. By Proposition 3.6 (5), J_u is tamed by a symplectic form ω on $B_1(C_\infty^1)$. Consequently, for $n \gg 1$ the almost complex structure J_n is tamed by ω as well. Define a Riemannian metric g on $B_1(C_\infty^1)$ by

$$g := \frac{1}{2}(\omega(J_u \cdot, \cdot) + \omega(\cdot, J_u \cdot)).$$

The analogously defined metrics g_n are Hermitian with respect to J_n and converge to g . By the energy identity [MS12, Lemma 2.2.1],

$$\lim_{n \rightarrow \infty} \mathbf{M}(\tilde{C}_n) = \lim_{n \rightarrow \infty} \delta_{\tilde{C}_n}(\omega) = \delta_{C_\infty^1}(\omega) < \infty.$$

Therefore, the mass of \tilde{C}_n with respect to g_n (and thus also g) can be bounded independent of n .

By Lemma 1.9, a subsequence of (\tilde{C}_n) geometrically converges to a J -holomorphic cycle

$$\tilde{C}_\infty = \sum_{i=1}^I \tilde{m}_i \tilde{C}_\infty^i.$$

Condition (5.2) guarantees that $\text{supp}(\tilde{C}_\infty) \neq C_\infty^1$. Therefore, without loss of generality, \tilde{C}_∞^1 is not contained in the zero section. Since

$$[\tilde{C}_\infty] = \sum_{i=1}^I \tilde{m}_i [\tilde{C}_\infty^i] = m_1 [C_\infty^1] = A,$$

\tilde{m}_1 is bounded by the divisibility of A ; hence: $\tilde{m}_1 \leq k$. Proposition 3.12 applies and the map φ defined there has degree \tilde{m}_1 . This contradicts J being k -rigid. \square

A Proofs of Proposition 1.3 and Proposition 1.4

Proof of Proposition 1.3. By a classic theorem of Thom, there is a closed, connected, oriented surface Σ and an embedding $u_0: \Sigma \rightarrow M$ with $u_*[\Sigma] = A$. After adding sufficiently many small 1-handles, we can assume that $g(\Sigma) \geq n$. Gromov's h -principle [Gro86, Section 3.4.2 Theorem (A)] implies that u_0 is C^0 -close to an embedding $u: \Sigma \rightarrow M$ with $u^* \omega > 0$.

Denote the image of u by C . The restriction of TM to C is the direct sum of the symplectic subbundle TC and its symplectic complement, which can be identified with NC . The space of complex structures on \mathbf{R}^{2n} compatible with a fixed non-degenerate 2-form is contractible. This implies that: (1) both TC and NC admit almost complex structures compatible with ω ; hence, $TM|_C$ admits an almost complex structure compatible with ω , and (2) any such almost complex structure on $TM|_C$ can be extended to an almost complex structure on TM compatible with ω . \square

Remark A.1. If (M, ω) is a symplectic 4-manifold, then every $A \in H_2(M)$ with $\langle [\omega], A \rangle > 0$ is represented by an *immersed* symplectic surface C with transverse double points. Such a surface is J -holomorphic for an almost complex structure compatible with ω if and only if all of its self-intersections are positive.

Proof of Proposition 1.4. Denote by \mathcal{M} the component of the universal moduli space of simple pseudo-holomorphic maps from a Riemann surface of genus g to M . This is a separable Banach manifold and the projection map $\pi: \mathcal{M} \rightarrow \mathcal{F}$ is a Fredholm map of class C^1 and index

$$(n-3)(2-2g) + 2\langle c_1(M, J), A \rangle;$$

see, e.g., [Wen10, Theorem 0; IP18, Proposition 5.1]. If (1.5) is violated, then this index is negative. The result thus follows from the Sard–Smale theorem [Sma65]. \square

References

- [ACGH85] E. Arbarello, M. Cornalba, P. A. Griffiths, and J. Harris. *Geometry of algebraic curves. Volume I*. Vol. 267. Grundlehren der Mathematischen Wissenschaften. 1985, pp. xvi+386. DOI: 10.1007/978-1-4757-5323-3. MR: 770932. Zbl: 0559.14017 (cit. on p. 1).
- [BP01] J. Bryan and R. Pandharipande. *BPS states of curves in Calabi–Yau 3-folds*. *Geometry and Topology* 5 (2001), pp. 287–318. DOI: gt.2001.5.287. MR: 1825668. Zbl: 1063.14068 (cit. on pp. 2, 5).
- [Cas89] G. Castelnuovo. *Ricerche di geometria sulle curve algebriche*. Italian. *Atti della Accademia delle Scienze di Torino* 24 (1889), pp. 346–373. Zbl: 21.0669.01 (cit. on p. 1).
- [DSS15] C. De Lellis, E. Spadaro, and L. Spolaor. *Regularity theory for 2-dimensional almost minimal currents III: blowup*. 2015. arXiv: 1508.05510 (cit. on pp. 3, 12).
- [DSS17a] C. De Lellis, E. Spadaro, and L. Spolaor. *Regularity theory for 2-dimensional almost minimal currents II: Branched center manifold*. *Annals of PDE* 3.2 (2017), Art. 18, 85. DOI: 10.1007/s40818-017-0035-7. MR: 3712561 (cit. on pp. 3, 12, 13).
- [DSS17b] C. De Lellis, E. Spadaro, and L. Spolaor. *Uniqueness of tangent cones for two-dimensional almost-minimizing currents*. *Communications on Pure and Applied Mathematics* 70.7 (2017), pp. 1402–1421. DOI: 10.1002/cpa.21690. MR: 3666570 (cit. on pp. 3, 12).
- [DSS18] C. De Lellis, E. Spadaro, and L. Spolaor. *Regularity theory for 2-dimensional almost minimal currents I: Lipschitz approximation*. *Transactions of the American Mathematical Society* 370.3 (2018), pp. 1783–1801. DOI: 10.1090/tran/6995. MR: 3739191. Zbl: 1381.49047 (cit. on pp. 3, 12).
- [DW17] A. Doan and T. Walpuski. *On counting associative submanifolds and Seiberg–Witten monopoles*. 2017. arXiv: 1712.08383 (cit. on p. 2).

- [DW18] A. Doan and T. Walpuski. *Equivariant Brill–Noether theory for elliptic operators and super-rigidity of J -holomorphic maps*. 2018 (cit. on pp. 3–5).
- [Eft16] E. Eftekhary. *On finiteness and rigidity of J -holomorphic curves in symplectic three-folds*. *Advances in Mathematics* 289 (2016), pp. 1082–1105. DOI: 10.1016/j.aim.2015.11.028. MR: 3439707. Zbl: 1331.53107 (cit. on pp. 2, 3, 5).
- [Fed69] H. Federer. *Geometric measure theory*. Die Grundlehren der mathematischen Wissenschaften, Band 153. 1969, pp. xiv+676. MR: 0257325. Zbl: 0176.00801 (cit. on p. 13).
- [Gro85] M. Gromov. *Pseudo holomorphic curves in symplectic manifolds*. *Inventiones Mathematicae* 82.2 (1985), pp. 307–347. DOI: 10.1007/BF01388806. MR: 809718. Zbl: 0592.53025 (cit. on p. 3).
- [Gro86] M. Gromov. *Partial differential relations*. Vol. 9. 1986. Zbl: 0651.53001 (cit. on p. 15).
- [Hum97] C. Hummel. *Gromov’s compactness theorem for pseudo-holomorphic curves*. Vol. 151. Progress in Mathematics. 1997, pp. viii+131. DOI: 10.1007/978-3-0348-8952-0. MR: 1451624. Zbl: 0870.53002 (cit. on p. 3).
- [IP18] E.-N. Ionel and T. H. Parker. *The Gopakumar–Vafa formula for symplectic manifolds*. *Annals of Mathematics* 187.1 (2018), pp. 1–64. DOI: 10.4007/annals.2018.187.1.1. MR: 3739228. Zbl: 06841536 (cit. on pp. 2, 16).
- [IS99] S. Ivashkovich and V. Shevchishin. *Structure of the moduli space in a neighborhood of a cusp-curve and meromorphic hulls*. *Inventiones Mathematicae* 136.3 (1999), pp. 571–602. DOI: 10.1007/s002220050319. MR: 1695206. Zbl: 0930.32017 (cit. on pp. 4, 5).
- [Lan05] U. Lang. *Introduction to Geometric Measure Theory*. 2005. URL: https://people.math.ethz.ch/~lang/diab05_notes.pdf (cit. on pp. 11, 13).
- [Lio5] T.-J. Li. *Existence of symplectic surfaces*. *Geometry and topology of manifolds. Papers from the conference held at McMaster University, Hamilton, ON, USA, May 14–18, 2004*. 2005, pp. 203–217. Zbl: 1099.57026 (cit. on p. 2).
- [MS12] D. McDuff and D. Salamon. *J -holomorphic curves and symplectic topology*. Second. Vol. 52. American Mathematical Society Colloquium Publications. 2012, pp. xiv+726. MR: 2954391. Zbl: 1272.53002 (cit. on pp. 1, 4, 15).
- [PW93] T. H. Parker and J. G. Wolfson. *Pseudo-holomorphic maps and bubble trees*. *Journal of Geometric Analysis* 3.1 (1993), pp. 63–98. DOI: 10.1007/BF02921330. MR: 1197017. Zbl: 0759.53023 (cit. on pp. 3, 12).
- [RT09] T. Rivière and G. Tian. *The singular set of 1–1 integral currents*. *Annals of Mathematics* 169.3 (2009), pp. 741–794. DOI: 10.4007/annals.2009.169.741. MR: MR2480617. Zbl: 1182.32010 (cit. on pp. 3, 12).

- [Sim83] L. Simon. *Lectures on geometric measure theory*. Vol. 3. Proceedings of the Centre for Mathematical Analysis, Australian National University. Canberra, 1983, pp. vii+272. MR: 756417 (cit. on p. 13).
- [Sma65] S. Smale. *An infinite dimensional version of Sard's theorem*. *American Journal of Mathematics* 87 (1965), pp. 861–866. DOI: 10.2307/2373250. MR: 0185604. Zbl: 0143.35301 (cit. on p. 16).
- [Tau96a] C. H. Taubes. $SW \Rightarrow Gr$: *from the Seiberg–Witten equations to pseudo-holomorphic curves*. *Journal of the American Mathematical Society* 9.3 (1996), pp. 845–918. DOI: 10.1090/S0894-0347-96-00211-1. MR: MR1362874. Zbl: 0867.53025 (cit. on pp. 3, 12).
- [Tau96b] C.H. Taubes. *Counting pseudo-holomorphic submanifolds in dimension 4*. *Journal of Differential Geometry* 44.4 (1996), pp. 818–893. MR: 1438194. Zbl: 0883.57020. URL: <http://projecteuclid.org/euclid.jdg/1214459411> (cit. on p. 5).
- [Tau98] C.H. Taubes. *The structure of pseudo-holomorphic subvarieties for a degenerate almost complex structure and symplectic form on $S^1 \times B^3$* . *Geometry and Topology* 2 (1998), pp. 221–332. MR: 1658028. Zbl: 0908.53013. URL: <https://doi.org/10.2140/gt.1998.2.221> (cit. on pp. 11, 12).
- [Wen10] C. Wendl. *Automatic transversality and orbifolds of punctured holomorphic curves in dimension four*. *Commentarii Mathematici Helvetici* 85.2 (2010), pp. 347–407. DOI: 10.4171/CMH/199. MR: 2595183. Zbl: 1207.32021 (cit. on pp. 4, 5, 16).
- [Wen16] C. Wendl. *Transversality and super-rigidity for multiply covered holomorphic curves*. 2016. arXiv: 1609.09867 (cit. on pp. 3, 5, 7).
- [Ye94] R. Ye. *Gromov's compactness theorem for pseudo holomorphic curves*. *Transactions of the American Mathematical Society* 342.2 (1994), pp. 671–694. DOI: 10.2307/2154647. MR: 1176088. Zbl: 0810.53024 (cit. on p. 3).